

**Analytic Number Theory**  
**Homework #4**

*(due Thursday, April 28, 2016)*

**Problem 1:** Let  $\chi$  be a Dirichlet character (mod  $q$ ) for some integer  $q > 1$ . Prove that  $L(1, \chi) \ll \log q$ .

**Problem 2:** Fix a prime  $p$ . Show that

$$\Gamma_0(p) := \left\{ \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL(2, \mathbb{Z}) \mid c \equiv 0 \pmod{p} \right\}$$

is a subgroup of  $SL(2, \mathbb{Z})$ .

**Problem 3:** Define

$$P(z) := \sum_{\substack{c \in \mathbb{Z} \\ d \in \mathbb{Z} \\ (c,d)=1}} \frac{e^{2\pi i \cdot \frac{az+b}{cz+d}}}{(cz+d)^{2k}}.$$

Here, for every pair of coprime integers  $c, d$  we choose integers  $a, b$  so that  $ad - bc = 1$ . Show that the above series is independent of the choice of  $a, b$ . Also show that the above series converges absolutely for  $k > 1$ .

**Problem 4:** Rewrite  $P(z)$  as a sum involving  $j(\gamma, z) = cz + d$  for  $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ . If  $k > 1$  is an integer, show that  $P(z)$  is a holomorphic modular form of weight  $2k$  for  $SL(2, \mathbb{Z})$ .

**Problem 5:** For  $s \in \mathbb{C}$  with  $\Re(s) > 8$ , let

$$L(s) := \sum_{n=1}^{\infty} a(n)n^{-s}, \quad (a(n) \in \mathbb{C} \text{ for } n = 1, 2, \dots)$$

where  $a(1) = 1$  and  $|a(n)| \ll n^7$  (for  $n = 1, 2, \dots$ ). Assume that the function  $\Phi(s) := (2\pi)^{-s}\Gamma(s)L(s)$  is an entire function which is bounded in any fixed vertical strip  $\{s \in \mathbb{C} \mid a < \Re(s) < b\}$  and satisfies the functional equation

$$\Phi(s) = \Phi(12 - s)$$

for all  $s \in \mathbb{C}$ . Using the inverse Mellin transform, prove that

$$\sum_{n=1}^{\infty} a(n)e^{2\pi inz}, \quad (z \in \mathfrak{h})$$

is the Ramanujan cusp form of weight 12.