

A NOTE ON CLOSED 3-BRAIDS*

JOAN S. BIRMAN[†]

*Department of Mathematics, Barnard College and Columbia University
2990 Broadway
New York, NY 10027, USA
jb@math.columbia.edu*

WILLIAM W. MENASCO[‡]

*Department of Mathematics, University at Buffalo
244 Mathematics Building
Buffalo, New York 14260-2900, USA
menasco@math.buffalo.edu*

Dedicated to the memory of Xiao-Song Lin

Received 8 February 2008
Revised 13 May 2008

This is a review article about knots and links of braid index 3. Its goal is to gather together, in one place, some of the tools that are special to knots and links of braid index 3, in a form that could be useful for those who have a need to calculate, and need to know precisely all the exceptional cases.

Keywords: Braids; 3-braids; flypes; transversally simple.

Mathematics Subject Classification 2000: 57M25, 57M27

Knots and links which are closed 3-braids are a very special class. Like 2-bridge knots and links, they are simple enough to admit a complete classification. At the same time they are rich enough to serve as a source of examples on which, with luck, a researcher may be able to test various conjectures. As an example, non-invertibility is a property of knots which is fairly subtle, yet we know (see Theorem 1.2 below) precisely which links of braid index 3 are and are not invertible.

*This article is an expanded version of the preprint “A note on transversal knots which are closed 3-braids”, arXiv math/0703669.

[†]Partially supported by the US National Science Foundation Grant DMS 0405586.

[‡]Partially supported by the US National Science Foundation Grant DMS 0306062.

Therefore 3-braids could be very useful for testing potential invariants that might detect non-invertibility.

The purpose of this note is to gather together, in one place, some of the tools that are special to knots and links of braid index 3, in a form that could be useful for those who have a need to calculate, and need to know precisely all the exceptional cases. It includes the work in an earlier preprint [4] which was restricted to 3-braid representatives of transversal knots. The preprint [4] was written when [26] was in preparation, and the authors of [26] asked us about low crossing examples to test their then-new invariants of transversal knots.

1. Three Basic Theorems

In this section we present the basic lemmas and theorems about links that are closed three-braids. Initially, we will use the classical presentation for the braid group B_3 :

$$\langle \sigma_1, \sigma_2; \sigma_1\sigma_2\sigma_1 = \sigma_2\sigma_1\sigma_2 \rangle \tag{1.1}$$

In the presentation (1.1) the generator σ_i is a positive interchange of strands i and $i + 1$, where $i = 1$ or 2 .

Before we can state our first theorem, we need a definition. Let \mathcal{L} be a closed 3-braid, with representative $L \in B_3$. Then:

- \mathcal{L} is said to *admit a flype* if the conjugacy class of L has a representative which is in one of the special forms illustrated in each of the four sketches in Figure 1. In particular its associated (open) braids are conjugate to:

$$L = \sigma_1^u \sigma_2^v \sigma_1^w \sigma_2^\epsilon, \text{ for some } u, v, w, \epsilon \in \mathbb{N}, \epsilon = \pm 1. \tag{1.2}$$

- A flype is *positive* or *negative*, according as ϵ is $+1$ or -1 .
- A flype is *non-degenerate* when the braids $L = \sigma_1^u \sigma_2^v \sigma_1^w \sigma_2^\epsilon$ and $L' = \sigma_1^w \sigma_2^v \sigma_1^u \sigma_2^\epsilon$ are in distinct conjugacy classes. We are interested only in non-degenerate flypes.




Fig. 1. Positive and Negative 3-braid flypes

Theorem 1.1 (The classification theorem [5]). *Let \mathcal{L} be a link type which is represented by the closure of a 3-braid L . Then one of the following holds:*

- (1) \mathcal{L} has braid index 3 and every 3-braid which represents \mathcal{L} is conjugate to L .
- (2) \mathcal{L} has braid index 3, and is represented by exactly two distinct conjugacy classes of closed 3-braids. This happens if and only if the conjugacy class of L contains a braid which admits a non-degenerate flype.
- (3) The braid L is conjugate to $\sigma_1^k \sigma_2^{\pm 1}$ for some $k \in \mathbb{N}$. These are precisely the links which are defined by closed 3-braids, but have braid index less than 3.

Remark. It is immediately clear that in order to use Theorem 1.1, the reader will need definitive tests for deciding when two 3-braids are conjugate, and when a conjugacy class admits a non-degenerate flype, and when a conjugacy class contains $\sigma_1^k \sigma_2^{\pm 1}$ for some k . These matters will be discussed in §2 below.

Let \mathcal{K} be a knot. The standard knot tables classify unoriented knots. A sharper classification assumes that we fix an orientation on both S^3 and on a representative \vec{K} . Let \overleftarrow{K} denote \vec{K} with its orientation reversed. We then define \mathcal{K} to be *invertible* if there is an orientation-preserving homeomorphism h of S^3 such that $h(\vec{K})$ is smoothly isotopic to \overleftarrow{K} . Our second classification theorem treats the question of when an oriented link of braid index 3 fails to be invertible. As was mentioned earlier, this is an interesting issue, because at this writing there is no effective test that recognizes whether an arbitrary link is or is not invertible. Using the standard presentation (1.1) for B_3 , we note that if \vec{K} is defined by a cyclic word in the standard generators, then \overleftarrow{K} will be defined by the same cyclic word, read backwards.

Theorem 1.2 (The invertibility theorem [5]). *Let \mathcal{K} be a link of braid index 3 with oriented 3-braid representative \vec{K} . Then \mathcal{K} is non-invertible if and only if \vec{K} and \overleftarrow{K} are in distinct conjugacy classes, and the class of \vec{K} does not contain a representative which admits a non-degenerate flype.*

Proof. If \vec{K} admits a non-degenerate flype, then the isotopy of S^3 which realizes the flype takes $\vec{K} \rightarrow \overleftarrow{K}$, so that \mathcal{K} is invertible. If \vec{K} does not admit a non-degenerate flype, then by Theorem 1.1 the class of \vec{K} is the only class that represents the oriented link \mathcal{K} . The assertion follows. □

A knot is said to be *transversal* if it is everywhere transverse to the standard contact structure in \mathbb{R}^3 . A *transversal knot type* is the equivalence class of a transversal knot, under isotopies which are restricted so that every intermediate representative is transversal. Our third classification theorem relates to transversal knot types that are closed 3-braids. It uses Theorem 1.1 and is related to Theorem 1.2, however in Theorem 1.3 below we are interested explicitly in the conjugacy classes which admit a non-degenerate negative flype and do not at the same time admit a positive flype, because a positive flype can be realized by a transversal isotopy. Recall that

a transversal knot type is *transversally simple* if it is determined completely by its topological link type and its Bennequin invariant.

Theorem 1.3 (The transversal classification theorem [6]). *Let \mathcal{TK} be a transversal knot type which is represented by a closed 3-braid $K_1 = \sigma_1^u \sigma_2^v \sigma_1^w \sigma_2^{-1}$. Assume that \mathcal{K}_1 does not also admit a positive flype. Then the braids K_1 and $K_2 = \sigma_1^w \sigma_2^v \sigma_1^u \sigma_2^{-1}$ close to distinct transversal knot types $\mathcal{TK}_1, \mathcal{TK}_2$ that are topologically invertible but are not transversally invertible. In particular, \mathcal{TK}_1 and \mathcal{TK}_2 are not transversally simple.*

2. Solving the Conjugacy Problem in B_3

As noted above, to use Theorem 1.1 the reader will need definitive tests for deciding (i) when two 3-braids are conjugate; (ii) when a conjugacy class is one of the exceptional ones which admits a non-degenerate flype; and (iii) when a conjugacy class contains $\sigma_1^k \sigma_2^{\pm 1}$ for some k . In this section we discuss (i).

The conjugacy problem in B_3 was solved long ago by O. Schreier [33], making use of the fact that B_3 is a free product with amalgamations of cyclic groups of orders 2 and 3. His work did not generalize to B_n , $n > 3$, where the braid groups have a more complicated structure than they do when $n = 3$.

The first solution to the conjugacy decision and conjugacy search problems for all braid groups B_n , $n \geq 3$, was discovered by F. Garside in 1965 [13]. Garside’s solution was exponential in word length and in braid index, and that fact has led to major efforts to find a polynomial-time solution to the conjugacy decision and search problems in B_n , which continue at this writing. In 1992, motivated by the work of Garside [13] and the later work of Birman and Menasco on 3-braids in [5], P.J. Xu discovered in [40] a new and fast solution to the conjugacy problem in the special case of B_3 . Her work led naturally to an algorithm to solve the shortest word problem in B_3 , and so (by special properties of links that are closed 3-braids, established in [5]) to compute the genus of a knot of braid index 3. One surprising feature of it, that was generalized later to all the groups B_n , $n \geq 2$, in [3] and thence to many Garside groups, was that it gave a ‘dual’ Garside structure on the braid groups. This fact played an important role in the isolation of *Garside groups* as an interesting class [8, 7]. In describing the basic theorems about 3-braids we choose Xu’s work over Schreier’s because it generalizes to all n and so may give insights that might otherwise be missed. We also choose the band presentation (described below) and its Garside structure over the classical presentation (1.1) and its Garside structure because it seems simpler, from a computational point of view. None of the solutions to the conjugacy problem in B_3 are particularly suited to problems (ii) and (iii) above.

2.1. The band presentation

The *band presentation* of B_3 uses a cyclically ordered set of generators which includes the generators of (1.1):

$$\langle a_1, a_2, a_3 \mid a_2 a_1 = a_3 a_2 = a_1 a_3 \rangle, \text{ where } a_1 = \sigma_1, \quad a_2 = \sigma_2, \quad a_3 = \sigma_1^{-1} \sigma_2 \sigma_1. \tag{2.1}$$

In terms of the new generators, a 3-braid L admits a flype (see (1.2)) if its conjugacy class contains $a_1^u a_2^v a_1^w a_2^{\pm 1}$ for some u, v, w, ϵ . The constraints on u, v, w, ϵ are as before. In working with the presentation (2.1) subscripts are always defined mod 3. Let $\delta = a_2 a_1 = a_3 a_2 = a_1 a_3$. The following formulas follow immediately from the defining relations in (2.1).

$$\begin{aligned} (i) \quad & a_i^{-k} = \delta^{-k} a_{i-k+2} a_{i-k+1} \cdots a_i a_{i+1} \quad \text{for all } k \geq 1, \\ (ii) \quad & a_j \delta^m = \delta^m a_{j+m} \quad \text{for all } m \in \mathbb{N}. \end{aligned} \tag{2.2}$$

They will be used repeatedly in what follows. Note that δ^3 generates the center of B_3 , because subscripts are defined mod 3.

2.2. The word problem

Xu’s solution to the word problem builds directly on the analogous ideas in the work of Garside [13]. Choose any word W in the symbols a_1, a_2, a_3 and their inverses, representing an element $\beta \in B_3$. Using (2.2)(i), all inverses of generators can be traded for powers of δ^{-1} . Using (2.2)(ii), all powers of δ may be collected, say, at the left, so that one may rewrite W in the form $\delta^k P$, where $k \in \mathbb{N}$ and P is a *positive* word. Positive words contain powers of δ if and only if two adjacent letters $a_{i+1} a_i$ appear, so after a simple search we may change $\delta^k P$ to a word of the form $\delta^q a_{r_1} a_{r_2} \cdots a_{r_s}$ where q is maximal and the positive word $a_{r_1} a_{r_2} \cdots a_{r_s}$ contains no adjacent letters of the form $a_{i+1} a_i$. Rewriting our word in syllable form $X = \delta^q a_{r_1}^{s_1} \cdots a_{r_t}^{s_t}$, we see that the subscripts must be strictly increasing. The integer q is the *power*, and the positive word is the *tail*. This solves the word problem in B_3 .

We give an example. The following set of moves bring the flype-admissible braid $W = a_1^{-2} a_2^{-3} a_1^5 a_2$, which is not in normal form, to its normal form $\delta^{-4} a_2^2 a_3 a_1^5 a_2$: Use (i) to modify $W = a_1^{-2} a_2^{-3} a_1^5 a_2$ to $(\delta^{-1} a_2)^2 (\delta^{-1} a_3)^3 a_1^5 a_2$. Now use (ii) to modify $(\delta^{-1} a_2)^2 (\delta^{-1} a_3)^3 a_1^5 a_2$ to $\delta^{-2} a_1 a_2 \delta^{-3} a_1 a_2 a_3 a_1^5 a_2$ and then to $\delta^{-5} a_1 a_2 a_1 a_2 a_3 a_1^5 a_2$. Finally, observe that $\delta = a_2 a_1$ is a subword of the tail of $\delta^{-5} a_1 a_2 a_1 a_2 a_3 a_1^5 a_2$, so we can move it to the left to obtain $\delta^{-4} a_2^2 a_3 a_1^5 a_2$. The subscripts in the tail are strictly increasing, so we are done.

2.3. The conjugacy problem for 3-braids

The conjugacy problem is close-by. The *summit set* of W , denoted $SS(W)$, is the subset of those elements in the conjugacy class of W which, when represented by words in normal form, have maximal power. It is, by definition, invariant under conjugation by powers of δ . Even more, it must be finite, for the following reasons. The defining relations in the band presentation preserve the algebraic sum of the exponents of an element in B_3 , and conjugation does too. Therefore the algebraic sum is a class invariant, and from this it follows that every time that the power is increased by conjugation, the tail goes down in letter length, until we reach the summit set. As for the summit set, it must be finite because the letter length of the

tail of a word in $SS(W)$ is finite and is an invariant. The summit set is Xu’s basic invariant of the conjugacy class of W . This was the invariant in Garside’s algorithm in [13], however in the special case of B_3 , using the band presentation, Xu was able to determine $SS(W)$ rapidly, and choose a unique symbol $\Sigma^*(W)$ with the property that two elements W, W' are conjugate if and only if $\Sigma^*(W) = \Sigma^*(W')$. Her unique symbol is defined as follows. Suppose that $X = \delta^m a_{i_1}^{l_1} a_{i_2}^{l_2} \cdots a_{i_t}^{l_t}$ is in the summit set of W . She associates to X the array $\Sigma(X) = (m, (l_1, l_2, \dots, l_t))$. The integer m is fixed (and is maximal for the conjugacy class. Her representative $\Sigma^*(W)$ is chosen so that t is minimal, and so that among all sequences with the same t it is lexicographically smallest. Thus, for example, $(5, (2, 2)) < (5, (1, 2, 1))$ and $(5, (1, 3)) < (5, (2, 2))$. She proves that $W, W' \in B_3$ are conjugate if and only if $\Sigma^*(W) = \Sigma^*(W')$. We illustrate the algorithm via an example.

Step 1: The first part of the algorithm is to find a single element in $SS(W)$. Let us suppose that we are given an element $W = \delta^q a_{\tau_1}^{s_1} \dots a_{\tau_t}^{s_t} \in B_3$, which is in normal form but may or may not be in its summit set. To test whether W is in $SS(W)$, we modify $W = \delta^q a_{\tau_1}^{s_1} \dots a_{\tau_t}^{s_t}$ by ‘twisted cyclic permutation’ of the tail of W . That is, we push the left-most syllable $a_{\tau_1}^{s_1}$ in the tail to the left, past δ^q , using (2.2) and then move it to the right end of W . This may be achieved by a special kind of conjugation that we call a *tail move* on W :

$$W = \delta^q a_{\tau_1}^{s_1} \dots a_{\tau_t}^{s_t} \rightarrow (\delta^q a_{\tau_1}^{-s_1} \delta^{-q})(W)(\delta^q a_{\tau_1}^{s_1} \delta^{-q}) = \delta^q a_{\tau_2}^{s_2} \dots a_{\tau_t}^{s_t} (\delta^q a_{\tau_1}^{s_1} \delta^{-q}) = X.$$

Xu proves that W is in its summit set if and only if, after a single tail move, the power is not increased and the syllable length is not decreased. To understand why this is so, observe that the final syllable of X is $\delta^q a_{\tau_1}^{s_1} \delta^{-q} = a_j^{s_1}$ for some $j = 1, 2, 3$ and so there are 3 possibilities, namely $j = \tau_t + 1, \tau_t$ or $\tau_t - 1 \pmod{3}$. Xu proves that if $j = \tau_t + 1$ or τ_t , then W and also X are in normal form as written and are in $SS(W)$. The first part of the algorithm ends. For, if X is modified again by another tail move, then the modified X will have the same power as X and be in normal form, because W was in normal form, so that $\tau_2 = \tau_1 + 1 \pmod{3}$, and this property is preserved by the second tail move.

If X is not in normal form, then the only possibility is that $j = \tau_t - 1$, in which case we obtain a new factor δ at the interface between the final letter in W and the initial letter in the syllable that was permuted, because $\delta = a_{\tau_t} a_{\tau_t-1}$. An example is given by $W = \delta^{-3} a_1^4 a_2 a_3^3 a_1^2 a_2$. After a tail move it changes to $X = \delta^{-3} (a_2 a_3^3 a_1^2 a_2) (a_1^4) = \delta^{-3} a_2 a_3^3 a_1^2 \delta a_1^3 = \delta^{-2} a_3 a_1^3 a_2^2 a_1^3$, which has higher power than W . But in this example X is not in normal form, because when the factor δ that appeared after the tail move was moved to the left, an additional factor δ popped up at the interface between the final two syllables. Pushing the new factor δ to the left, we obtain $Y = \delta^{-2} a_3 a_1^3 a_2 \delta a_1^2 = \delta^{-1} a_1 a_3^3 a_3 a_1^2$. The power has been increased again. Normal form has been achieved, because the subscript array is increasing.

We now test whether Y in our example is in $SS(W)$, by applying a tail move. The answer is ‘yes’. More generally, the process of repeated applications of tail moves,

and passage to normal form, continues until we find an element whose power is unchanged after a tail move. In our example that element is Y and so $Y \in \text{SS}(W)$. In a more general example, another cycling might lead to additional powers of δ , but in the most general example the following holds: If a tail move does not change the power, then we have achieved summit power. Note that the tail move may, nevertheless, increase syllable length.

Step 2: Having achieved summit power, Xu’s method for choosing an invariant of the conjugacy class is the following: If $Y = \delta^m a_{\mu_1}^{l_1} \dots a_{\mu_k}^{l_k}$ belongs to $\text{SS}(W)$, one associates to Y its *Xu symbol* $\Sigma(Y) = (m; (l_1, l_2, \dots, l_k))$. In our example, $\Sigma(Y) = (-1; (1, 3, 1, 2))$.

Note that every element in $\text{SS}(W)$ has the same summit power m , but elements need not have the same syllable length. In particular, if in step 1 it had happened that, after a tail move on W , we found ourselves in the case $j = \tau_t$, then the syllable length would go down, because in X the permuted syllable $a_j^{s_1}$ could be amalgamated with the final syllable $a_{\tau_t}^{s_t}$ of W . Among all elements $Y \in \text{SS}(W)$, her defining symbol $\Sigma^*(W)$ is the one for which the sequence of exponents (l_1, l_2, \dots, l_k) is lexicographically smallest. In particular, the syllable length is minimized among elements in the summit set. It is now easy to see that if a tail move leaves both the power and the syllable length invariant then we are almost done with step 2.

We study $\Sigma(Y) = (-1; (1, 3, 1, 2))$ in our example. We seek the unique element in $\text{SS}(W)$ for which the tail symbol is lexicographically smallest. But now observe that the candidates are simply the 4 cyclic permutations of the tail symbol, i.e. $(1,3,1,2)$, $(3,1,2,1)$, $(1,2,1,3)$, $(2,1,3,1)$. (In a more general case the tail symbol will have, say, u entries, and so it will have u cyclic permutations, all of which occur for some element in $\text{SS}(W)$.) In our example $\Sigma^*(Y) = (-1; (1, 2, 1, 3))$ is the Xu invariant of the conjugacy class. We found it without calculating the entire summit set, in particular we ignored all the elements whose syllable length was bigger than 4, and we did not have to deal with the 3 possible subscript sequences of each element that has this Xu symbol. Therefore we are done. □

3. Using Theorems 1.1, 1.2 and 1.3

Theorems 1.1 and 1.2. We need two more results before we can make Theorems 1.1 and 1.2 into results that can be used computationally. First, we need to be able to recognize when the closure of a 3-braid has braid index less than 3. Table 1 gives the Xu symbols $\Sigma^*(W)$ in the cases when the closed braid associated to W has braid index less than 3. Using it, we can decide when a closed 3-braid is a link of braid index 3, by computing its Xu-invariant and verifying whether it is or is not the symbol for a 3-braid of braid index less than 3. In the table the symbol 1^u means u syllables of length 1. For example, Table 1 tells us that a 3-braid X whose Xu-invariant is $(-5; (1^4, 2)) = (-5; (1, 1, 1, 1, 2))$ is conjugate to the 3-braid $a_1^{-5}a_2$, which closes to the type (2,5) torus knot. Therefore the closure of X has braid index 2.

Table 1. Conjugacy classes of links of braid index less than 3

W	$\Sigma^*(W)$
$a_1^k a_2, k = 1 \text{ and } 2$	$(1; (k - 1))$
$a_1^k a_2, k \geq 3$	$(-1; (k + 1))$
$a_1^k a_2^{-1}, k \geq 2$	$(-1; (1, k))$
$a_1^k a_2, k \leq -2$	$(k; (1^{-k-1}, 2))$
$a_1^k a_2^{-1}, k \leq -1,$	$(k; (1^{-k-1}))$
$a_1^{-1} a_2 \text{ and } a_1 a_2^{-1}$	$(-1; (2))$
a_2	$(0; (1))$
a_2^{-1}	$(-1; (1))$

Next, if we are given two 3-braids W, W' and find that they have distinct Xu-invariants, we cannot be sure that they determine distinct links until we determine whether they are flype-admissible. To answer this question, we need the following Lemma, which is due to K.H. Ko and S.J. Lee:

Lemma 3.1 (K.H. Ko and S.J. Lee, [21]). *(i) The flype in (1.2) is non-degenerate if and only if $|v| \geq 2$, also $u, v + \epsilon, w$ are pairwise distinct, also $u, w \neq 0, \epsilon, 2\epsilon$. (ii) The conjugacy class of a negative flype-admissible braid L contains a braid that admits a positive flype if and only if $u = 1$ or $w = 1$, or $v = 2$.*

In Table 2 we give the Xu symbols for all non-degenerate flype-admissible braids,

Table 2. Conjugacy classes admitting non-degenerate flypes

ϵ	W	$\Sigma^*(W)$, up to cyclic permutation of tail symbol
<i>pos</i>	$a_1^p a_2^q a_1^r a_2$	$(3 ; (p - 2, q - 1, r - 2))$
	$a_1^{-p} a_2^q a_1^r a_2$	$(1 - p ; (q, r - 1, 1^p))$
	$a_1^p a_2^{-q} a_1^r a_2$	$(2 - q ; (p - 1, 1^{q-1}, r - 1))$
	$a_1^p a_2^q a_1^{-r} a_2$	$(1 - r ; (p - 1, q, 1^r))$
<i>neg</i>	$a_1^p a_2^q a_1^r a_2^{-1}$	$(0 ; (p, q - 1, r))$
	$a_1^{-p} a_2^q a_1^r a_2^{-1}$	$(-p ; (q, r + 1, 1^{p-2}))$
	$a_1^p a_2^{-q} a_1^r a_2^{-1}$	$(-q - 1 ; (r + 1, p + 1, 1^{q-1}))$
	$a_1^p a_2^q a_1^{-r} a_2^{-1}$	$(-r ; (p + 1, q, 1^{r-2}))$

up to cyclic permutation of the tail symbol. There are 16 cases, because the exponents u, v, w, ϵ can either be positive or negative. To keep track of the signs, replace u, v, w by $\pm p, \pm q, \pm r$, where $p, q, r > 0$. Now observe that only 8 of the 16 cases are needed, because a given braid K admits a non-degenerate flype if and only if the Xu-symbol for K or for K^{-1} coincides with one of the 8 symbols that are given in Table 2.

We remark that the data in Table 2 is only given up to cyclic permutation of the tail symbol, because the lexicographically smallest choice depends upon the relative numerical values of p, q, r .

We now have essentially all the data needed to implement Theorems 1.1, 1.2 and 1.3, however an additional issue arises if one wishes to use any of our three

theorems on low crossing examples which admit non-degenerate flypes. For example, one might wish to test a potential invariant which was difficult to compute on knots which had large crossing number. We give, in Table 3, a list of all of the knots of braid index 3 which admit non-degenerate negative flypes, and do not also admit positive flypes, and have braid crossing number $c_b = |u| + |v| + |w| + 1 \leq 12$. Note that c_b is not the same as minimum crossing number for all possible projections of the knot associated to our closed braid (although for all but one of the examples in Table 1 the two are actually the same), rather it is crossing number for closed 3-braid representatives which are in flype position.

The knots are identified, in Table 3, using both classical notation (e.g as in [32])

Table 3. A list of all examples of non-transversally simple closed 3-braids with braid crossing number $c_b \leq 12$

$\mathcal{K}(\text{classical})$	$\mathcal{K}(\text{Dowker-Thistlethwaite})$	β	c_b	(u, v, w)	(w, v, u)
8 ₁₀	8 _{a3}	-1	8	(3,-2,2)	(2,-2,3)
10 ₄₇	10 _{a15}	+1	10	(5,-2,2)	(2,-2,5)
10 ₆₂	10 _{a41}	+1	10	(3,-2,4)	(4,-2,3)
10 ₄₈	10 _{a79}	-3	10	(3,-4,2)	(2,-4,3)
10 ₁₂₇	10 _{n16}	-9	10	(-5,-2,2)	(2,-2,-5)
10 ₁₄₃	10 _{n26}	-7	10	(3,-2,-4)	(-4,-2,3)
***	11 _{a240}	+7	12	(5,3,3)	(3,3,5)
***	12 _{a146}	+3	12	(7,-2,2)	(2,-2,7)
***	12 _{a369}	+3	12	(5,-2,4)	(4,-2,5)
***	12 _{a576}	+3	12	(3,-2,6)	(6,-2,3)
***	12 _{a835}	-5	12	(3,-6,2)	(2,-6,3)
***	12 _{a878}	-1	12	(5,-4,2)	(2,-4,5)
***	12 _{a1027}	-1	12	(3,-4,4)	(4,-4,3)
***	12 _{a1233}	+1	12	(5,-3,3)	(3,-3,5)
***	12 _{n234}	-11	12	(-7,-2,2)	(2,-2,-7)
***	12 _{n466}	-9	12	(-5,-3,3)	(3,-3,-5)
***	12 _{n467}	-7	12	(-3,-4,4)	(4,-4,-3)
***	12 _{n468}	-5	12	(-3,-3,5)	(5,-3,-3)
***	12 _{n472}	-15	12	(-3,-4,-4)	(-4,-4,-3)
***	12 _{n570}	-9	12	(-3,-5,3)	(3,-5,-3)

and the newer notation of Dowker-Thistlethwaite (used in all the newer tables). The former only exist through 10 crossings.

To assemble the data in Table 3 we began by listing all possible numbers $(\pm|u|, \pm|v|, \pm|w|)$ with $c_b = |u| + |v| + |w| + 1$, deleting those that do not satisfy the constraints of the two lemmas. Next we used Knotscape [15] to sort the survivors into classes which had the same topological knot type. For each topological knot type, some of the examples had non-minimal braid crossing number, so we deleted those. For example, $(-3, 4, 2)$ appeared as an example with $c_b = 10$, but its conjugacy class also has an 8-crossing example, $(3, -2, 2)$, so we deleted $(-3, 4, 2)$. After that, for each topological knot type we divided the representing closed braids into conjugacy classes belonging to the braids before and after the

flype. In every case it turned out that there were two distinct conjugacy classes in B_3 , as must be the case, by Theorem 1.1. Finally, we chose exactly one representative from each conjugacy class, whenever it happened that the list of survivors had more than one representative. In this regard we note that, for example, $\sigma_1^{-5}\sigma_2^3\sigma_1^{-3}\sigma_2^{-1} \approx \sigma_1^2\sigma_2^{-2}\sigma_1^{-5}\sigma_2^{-1} \approx \sigma_1^{-3}\sigma_2^{-4}\sigma_1^2\sigma_2^{-1}$. This shows that conjugate braids which admit flypes need not have the same braid crossing number c_b . In this case the first has braid crossing number 12 and the other two have crossing number 10. There are examples where all three have the same crossing number, and examples where one has minimal crossing number and the other two have higher crossing number. This issue arises when we enumerate the low-crossing number examples.

The examples in Table 3 are given by specifying, for each pair of transversal knot types, the topological knot type \mathcal{K} in the standard tables, the Bennequin number $\beta = u + v + w - 4$, the minimum closed braid crossing number $c_b = |u| + |v| + |w| + 1$, and the two closed 3-braid representatives $\mathcal{TK}_1, \mathcal{TK}_2$. The closed braids are determined by the triplets (u, v, w) and (w, v, u) , i.e. as negative flype-related braids $\sigma_1^u\sigma_2^v\sigma_1^w\sigma_2^{-1}$ and $\sigma_1^w\sigma_2^v\sigma_1^u\sigma_2^{-1}$.

4. Some Open Questions and a Survey of Known Results About Links of Braid Index 3

In this section we survey the literature and discuss some unknown properties of knots and links of braid index 3. Our choices are guided by what we know best, and by our wish to illustrate ways in which Theorems 1.1, 1.2 and 1.3 have been and might be useful.

- (1) We made inquiries, but were unable to find a computer program that implements Xu’s solution to the conjugacy problem in B_3 . If anyone is interested, it would be useful to have such a program, if it is freely available to researchers.
- (2) In our judgment, the most striking feature of closed 3-braids that has not been used in any substantial way in the literature is that we know precisely which links of braid index 3 are invertible, and which are not. In the early history of knot theory it was not known whether non-invertible knots exist. The first proof that they do exist is due to H. Trotter [37], who showed that if $|p|, |q|, |r| \geq 2$ are distinct, then the assumption that the pretzel knots $K(p, q, r)$ are invertible leads to a contradiction about the structure of their groups. His indirect proof characterizes all the work that we know which treats non-invertibility. We do not know of any computable numerical invariant that detects non-invertible knots. Therefore, if anyone has a candidate for an invariant that detects non-invertibility, three-braids are an excellent source of examples, because ‘most’ of them are non-invertible. Note that all knots that admit a flype, degenerate or not, are invertible. Theorem 1.2 and Tables 1 and 2 tell us precisely how to find all non-invertible 3-braid examples.

- (3) A knot \vec{K} is *strongly invertible* if there is a representative $\vec{K} \subset S^3$ and an involution $\iota : (S^3, \vec{K}) \rightarrow (S^3, \overleftarrow{K})$. In the course of writing this note we observed the following:

Proposition 4.1. *In the special case of 3-braids, if a knot type \vec{K} is invertible, then it is strongly invertible.*

Question: Is it possible that this is the case for arbitrary knots?

- (4) We discuss next a problem which turns out to be closely related to problem 2 above, although initially it does not seem to be. A knot is *transversal* if it is everywhere transverse to the standard contact structure in \mathbb{R}^3 . The *transversal knot type* of a transversal knot is its equivalence class under isotopies through transversal knots. Known invariants of transversal knot types are their topological knot type and their Bennequin number and (a recent development at the time of this writing) certain invariants based upon Knot Floer Homology and Khovanov Homology [28, 26, 30, 31, 39, 14]. The first examples of pairs of knots which have the same topological knot type and the same Bennequin invariant, but are transversally distinct, were discovered via 3-braids [6]. The proof in [6] is indirect, and at this writing nobody has found an invariant which distinguishes even one of those pairs, however caution is needed: Each pair is an example of a pair of knots which are topologically invertible but not transversally invertible. In view of Problem 1 above they are bound to be extremely subtle to detect.
- (5) We discuss a different set of questions about 3-braids which has received much attention and which, like Problem 2 above, seems very difficult. When the Jones and HOMFLY polynomials were discovered [16] it was not known whether they were or were not injective on topological link types, but examples taken from 3-braids [2] quickly showed that the answer is ‘no’. However, at this writing we know very little about how good a job the Jones polynomial does. It is therefore a natural question to ask whether one can arrive at some understanding of the kernel of the Jones polynomial in the special case of braid index 3. This is especially tempting because when the braid index is 3 the Jones polynomial depends on just one small calculation: the trace of a certain 2×2 matrix. Nevertheless, the prohibitively complicated formulas in [36] tell us that this is simply not a good question, and that simplifications are needed before we can hope for a start in understanding the kernel of the Jones polynomial of links that are closed 3-braids.

An example is enlightening, even though it does not have much to do with 3-braids. In the class of torus links, two integers r, s are all that is needed to distinguish one torus link from another. On the other hand, if one wishes to distinguish the Jones polynomial of one torus link from that of another, the

matter is considerably more complicated. One has the following very complicated formula [17]:

$$V_{r,s}(t) = \frac{t^{\frac{1}{2}(r-1)(s-1)}}{1-t^2} \sum_{l=0}^n \binom{n}{l} t^{\frac{r}{n}(n-l)(1+\frac{s}{n}l)} (t^{(n-l)\frac{s}{n}} - t^{l+\frac{s}{n}}), \tag{4.1}$$

where $n = \gcd(r, s)$. It isn't even obvious that it gives a polynomial (although of course it does). The reader will notice that, knowing the general form of (4.1), $V_{r,s}(t)$ is completely determined by the two integers r and s .

- (6) Concluding this note, we point the reader, in no particular order, to some of the published work that has come to our attention about links of braid index 3:
- (a) K. Murasugi's monograph [24] was pioneering, in the sense that it was the first study that isolated knots and links of braid index 3 as a special class worthy of study. It is filled with detailed information and the author's tireless, error-free and very useful calculations. Murasugi used Schreier's solution to the conjugacy problem [33], but carried things a step further by partitioning Schreier's conjugacy classes into groups which were later realized to coincide with pseudo-Anosov, finite order and reducible braids. In this sense it was prescient, because it foresaw the Nielsen-Thurston trichotomy that has played such a major role in understanding mapping class groups of surfaces, in the special case of 3-braids. This seemed so striking that many people tried to carry generalize Murasugi's work to B_n , but alas the generalizations did not seem to work. One may also find, in [24], an example of the flypes which play such an important role in Theorems 1.1, 1.2 and 1.3.
 - (b) Two algorithmic solutions to the shortest word problem for 3-braids appeared in the literature essentially simultaneously. They were due to P.J. Xu [40] and M. Berger [1], and they used the band and classical presentations respectively. A year later R.D. Keever [19] gave yet another solution, duplicating both of the preceding. (His paper appeared in the same journal as Xu's paper, a year later, and neither Keever nor anyone else seems to have realized the duplication!). Of these three, only Xu's paper does more, as it not only also gives the pleasant solution to the conjugacy problem that we discussed in §2.3 but also proves that the shortest word problem implies an algorithm for computing link genus. Very recently Usman Ali [38] posted a paper on the arXiv that chooses a unique representative in $SS(W)$ algorithmically when W is a 3-braid, but using the classical presentation rather than the band presentation of Xu. All these results are of interest because it has been proved by Paterson and Razborov [29] that the shortest word problem in B_n , using the classical presentation, is NP complete.
 - (c) In a separate paper [41] Xu uses her earlier work to compute the growth functions for the positive braid semigroup. Neither [1] nor [19] consider that problem.

- (d) In a different direction S. Orevkov [27] discovered an algorithm for deciding whether links of braid index 3 are quasi-positive, a question which is of interest in the study of the topology of plane real algebraic curves.
- (e) In [25] Y. Ni used knot Floer homology to determine which links of braid index 3 are fibered. He introduced there the interesting concept of ‘almost fibered’. The question of when 3-braids are fibered was also studied by A. Stoimenow in [34].
- (f) M.T. Lozano and J. Przytycki [22] classified horizontal incompressible surfaces in the complement of a link of braid index 3. Using a different set of techniques, E. Finkelstein obtained related results in [12].
- (g) D. Erle has computed the signature of links of braid index 3 in [9].
- (h) A. Stoimenow gives a classification of alternating links of braid index 3 in [35].
- (i) W. Menasco and X. Zhang prove, in [23], that knots which are closed 3-braids have Property P.
- (j) In [20] E. Kin uses 3-braids to study a problem in dynamics.

Acknowledgment

We thank Sang-Jin Lee, Lenny Ng, Peter Ozsvath and Dylan Thurston for useful comments on the precursor of this manuscript, the preprint [4]. We thank Usman Ali for pointing out errors in Table 1.

References

- [1] M. Berger, Minimum crossing numbers for 3-braids, *J. Phys. A* **27** (1994) 6205–6213.
- [2] J. S. Birman, On the Jones polynomial of closed 3-braids, *Invent. Math.* **81** (1985) 287–294.
- [3] J.S. Birman, K.H. Ko and S.J. Lee, A new approach to the word and conjugacy problems in the braid groups, *Adv. Math.* **139** (1998) 322–353.
- [4] J.S. Birman and W.W.Menasco, A note on transversal knots which are closed 3-braids, preprint, arXiv:math/0703669.
- [5] J. S. Birman and W. W. Menasco, Studying links via closed braids III: Classifying links which are closed 3-braids, *Pacific J. of Math.* **161** (1993) 25–113.
- [6] J. S. Birman and W. W. Menasco, Stabilization in the braid groups II: Transversal simplicity of knots, *Geometry and Topology* **10** (2006) 1425–1452.
- [7] P. Dehornoy, Groupes de Garside, *Ann. Sci. cole Norm. Sup. (4)* **35** (2002) 267–306.
- [8] P. Dehornoy and L. Paris, Gaussian groups and Garside groups, two generalisations of Artin groups, *Proc. London Math. Soc. (3)* **79** (1999) 569–604.
- [9] D. Erle, Calculation of the signature of a 3-braid link, *Kobe J. Math.* **16** (1999) 161–175.
- [10] J. Etnyre and K. Honda, Cabling and transverse simplicity, *Annals of Math.(2)* **162** (2005) 1305–1333.
- [11] E. El-Rifai and H. Morton, Algorithms for positive braids, *Quart. J. Math. Oxford Ser. (2)* **45** (1994) 479–497.

- [12] E. Finkelstein, Closed incompressible surfaces in closed braid complements, *J. Knot Theory Ramifications* **7** (1998) 335–379.
- [13] F. A. Garside, The braid group and other groups, *Quart. J. Math. Oxford Ser. (2)* **20** (1969) 235–254.
- [14] S. Harvey, K. Kawamuro and O. Plamenevskaya, Transverse knots, branched double covers and Heegaard Floer contact invariants, preprint, arXiv:0712.1557.
- [15] J. Hoste and M. Thistlethwaite (with help from Bruce Ewing, Ken Millett, Ken Stephenson and Jeff Weeks), *Knotscape 1.01 4-1-99*, <http://www.math.utk.edu/morwen/knotscape.html>.
- [16] V.F.R.Jones, Hecke algebra representations of braid groups and link polynomials, *Ann. of Math. (2)* **126** (1987) 335–388.
- [17] Private communication from Vaughan Jones.
- [18] E.S. Kang, K.H. Ko and S.J.Lee, Band-generator presentation for the 4-braid group, *Topology Appl.* **78**, (1997) 39–60.
- [19] R.D. Keever, Minimal 3-braid representations, *J. Knot Theory Ramifications* **3** (1994) 163–177.
- [20] E. Kin, The forcing partial order on a family of braids forced by pseudo-Anosov 3-braids, preprint, arXiv:0711.4398.
- [21] K.H. Ko and S.J. Lee, Flypes of closed 3-braids in the standard contact space, *J. of the Korean Math. Soc.* **36** (1999) 51–71.
- [22] M.T.Lozano and J. Przytycki, Incompressible surfaces in the exterior of a closed 3-braid. I. Surfaces with horizontal boundary components, *Math. Proc. Cambridge Philos. Soc.* **98** (1985) 275–299.
- [23] W. Menasco and X. Zhang, Positive Knots And Knots With Braid Index Three Have Property-P, *J. Knot Theory Ramifications* **12** (2003) 427–444.
- [24] K. Murasugi, On closed 3-braids, *Memoirs of the AMS* **151** (1974).
- [25] Y. Ni, Closed 3-braids are nearly fibered, preprint, arXiv:math.GT/0510243.
- [26] L. Ng, P. Ozsváth and D. Thurston, Transverse knots distinguished by knot Floer homology, preprint, arXiv:math.GT/0703446.
- [27] S. Orevkov, Quasipositivity problem for 3-braids, *Turkish J. Math.* **28** (2004) 89–93.
- [28] P. Ozsváth, Z. Szabó, and D. Thurston, Legendrian knots, transverse knots and combinatorial Floer homology, preprint, arXiv:math.GT/0611841.
- [29] M. S. Paterson and A. A. Razborov, The set of minimal braids is co-NP-complete, *J. Algorithms* **12** (1991) 393–408.
- [30] O. Plamenevskaya, Transverse knots, double branched double covers and Heegaard Floer Knot Homology, *J. Symplectic Geometry* **4** (2006) 149–170
- [31] O. Plamenevskaya, Transverse knots and Khovanov homology, *Math Research Letters* **13** (2006) 571–586.
- [32] D. Rolfsen, *Knots and Links*, Publish or Perish, 1976 and 1990.
- [33] O. Schreier, Uber die Gruppen $A^a B^b = 1$, *Abh. Math. Sem. Univ. Hamburg* **3** (1924) 167–169.
- [34] A. Stoimenow, Properties of closed 3-braids and other link braid representations, preprint, arXiv:math/0606435v2.
- [35] A. Stoimenow, The skein polynomial of closed 3-braids, *J. Reine Angew. Math.* **564** (2003) 167–180.
- [36] M. Takahashi, Explicit formulas for the Jones polynomials of closed 3-braids, *Comment. Math. Univ. St. Paul* **38** (1989) 129–167.
- [37] H. Trotter, Non-invertible knots exist, *Topology* **2** (1963) 275–280.
- [38] A. Usman, Conjugacy classes of 3-braid group B_3 , preprint, arXiv:0801.4443.

- [39] H. Wu, Braids, transversal links and the Khovanov-Rozansky theory, *Trans. Amer. Math. Soc.* **360** (2008) 3365–3389.
- [40] P.J. Xu, The genus of closed 3-braids, *J. Knot Theory Ramifications* **1** (1992) 303–326.
- [41] P.J.Xu, Growth of the positive braid semigroups, *J. Pure Appl. Algebra* **80** (1992) 197–215.