

A note on critical points of integrals of soliton equations

I. M. Krichever · D. V. Zakharov

Received: 29 September 2010 / Accepted: 9 November 2010 / Published online: 21 December 2010
© Springer Basel AG 2010

Abstract We consider the problem of extending the integrals of motion of soliton equations to the space of all finite-gap solutions. We consider the critical points of these integrals on the moduli space of Riemann surfaces with marked points and jets of local coordinates. We show that the solutions of the corresponding variational problem have an explicit description in terms of real-normalized differentials on the spectral curve. Such conditions have previously appeared in a number of problems of mathematical physics.

1 Introduction

In the modern theory of integrable systems, the constants of motion of a soliton equation are defined as integrals of certain differential polynomials involving the solution. These integrals are then regarded as Hamiltonians of compatible commuting flows, and the critical points of these Hamiltonians are stationary points of the corresponding flows. The purpose of this paper is to revisit this fundamental concept of the soliton theory in a more general setting and to present unexpected connections of the critical points of the soliton integrals with physical problems, including the theory of 2D

Research is supported in part by National Science Foundation under the grant DMS-04-05519.

I. M. Krichever (✉) · D. V. Zakharov
Columbia University, New York, USA
e-mail: krichev@math.columbia.edu

D. V. Zakharov
e-mail: zakharov@math.columbia.edu

I. M. Krichever
Landau Institute for Theoretical Physics, Moscow, Russia

I. M. Krichever
Kharkevich Institute for Information Transmission Problems, Moscow, Russia

quantum gravity, the Whitham approach to string equations, the theory of orthogonal polynomials, and the study of Hele–Shaw flows.

First of all, it is necessary to emphasize that the Hamiltonian theory of periodic non-singular solutions of soliton equations is well-defined only for solutions of a fixed period. It is not a priori clear that the Hamiltonians considered as functionals on the space of all periodic functions with variable periods have critical points. In fact, we will show that in the naïve formulation this problem has no solutions in the simplest case of the stationary KdV equation.

The problem becomes non-trivial and a posteriori interesting when extended onto the space of all finite-gap (algebro-geometric) solutions. Although these solutions are in general singular and complex functions of their arguments, their integrals can be defined in a pure algebraic-geometrical form and can be seen as functions on the moduli spaces of algebraic curves with fixed jets of local coordinates in the neighborhoods of punctures.

It is instructive to present the formulation of the extended problem and its solution in the case of the KdV equation

$$4u_t = 6uu_x - u_{xxx}. \quad (1.1)$$

This equation has an infinite number of integrals of motion, which in the periodic case have the form

$$H_n = \frac{1}{T} \int_0^T P_n(u, u', \dots, u^{(n+1)}) dx, \quad n = -1, 1, 3, 5, \dots, \quad (1.2)$$

where the P_n are certain differential polynomials in u :

$$P_{-1} = -\frac{u}{2}, \quad P_1 = -\frac{u^2}{8}, \quad P_3 = -\frac{u_x^2 + 2u^3}{32}, \dots \quad (1.3)$$

Consider an arbitrary linear combination $H = H_{2n+1} + c_{2n-1}H_{2n-1} + \dots + c_{-1}H_{-1}$ of the integrals on the space of functions with fixed period T , where the coefficients c_i are real. The critical points of such a functional are solutions of an ordinary differential equation known as the *n*th stationary KdV equation [1]. For example, the stationary points of the functional

$$H = H_3 - \frac{g_2 H_{-1}}{8} \quad (1.4)$$

are stationary solutions of the usual KdV equation:

$$16 \frac{\delta H}{\delta u} = u_{xx} - 3u^2 + g_2 = 0. \quad (1.5)$$

Let us now consider H as a functional on the space of all nonsingular periodic functions. Evaluating the variation of H with respect to the period T and setting it equal

to zero, we obtain the additional condition

$$\int_0^T u_x^2 dx = 0, \quad (1.6)$$

which has no solutions when u is real and smooth.

We now show how to extend the functional H to the space of all periodic solutions of Eq. (1.5). The general solution of Eq. (1.5) is given by the formula

$$u(x) = 2\wp(x + x_0), \quad (1.7)$$

where $\wp(x) = \wp(x|\omega, \omega')$ is the Weierstrass \wp -function corresponding to the elliptic curve Γ defined by the equation

$$Y^2 = E^3 - g_2 E - g_3 = (E - E_0)(E - E_1)(E - E_3). \quad (1.8)$$

The function u given by Eq. (1.7) is real and smooth when the branch points E_i of the elliptic curve are real, in which case we assume that $E_0 < E_1 < E_2$. For a generic elliptic curve, this solution is a quasi-periodic meromorphic function. For such functions, formula (1.2) requires regularization.

For a real smooth solution $u(x)$ of (1.5) defined by an elliptic curve Γ with real branch points, the values of the integrals (1.2) can be expressed in the following way (for details see [2]). There exists a unique meromorphic differential dQ on Γ , called the *quasi-momentum differential*, with a second order pole at infinity with principal part

$$dQ \sim \frac{dz}{z^2}, \quad z = \frac{1}{\sqrt{E}}, \quad (1.9)$$

and no other singularities, and which in addition satisfies the normalization condition

$$\int_{E_1}^{E_2} dQ = 0. \quad (1.10)$$

Then the values of the integrals (1.2) of a solution $u(x)$ corresponding to the curve Γ are the coefficients of the corresponding Abelian integral Q (which is the quasi-momentum of the auxiliary Sturm–Liouville operator $L = -\partial_x^2 + u$) at infinity:

$$Q = z^{-1} + \sum_{n=0}^{\infty} H_{2n-1} z^{2n+1}. \quad (1.11)$$

For an elliptic curve with real branch points, the coefficients H_n are real-valued.

As one of the authors has pointed out in [3], the quasi-momentum differential for a generic spectral curve is uniquely defined by Eq. (1.9) and by the condition that all its periods are *real*:

$$\operatorname{Im} \oint_{\gamma} dQ = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (1.12)$$

For elliptic curves with real branch points the two normalization conditions (1.10) and (1.12) coincide. We can then define the functions H_n on the moduli space of smooth elliptic curves as the coefficients of dQ at infinity, i.e. using formula (1.11). These functions are in general complex-valued. We then define the real-valued extension of the KdV integrals (1.2) to be real parts of these functions, i.e. we set

$$\mathcal{H}_n = \operatorname{Re} H_n. \quad (1.13)$$

The main results of this paper are Theorems 2.2 and 3.2, which describe the critical points of such functionals. In the case under consideration, Theorem 2.2 implies that an elliptic curve Γ given by Eq. (1.8) is a critical point of the extended Hamiltonian $\mathcal{H} = \mathcal{H}_3$ if and only if the following condition is satisfied:

$$\operatorname{Im} \oint_{\gamma} Y dE = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (1.14)$$

This equation has already appeared in the framework of the Whitham approach to string equations [4–8]. Later similar conditions were introduced in the theory of 2D-gravity [9, 10] and in the study of distribution of zeros in orthogonal polynomials [11]. Recently, the physical meaning of this condition in the theory of Hele-Shaw flow was revealed in [12].

Equation (1.14) was solved numerically in [6]. It turns out that there is a unique solution in the space of smooth elliptic curves. For real $g_2 > 0$ the value of g_3 of the corresponding curve is

$$g_3 = \frac{4h^2 + 1}{(4h^2 - 3)^{3/2}} g_2^{3/2}, \quad (1.15)$$

where $h \approx 3.2463822253744278875676$.

The original goal of the authors was to study the geometry of the moduli spaces of smooth curves using the extended soliton integrals as Morse functions. The connection of the variational problem for these integrals with important physical problems is to some extent an unexpected byproduct of the authors' original goal.

In the most general setting, the extended soliton integrals are defined as functions on the moduli space $\mathcal{M}_{g,N}(\mathbf{n}, \mathbf{m}) = \{(\Gamma, P_\alpha, [z_\alpha], E, Q)\}$ of smooth genus g algebraic curves Γ with N marked points P_1, \dots, P_N with jets of local coordinates $[z_\alpha]$ at P_α and with a pair of Abelian integrals, denoted E and Q for historical reasons, having poles of orders n_α and m_α at the marked points P_α . This moduli space is central to several

theories with very distinct goals and origins. These include the non-linear WKB (or Whitham) theory, two-dimensional topological models, Seiberg–Witten exact solutions of $N = 2$ supersymmetric gauge theories (see [13–15] and references therein), and the theory of Hele–Shaw flows [12]. Recently, in [16, 17] it was shown that the constructions of the Whitham theory on the moduli space $\mathcal{M}_{g,N}^{\mathbf{m}}(\mathbf{n})$ provide a geometrical explanation of cohomological vanishing results on the moduli spaces $\mathcal{M}_{g,n}$ of smooth curves with marked points.

2 The one-point case

The object of our study in this paper is the moduli space of Riemann surfaces with marked points and with jets of local coordinates at the marked points. Given a Riemann surface with jets of local coordinates at marked points, we construct a pair of real-normalized Abelian integrals on the surface. We then define a collection of functions on the moduli space by considering the periods of these integrals and the coefficients of their expansions at the marked points, and we consider the critical points of these functions on the leaves of a hierarchy of foliations defined on the moduli space. We show that the critical points of these functions admit an explicit analytic description. To clarify the exposition, in this section we describe the case of one marked point.

Definition Let Γ be a Riemann surface, let P be point on Γ . An n -jet of local coordinates at P is an equivalence class of local coordinates at P , with z equivalent to z' if $z' = z + O(z^{n+1})$.

Definition Let ω be a meromorphic differential on a Riemann surface Γ . We say that ω is *real-normalized* if the integral of ω along any cycle in Γ is a real number, i.e. if

$$\operatorname{Im} \oint_{\gamma} \omega = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.1)$$

The period matrix of a Riemann surface has positive definite imaginary part, therefore a holomorphic real-normalized differential is identically zero. More generally, given a collection of principal parts with imaginary residues, there exists a unique real-normalized meromorphic differential with these principal parts:

Proposition 2.1 *Let Γ be a Riemann surface, let P_1, \dots, P_N be points on Γ , let z_α be local coordinates in neighborhoods of P_α , and let $c_{\alpha,k}$, for $\alpha = 1, \dots, N, k = 1, \dots, n_\alpha$ be complex numbers, such that $c_{\alpha,1}$ are purely imaginary and $\sum c_{\alpha,1} = 0$. Then there exists a unique real-normalized meromorphic differential ω on Γ with poles at P_α of the form*

$$\omega = \left(\frac{c_{\alpha,n_\alpha}}{z_\alpha^{n_\alpha}} + \dots + \frac{c_{\alpha,1}}{z_\alpha} + O(1) \right) dz_\alpha \text{ near } P_\alpha, \quad (2.2)$$

and with no other singularities.

Definition Let Γ be a Riemann surface and let P_0 be a point on Γ . An *Abelian integral with fixed branch P_0* is a multi-valued function $A(P, \gamma)$ on Γ depending on a choice of path of integration γ from P_0 to P , and such that its differential dA is a single-valued meromorphic differential on Γ . We say that an Abelian integral is *real-normalized* if the corresponding differential is.

In the neighborhood of any point that is not a residue of dA , the Abelian integral A is locally a single-valued function. In particular, if dA has no residues at P_0 , then in a sufficiently small neighborhood of P_0 the Abelian integral A has a selected branch corresponding to the trivial path from P_0 .

We consider the following moduli spaces:

- $\mathcal{M}_{g,1}(n)$ is the moduli space of Riemann surfaces Γ with a marked point P and an $(n+1)$ -jet of local coordinates $[z]$ at P . This space is our principal object of interest.
- $\mathcal{M}_{g,1}^m(n)$ is the moduli space of Riemann surfaces Γ with a marked point P , an $(n+1)$ -jet of local coordinates $[z]$ at P , and a real-normalized Abelian integral Q with a fixed branch at P , having a pole of order m at P and no other singularities. This space is the natural domain of the extended Hamiltonian functions that we are studying.
- $\mathbf{M}_{g,1}(m, n)$ is the moduli space of Riemann surfaces Γ with a marked point P , an $(n+1)$ -jet of local coordinates $[z]$ at P , and a pair of Abelian integrals E and Q (not necessarily real-normalized) with fixed branches at P , having poles of order n and m at P , respectively, and no other singularities, where we in addition assume that the selected branch of E at P has the form $E = z^{-n} + O(z)$. This space is a slight modification of the moduli space introduced in the paper [14].

These spaces are related by the following maps:

$$\begin{array}{ccc} \mathcal{M}_{g,1}^m(n) & \hookrightarrow & \mathbf{M}_{g,1}(n, m) \\ \downarrow & & \\ \mathcal{M}_{g,1}(n) & & \end{array} \quad (2.3)$$

where the vertical map consists in forgetting the integrals.

Let Γ be a Riemann surface, and let $[z]$ be an $(n+1)$ -jet of local parameters at a point P on Γ . By Proposition 2.1 there exists a unique real-normalized Abelian integral E on Γ with a fixed branch at P , where it has a pole of order n of the form $E = z^{-n} + O(z)$, and with no other singularities. There exists a unique local coordinate z in the jet $[z]$ such that the fixed branch of E can be written as

$$E = z^{-n} \text{ near } P, \quad (2.4)$$

where the equality is now assumed to be exact. Therefore, the notion of real-normalized Abelian integrals gives us a natural choice of a representative in the jet $[z]$.

Let Q be another Abelian integral on Γ . We can expand it in terms of this chosen local coordinate:

$$Q = \sum_{j=0}^m q_j z^{-j} + O(z) \text{ near } P. \quad (2.5)$$

By virtue of Proposition 2.1 we can identify the Abelian integral Q with the polynomial $q(t) = q_m t^m + \dots + q_0$. Therefore, the vertical map in the diagram (2.3) is a trivial bundle with fiber \mathbb{C}^{m+1} . Any polynomial $q(t)$ of degree m determines a section of this bundle, i.e. an embedding of $\mathcal{M}_{g,1}(n)$ into $\mathcal{M}_{g,1}^m(n)$ and hence into $\mathbf{M}_{g,1}(n, m)$.

We now consider a collection of functions on the space $\mathbf{M}_{g,1}(n, m)$. Let $(\Gamma, P, [z], E, Q)$ be a point of $\mathbf{M}_{g,1}(n, m)$. Consider the coefficients of the expansion of the differential $Q dE$ at the point P with respect to the chosen local coordinate z :

$$Q dE = \left(\sum_{k=0}^{n+m} T_k z^{-k-1} + \sum_{j=1}^{\infty} H_j z^{j-1} \right) dz. \quad (2.6)$$

The coefficients of the principal part T_k act as coordinates on the space $\mathbf{M}_{g,1}(n, m)$. The coefficients of the regular part H_j are the Hamiltonian functions that are of interest to us.

The following additional functions are only locally well defined on $\mathbf{M}_{g,1}(n, m)$. Fix a basis of cycles A_i, B_i in $H_1(\Gamma, \mathbb{Z})$ with canonical intersection form, such that no cycle passes through P or the zero-divisors of dE or dQ . Let $\bar{\Gamma}$ denote the Riemann surface cut along these cycles and let $\partial\Gamma$ denote the boundary of $\bar{\Gamma}$. The branches E and Q fixed at P then extend to single-valued functions on $\bar{\Gamma}$. Consider the real-valued periods of the differentials dE and dQ along the chosen cycles:

$$\tau_{A_i}^E = \oint_{A_i} dE, \quad \tau_{B_i}^E = \oint_{B_i} dE, \quad \tau_{A_i}^Q = \oint_{A_i} dQ, \quad \tau_{B_i}^Q = \oint_{B_i} dQ, \quad i = 1, \dots, g, \quad (2.7)$$

and the A -periods of the differential $Q dE$:

$$a_i = \oint_{A_i} Q dE, \quad i = 1, \dots, g. \quad (2.8)$$

The residue of the differential $Q dE$ at P is equal to its integral over the boundary $\partial\Gamma$, which gives the only relation between these functions:

$$2\pi i T_0 = \sum_{i=1}^g \tau_{A_i}^Q \tau_{B_i}^E - \tau_{B_i}^Q \tau_{A_i}^E. \quad (2.9)$$

Therefore, we can exclude T_0 from consideration.

The functions T_k and H_j do not depend on the choice of basis for $H_1(\Gamma, \mathbb{Z})$ and are well-defined on all of $\mathbf{M}_{g,1}(n, m)$, while the other functions defined above depend on

the choice of cycles. However, in the neighborhood of any point $\mathbf{M}_{g,1}(n, m)$ this choice can be made in a consistent way for all curves, therefore these functions are *locally* well defined. In [14] it was shown that these functions can be used as local coordinates on $\mathbf{M}_{g,1}(n, m)$:

Theorem 2.1 *Let $(\Gamma, P, [z], E, Q)$ be a point $\mathbf{M}_{g,1}(n, m)$ such that the zero-divisors of dE and dQ on Γ do not intersect. Then in a neighborhood of this point the functions T_k (excluding T_0), $\tau_{A_i}^E, \tau_{B_i}^E, \tau_{A_i}^Q, \tau_{B_i}^Q$ and a_i (defined with respect to a consistent choice of cycles) have linearly independent differentials and therefore define a local holomorphic coordinate system for $\mathbf{M}_{g,1}(n, m)$.*

We now proceed as follows. Fix a polynomial $q(z)$ of degree m , and consider the embedding $\mathcal{M}_{g,1}(n) \hookrightarrow \mathcal{M}_{g,1}^m(n) \hookrightarrow \mathbf{M}_{g,1}(n, m)$ defined by that polynomial. We then define a hierarchy of foliations on $\mathbf{M}_{g,1}(n, m)$ and consider their restrictions to $\mathcal{M}_{g,1}(n)$. Finally, we define a collection of functions on $\mathcal{M}_{g,1}(n)$ and study the critical points of their restrictions to the leaves of the various foliations.

The period functions (2.7) are defined with respect to a choice of basis of $H_1(\Gamma, \mathbb{Z})$, and any two such choices are related by an element of $Sp(2n, \mathbb{Z})$. Therefore, at any point $p \in \mathbf{M}_{g,1}(n)$ the intersection of the null spaces of the differentials $d\tau_{A_i}^E$ and $d\tau_{B_i}^E$ in the tangent space $T_p(\mathbf{M}_{g,1}(n))$ does not depend on the choice of basis and is therefore a well-defined subspace $T_p^E(\mathbf{M}_{g,1}(n))$ of $T_p(\mathbf{M}_{g,1}(n))$. Moreover, these subspaces are the tangent spaces to the level sets of some locally well-defined functions, therefore they form an integrable distribution on $\mathbf{M}_{g,1}(n, m)$. Similarly, the intersection of the null spaces of $d\tau_{A_i}^Q$ and $d\tau_{B_i}^Q$ is a well-defined integrable distribution $T_p^Q(\mathbf{M}_{g,1}(n))$ on $\mathbf{M}_{g,1}(n)$.

The embedding $\mathcal{M}_{g,1}^m(n) \hookrightarrow \mathbf{M}_{g,1}(n, m)$ is locally defined by the condition that the functions (2.7) are real-valued. This condition does not depend on the choice of basis for $H_1(\Gamma, \mathbb{Z})$. Therefore, the distributions defined above descend to $\mathcal{M}_{g,1}^m(n)$. Furthermore, the space $\mathcal{M}_{g,1}(n)$ is a level set of the functions T_n, \dots, T_{n+m} , and therefore by Proposition 2.1 these distributions descend to $\mathcal{M}_{g,1}(n)$.

Let $\Lambda \subset \mathcal{M}_{g,1}(n)$ denote a joint level set of the functions T_1, \dots, T_{n-1} . Let T_p^E denote the intersection of the null spaces of the differentials $d\tau_{A_i}^E$ and $d\tau_{B_i}^E$ on the space $T_p(\Lambda)$. Let $\mathcal{L}_E \subset \Lambda$ denote an integral submanifold of this distribution. Along \mathcal{L}_E , the periods of dE are constant:

$$\delta \oint_\gamma dE = 0 \quad \forall \delta \in T_p(\mathcal{L}_E) = T_p^E \quad \text{and} \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.10)$$

Similarly, let $T_p^Q \subset T_p(\Lambda)$ denote the directions along which the periods of dQ do not change, and let $\mathcal{L}_Q \subset \Lambda$ denote the corresponding integral submanifold:

$$\delta \oint_\gamma dQ = 0 \quad \forall \delta \in T_p(\mathcal{L}_Q) = T_p^Q \quad \text{and} \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.11)$$

Finally, let $\mathcal{L} = \mathcal{L}_E \cap \mathcal{L}_Q$ denote a submanifold along which the periods of both dE and dQ are constant:

$$\delta \oint_{\gamma} dE = \delta \oint_{\gamma} dQ = 0 \quad \forall \delta \in T_p(\mathcal{L}) = T_p^E \cap T_p^Q \quad \text{and} \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.12)$$

We now define a real-valued function on $\mathcal{M}_{g,1}(n)$ and study its critical points on the leaves of the foliations defined above. Let c_j, d_j , for $j = 1, \dots, M$ be real coefficients. Consider the following function on $\mathcal{M}_{g,1}(n)$:

$$\mathcal{H} = \sum_{j=1}^M (c_j \operatorname{Re} H_j + d_j \operatorname{Im} H_j). \quad (2.13)$$

The following theorem, which is the principal result of this chapter, describes the critical points of \mathcal{H} restricted to the leaves of the foliations defined above:

Theorem 2.2 *Let $(\Gamma, P, [z])$ be a point in $\mathcal{M}_{g,1}(n)$, which we assume to be embedded in $\mathcal{M}_{g,1}^m(n)$ via the polynomial $q(z)$. Let E and Q be the real-normalized Abelian integrals defined by Eqs. (2.4) and (2.5), and let D denote the intersection of the zero-divisors of dE and dQ .*

- A. *The point $(\Gamma, P, [z])$ is a critical point of \mathcal{H} restricted to \mathcal{L} if and only if*
 - (i) *There exists a meromorphic function Y on Γ with a pole at P with the principal part*

$$Y = \sum_{j=1}^{\infty} (c_j - id_j) z^{-j} + O(z) \text{ near } P, \quad (2.14)$$

with poles at the divisor D , and no other singularities.

- B. *The point $(\Gamma, P, [z])$ is a critical point of \mathcal{H} restricted to \mathcal{L}_E if and only if there exists a function Y satisfying condition (i) and in addition the following condition:*
 - (ii) *the differential $Y dE$ is real-normalized, i.e.*

$$\operatorname{Im} \oint_{\gamma} Y dE = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.15)$$

- C. *The point $(\Gamma, P, [z])$ is a critical point of \mathcal{H} restricted to \mathcal{L}_Q if and only if there exists a function Y satisfying condition (i) and in addition the following condition:*
 - (iii) *the differential $Y dQ$ is real-normalized, i.e.*

$$\operatorname{Im} \oint_{\gamma} Y dQ = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.16)$$

- D.** *The point $(\Gamma, P, [z])$ is a critical point of \mathcal{H} restricted to Λ if and only if there exists a function satisfying conditions (i), (ii) and (iii).*

Remark. The correspondence between critical points of integrals of soliton equations and meromorphic functions with prescribed singularities on the spectral curve, in other words part A of our theorem, has long been a central notion in the theory of finite-gap integration (see, for example, the survey [2]). However, this result has never before been stated in such an explicit form. Moreover, the known results only apply to the case when the equations are Hamiltonian, which corresponds to the case when the zero-divisors of the differentials dE and dQ do not intersect.

The Hamiltonian structure of the integrable systems associated with the moduli space $\mathcal{M}_{g,N}(n)$ is determined by the Novikov–Veselov–Witten–Seiberg symplectic form, which is defined as follows. Consider a leaf \mathcal{L} of the foliation defined above, and consider the fibration \mathcal{N}^g over \mathcal{L} whose fiber over a point $(\Gamma, P, [z])$ is the g -th symmetric power of Γ . This fibration has dimension $2g$ and is the phase space for the corresponding integrable system. The Novikov–Veselov–Witten–Seiberg symplectic form is defined as

$$\omega_{\mathcal{M}} = \sum_{i=1}^g da_i \wedge d\omega_i, \quad (2.17)$$

where ω_i are normalized holomorphic differentials on Γ .

If $(\Gamma, P, [z])$ is a point at which the zero-divisors of dE and dQ intersect, then according to Theorem 2.1 the coordinates (2.8) are not independent on the leaf \mathcal{L} . Therefore, the symplectic form $\omega_{\mathcal{M}}$ is degenerate at $(\Gamma, P, [z])$, and therefore the soliton equation is not Hamiltonian.

Example Genus one solutions of KdV. Before proceeding with the proof of the theorem, we show how it applies to the example considered in the introduction. Assume that $g = 1, n = 2$ and $m = 1$, and consider the embedding $\mathcal{M}_{1,1}(2) \hookrightarrow \mathcal{M}_{1,1}^1(2)$ given by the polynomial $q(z) = z$. Suppose furthermore that we are on a leaf \mathcal{L}_E along which the periods of dE are identically zero. For every point $(\Gamma, P, [z]) \in \mathcal{L}_E$, Γ is an elliptic curve and the Abelian integral E is a single-valued function on Γ of degree two, having a second order pole at P of the form $E = z^{-2}$ and no other singularities.

The differential $Q dE$ is odd with respect to the hyperelliptic involution and therefore has the following expansion at P :

$$Q dE = -2 \left(z^{-4} + T_1 z^{-2} + H_1 + H_3 z^2 + O(z^4) \right) dz. \quad (2.18)$$

We consider the critical points of the function $\mathcal{H} = \operatorname{Re} H_3$, which are the same as those of the function (1.4), since $H_{-1} = T_1$ is constant along \mathcal{L}_E . According to the theorem, a point $(\Gamma, P, [z])$ is a critical point of H restricted to \mathcal{L} if there exists a meromorphic function Y on Γ with a triple pole at P of the form

$$Y = z^{-3} + O(z) \text{ near } P. \quad (2.19)$$

The function Y then satisfies an equation of the form (1.8), so $(\Gamma, P, [z], E, Y, Q)$ are the spectral data of a stationary genus one solution of the KdV equation. If the point $(\Gamma, P, [z])$ is a critical point of \mathcal{H} on all of \mathcal{L}_E , then the function Y satisfies the condition

$$\text{Im} \oint_{\gamma} Y dE = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.20)$$

The main tool in the proof of the theorem is a homomorphism

$$f : T_p(\Lambda) \rightarrow \Omega(\Gamma, P, E, Q) \quad (2.21)$$

from the tangent space to Λ at a point $p = (\Gamma, P, [z])$ to a space of certain multi-valued differentials on Γ . This homomorphism was introduced in Whitham theory in [13] (see more details in [14, 15]) and modified for the case real normalized differentials in [18]. It is defined as follows.

Let $\bar{\Gamma}$ denote the surface cut along a choice of basis cycles, and let $\sum v_s E_s$ be the zero divisor of the differential dE . On $\bar{\Gamma}$, the branch of E fixed at P can be used as a local coordinate in the neighborhood of any point except E_s . This allows us to lift a tangent vector $\delta \in T_p(\Lambda)$ to a tangent vector on the total space of the universal curve over \mathcal{L}_E and defines the partial derivative of $Q = Q(E)$ for fixed levels of E , which we denote by $\delta Q|_E$.

The derivative $\delta Q|_E$ is holomorphic away from the points E_s , where it has poles of order v_s , and from the cuts. Along the two sides of a cut A_i or B_i , the values of E and Q have jumps equal to the corresponding periods of dE and dQ :

$$Q\left(E + \tau_{A_i}^E\right) - Q(E) = \tau_{A_i}^Q, \quad Q\left(E + \tau_{B_i}^E\right) - Q(E) = \tau_{B_i}^Q. \quad (2.22)$$

Therefore, the derivative $\delta Q|_E$ has jumps along the cuts A_i and B_i equal to

$$\begin{aligned} \delta Q\left(E + \tau_{A_i}^E\right) - \delta Q(E) &= \delta \tau_{A_i}^Q - \frac{dQ}{dE} \delta \tau_{A_i}^E, \\ \delta Q\left(E + \tau_{B_i}^E\right) - \delta Q(E) &= \delta \tau_{B_i}^Q - \frac{dQ}{dE} \delta \tau_{B_i}^E. \end{aligned} \quad (2.23)$$

The map f associates to a tangent vector δ the multi-valued meromorphic differential

$$f(\delta) = \Omega_{\delta} = (\delta Q|_E)dE. \quad (2.24)$$

Since all the higher order terms in the expansion (2.6) of $Q dE$ at P are fixed on Λ , the differential Ω_{δ} has at most a simple pole at P . At the points E_s , the zeroes of dE cancel the poles of $\delta Q|_E$, hence Ω_{δ} has no other singularities. Finally, along the cuts A_i and B_i the differential Ω_{δ} has jumps which are linear combinations of the differentials dE and dQ with real coefficients.

The space $\Omega(\Gamma, P, E, Q)$ of differentials with these properties has real dimension $6g$, the same as \mathcal{L} . It is clear from the construction that the image of a tangent vector lying inside one of the subspaces $T_p(\mathcal{L}_E)$ or $T_p(\mathcal{L}_Q)$ is a differential with jumps proportional only to dE or dQ , respectively, and that the image of their intersection $T_p(\mathcal{L}) = T_p(\mathcal{L}_E) \cap T_p(\mathcal{L}_Q)$ lies inside the space of holomorphic differentials $\Omega^h(\Gamma) \subset \Omega(\Gamma, P, E, Q)$. A complete description of the homomorphism (2.21) was given in [14]:

Proposition 2.2 *Let $D = \sum \mu_s Q_s$ denote the intersection of the zero divisors of dE and dQ . Then the image of the tangent space $T_p(\Lambda)$ under the homomorphism f of (2.21) is the subspace $\Omega_D(\Gamma, P, E, Q) \subset \Omega(\Gamma, P, E, Q)$ of differentials having zeroes at Q_s of orders μ_s . In particular, if the zero divisors of dE and dQ do not intersect, then the map (2.21) is an isomorphism.*

The space $\Omega(\Gamma, P, E, Q)$ is spanned over \mathbb{R} by the following collection of differentials:

- $2g$ single-valued holomorphic differentials $\Omega_{A_i}^h, \Omega_{B_i}^h, i = 1, \dots, g$ normalized by the conditions

$$\operatorname{Im} \left(\oint_{A_j} \Omega_{A_i}^h \right) = \delta_{ij}, \quad \operatorname{Im} \left(\oint_{B_j} \Omega_{A_i}^h \right) = 0, \quad j = 1, \dots, g, \quad (2.25)$$

$$\operatorname{Im} \left(\oint_{A_j} \Omega_{B_i}^h \right) = 0, \quad \operatorname{Im} \left(\oint_{B_j} \Omega_{B_i}^h \right) = \delta_{ij}, \quad j = 1, \dots, g. \quad (2.26)$$

- $2g$ differentials $\Omega_{A_i}^E, \Omega_{B_i}^E, i = 1, \dots, g$ having a simple pole at P and a single jump equal to dE along the cuts A_i or B_i , respectively:

$$\left(\Omega_{A_i}^E \right)^+ - \left(\Omega_{A_i}^E \right)^- = dE \text{ along } A_i, \quad \left(\Omega_{B_i}^E \right)^+ - \left(\Omega_{B_i}^E \right)^- = dE \text{ along } B_i, \quad (2.27)$$

normalized by the conditions

$$\begin{aligned} \operatorname{Im} \oint_{A_j} \Omega_{A_i}^E &= 0, & \operatorname{Im} \oint_{B_j} \Omega_{A_i}^E &= 0, & \operatorname{Im} \oint_{A_j} \Omega_{B_i}^E &= 0, \\ \operatorname{Im} \oint_{B_j} \Omega_{B_i}^E &= 0, & j &= 1, \dots, g. \end{aligned} \quad (2.28)$$

- $2g$ differentials $\Omega_{A_i}^Q, \Omega_{B_i}^Q, i = 1, \dots, g$ having a simple pole at P and a single jump equal to dQ along the cuts A_i or B_i , respectively:

$$\left(\Omega_{A_i}^Q\right)^+ - \left(\Omega_{A_i}^Q\right)^- = dQ \text{ along } A_i, \quad \left(\Omega_{B_i}^Q\right)^+ - \left(\Omega_{B_i}^Q\right)^- = dQ \text{ along } B_i, \quad (2.29)$$

normalized by the conditions

$$\begin{aligned} \operatorname{Im} \oint_{A_j} \Omega_{A_i}^Q &= 0, & \operatorname{Im} \oint_{B_j} \Omega_{A_i}^Q &= 0, & \operatorname{Im} \oint_{A_j} \Omega_{B_i}^Q &= 0, \\ \operatorname{Im} \oint_{B_j} \Omega_{B_i}^Q &= 0, & j &= 1, \dots, g. \end{aligned} \quad (2.30)$$

Remark. The integral of a multi-valued differential along a curve in $\bar{\Gamma}$ is not determined by the homology class of that curve in Γ . For this reason, the integrals in the above formulas are assumed to be taken over the particular choice of A_i and B_i that we have made.

Proof of Theorem 2.2. We first prove the statement of the theorem for the critical points of \mathcal{H} restricted to the open subset of Λ where the differentials dQ and dE have no common zeros.

Given a set of coefficients c_j, d_j there is a unique real-normalized meromorphic differential dY of the second kind whose Abelian integral has a single pole at P of the form (2.14). By the definition of \mathcal{H} the derivative $\delta\mathcal{H}$ along a tangent vector δ is equal to

$$\delta\mathcal{H} = \operatorname{Re} \operatorname{Res}_P (Y\delta Q|_E dE) = -\frac{1}{2\pi} \operatorname{Im} \left(\oint_{\partial\Gamma} Y\Omega_\delta \right), \quad (2.31)$$

where $\partial\Gamma$ is the boundary of $\bar{\Gamma}$. Hence, the point $(\Gamma, P, [z])$ is a critical point of \mathcal{H} restricted to \mathcal{L} if and only if

$$\operatorname{Im} \left(\oint_{\partial\Gamma} Y\Omega \right) = \sum_{i=1}^g \left(\oint_{A_i} dY \cdot \operatorname{Im} \oint_{B_i} \Omega - \oint_{B_i} dY \cdot \operatorname{Im} \oint_{A_i} \Omega \right) = 0 \quad (2.32)$$

for all holomorphic differentials $\Omega \in \Omega^h(\Gamma)$. Plugging the differentials $\Omega_{A_i}^h$ and $\Omega_{B_i}^h$ in this equation, we see that all the periods of Y are zero:

$$\oint_{\gamma} dY = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (2.33)$$

Therefore Y is a single-valued meromorphic function, which proves part A of the theorem. To prove parts B and C of the theorem, we plug the remaining differentials $\Omega_{A_i}^E, \Omega_{B_i}^E, \Omega_{A_i}^Q, \Omega_{B_i}^Q$ into Eq. (2.32), which gives us conditions (2.15) and (2.16). Finally, since the tangent space $T_p(\Lambda)$ is spanned by $T_p(\mathcal{L}_E)$ and $T_p(\mathcal{L}_Q)$, part D follows from parts B and C.

The proof of the theorem in the general case goes along the same lines. Suppose that $p = (\Gamma, P, [z])$ is a critical point of \mathcal{H} restricted to \mathcal{L} , and let

$$D = \sum_s \mu_s Q_s = (dE)_0 \cap (dQ)_0. \quad (2.34)$$

be the intersection of the zero-divisors of dE and dQ . The image of the tangent subspace $T_p(\mathcal{L})$ under f is the space of holomorphic differentials on Γ vanishing at D , which we denote $\Omega_D^h(\Gamma)$. Let \mathcal{Y}_{D,c_i} denote space of real-normalized Abelian integrals on Γ having poles of orders μ_s at Q_s and a pole at the marked point P of the form (2.14). This is an affine space of real dimension $\dim_{\mathbb{R}} \mathcal{Y}_{D,c_i} = 2 \sum_s \mu_s = 2d$. The period vectors of the Abelian integrals in \mathcal{Y}_{D,c_i} form an affine subspace Π_{D,c_i} of \mathbb{R}^{2g} of dimension $2d - 2h^0(\Gamma, D) + 2$.

It is easy to see that we can evaluate the derivative $\delta\mathcal{H}$ by the same formula (2.31) using any $Y \in \mathcal{Y}_{D,c_i}$. Therefore, Eq. (2.32) holds for all $Y \in \mathcal{Y}_{D,c_i}$ and for all $\Omega \in \Omega_D^h(\Gamma)$. This gives $\dim_{\mathbb{R}} \Omega_D^h(\Gamma) = 2h^1(\Gamma, D) = 2g - 2d + 2h^0(\Gamma, D) - 2$ linearly independent conditions on the elements of Π_{D,c_i} . Therefore, Π_{D,c_i} is actually a linear subspace of \mathbb{R}^{2n} . Therefore, there exists an Abelian integral $Y \in \mathcal{Y}_{D,c_i}$ with trivial periods, that is to say a single-valued meromorphic function, which proves part A of the theorem.

Let Y_{D,c_i} denote the subspace of single-valued functions in \mathcal{Y}_{D,c_i} , we have shown above that $\dim_{\mathbb{R}} Y_{D,c_i} = 2h^0(\Gamma, D) - 2$. The image of $T_p(\mathcal{L}_E)$ under f is the space of differentials with jumps proportional to dE with real coefficients, having a simple pole at P and zeroes at the divisor D . We denote this space by $\Omega_D(\Gamma, P, E)$. Let $J_e : \Omega_D(\Gamma, P, E) \rightarrow \mathbb{R}^{2g}$ be the map which associates to each differential its jumps along the basis cycles. Since $\dim_{\mathbb{R}} \Omega_D(\Gamma, P, E) = 4g - 2d$ and $\dim_{\mathbb{R}} \text{Ker } J_E = \dim_{\mathbb{R}} \Omega_D^h(\Gamma) = 2h^1(\Gamma, D)$, we see that $\dim_{\mathbb{R}} \text{Im } J_E = 2g + 2 - 2h^0(\Gamma, D) = 2g - \dim_{\mathbb{R}} Y_{D,c_i}$. Therefore, formula (2.32) gives $2g - \dim_{\mathbb{R}} Y_{D,c_i}$ linear relations on the $2g$ imaginary parts of the periods of YdE . Therefore, there exists a $Y \in Y_{D,c_i}$ such that the differential YdE is real-normalized, which proves part B. The proof of part C is identical.

Finally, to prove part D, consider the image $\Omega_D(\Gamma, P, E, Q)$ of $T_p(\Lambda)$ under f , and let $J_{E,Q} : \Omega_D(\Gamma, P, E, Q) \rightarrow \mathbb{R}^{2g} \oplus \mathbb{R}^{2g}$ be the map that associates to each differential its jumps, which are a real linear combination of dE and dQ . Since $\dim_{\mathbb{R}} \Omega_D(\Gamma, P, E, Q) = 6g - 2d$ and $\dim_{\mathbb{R}} \text{Ker } J_{E,Q} = \dim_{\mathbb{R}} \Omega_D^h(\Gamma) = 2h^1(\Gamma, D)$, formula (2.32) gives $\dim_{\mathbb{R}} \text{Im } J_{E,Q} = 4g + 2 - 2h^0(\Gamma, D)$ linear conditions on the $4g$ imaginary parts of the periods of YdE and YdQ . Therefore, there

exists a $Y \in Y_{D,c_i}$ such that both YdE and $Y dQ$ are real-normalized, which completes the proof of the theorem.

3 The N -point case

In this section, we extend our problem to the moduli space of curves with several marked points and with jets of local coordinates at those points. Given a curve with jets of local coordinates, we define a real-normalized Abelian integral with poles determined by those jets. The principal difference in the N -point case is that is that we can only fix a branch of an Abelian integral at one marked point. At all other marked points our Abelian integrals will be truly multi-valued functions. This means that we cannot use the integral E to specify a unique local coordinate in the jet, as we did in (2.4), and therefore the functions (2.6) are no longer globally defined on the moduli space. Furthermore, the expansion (2.5) of the Abelian integral Q depends on the choice of the local coordinate, therefore the map forgetting the second Abelian integral Q is in general no longer a trivial bundle. In this section, we show that under certain assumptions the imaginary parts of the functions (2.6) are well-defined, which allows us to construct a somewhat more limited extension of the results of the previous chapter.

Let $\mathbf{n} = (n_1, \dots, n_N)$ and $\mathbf{m} = (m_1, \dots, m_N)$ denote a pair of multi-indices. We assume that $n_\alpha > 0$ and $m_\alpha \geq 0$ for all α , and that $m_\alpha \leq n_\alpha$ for $\alpha = 2, \dots, N$; the meaning of the last condition will become clear later. We consider the following spaces:

- $\mathcal{M}_{g,N}(\mathbf{n})$ is the moduli space of Riemann surfaces Γ with marked points P_1, \dots, P_N , with an $(n_1 + 1)$ -jet of local coordinates $[z_1]$ at P_1 and with n_α -jets of local coordinates $[z_\alpha]$ at P_α for $\alpha = 2, \dots, n$.
- $\mathcal{M}_{g,N}^{\mathbf{m}}(\mathbf{n})$ is the moduli space of Riemann surfaces Γ with marked points P_1, \dots, P_N , with an $(n_1 + 1)$ -jet of local coordinates $[z_1]$ at P_1 and with n_α -jets of local coordinates $[z_\alpha]$ at P_α for $\alpha = 2, \dots, N$, and with a real-normalized Abelian integral Q with fixed branch at P_1 , having poles of orders m_α at P_α and no other singularities.
- $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$ is the moduli space of Riemann surfaces Γ with marked points P_1, \dots, P_N , with an $(n_1 + 1)$ -jet of local coordinates $[z_1]$ at P_1 and with n_α -jets of local coordinates $[z_\alpha]$ at P_α for $\alpha = 2, \dots, n$, and with a pair of Abelian integrals E and Q (not necessarily real-normalized) with fixed branches at P_1 , having poles of orders n_α and m_α at P_α , respectively, and no other singularities, where we in addition assume that $E = z_1^{-n_1} + O(z_1)$ near P_1 and $E = z_\alpha^{-n_\alpha} + O(1)$ near P_α for $\alpha = 2, \dots, N$. Here we assume for simplicity that the integral E has no logarithmic singularities.

These spaces are related by the following maps:

$$\begin{array}{ccc} \mathcal{M}_{g,N}^{\mathbf{m}}(\mathbf{n}) & \hookrightarrow & \mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m}) \\ \downarrow & & \\ \mathcal{M}_{g,N}(\mathbf{n}) & & \end{array} \tag{3.1}$$

where the vertical map consists in forgetting the integrals.

Let $[z_1]$ be a $(n_1 + 1)$ -jet of local coordinates at P_1 and let $[z_\alpha]$ be n_α -jets of local coordinates at P_α for $\alpha = 2, \dots, N$. According to Proposition 2.1 there exists a unique real-normalized Abelian integral E on Γ with a fixed branch at P_1 and with poles of orders n_α at P_α of the form

$$E = z_1^{-n_1} + O(z_1) \text{ near } P_1, \quad E = z_\alpha^{-n_\alpha} + O(1) \text{ near } P_\alpha, \quad \alpha = 2, \dots, N, \quad (3.2)$$

and with no other singularities. There exists a unique local coordinate z_1 in the jet $[z_1]$ such that $E = z_1^{-n_1}$ exactly. However, at the other points P_α the Abelian integral E does not have a selected branch, so we cannot choose a representative of the jet $[z_\alpha]$ in this way.

Let E' and E'' denote two branches of the Abelian integral E at a point P_α . We have $E'' = E' + C$, where the constant C is real if the integral E is real-normalized. Let z'_α and z''_α denote the unique representatives of the jet $[z_\alpha]$ such that $E' = (z'_\alpha)^{-n_\alpha}$ and $E'' = (z''_\alpha)^{-n_\alpha}$ near P_α . It is easy to check that z'_α and z''_α are related as follows:

$$z''_\alpha = z'_\alpha + \sum_{j=1}^{\infty} C_j (z'_\alpha)^{jn_\alpha + 1}, \quad (3.3)$$

where there the constants C_j are polynomials in C with rational coefficients.

Now let Q be another Abelian integral on Γ . At the point P_1 it can be expanded in terms of the chosen local coordinate z_1 :

$$Q = \sum_{j=0}^{m_1} q_{1,j} z_1^{-j} + R_1^Q \log z_1 + O(z_1) \text{ near } P_1. \quad (3.4)$$

To expand Q at the other points P_α , we need to choose local coordinates at those points. Let z_α denote a choice of local coordinate in the jet $[z_\alpha]$ corresponding to some branch of E . Then Q can be expanded at P_α :

$$Q = \sum_{j=1}^{m_\alpha} q_{\alpha,j} z_\alpha^{-j} + R_\alpha^Q \log z_\alpha + O(1) \text{ near } P_\alpha, \quad \alpha = 2, \dots, N. \quad (3.5)$$

If we now assume that $m_\alpha \leq n_\alpha$, then changing the representative of $[z_\alpha]$ according to (3.3) does not affect the principal terms in the expansion of Q . Therefore, if $m_\alpha \leq n_\alpha$ for $\alpha = 2, \dots, N$, then the map $\mathcal{M}_{g,N}^m(n) \rightarrow \mathcal{M}_{g,N}(n)$ is a trivial fiber bundle with fiber $\mathbb{C}^{m_1+1} \oplus \mathbb{C}^{m_2} \oplus \dots \oplus \mathbb{C}^{m_N} \oplus \mathbb{R}^{N-1}$. To define an embedding $\mathcal{M}_{g,N}(\mathbf{n})$ into $\mathcal{M}_{g,N}^m(\mathbf{n})$, we need to choose polynomials $q_\alpha(t)$ of degrees m_α , such that $q_\alpha(t)$ have no constant terms for $\alpha = 2, \dots, N$, and N imaginary numbers R_α^Q adding up to zero. If we do not make the assumption that $m_\alpha \leq n_\alpha$ for $\alpha = 2, \dots, N$, then the map is only locally trivial, and we cannot define a global embedding $\mathcal{M}_{g,N}(\mathbf{n}) \hookrightarrow \mathcal{M}_{g,N}^m(\mathbf{n})$.

We now define a collection of functions on $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$. First, we consider the residues of the differential dQ :

$$R_\alpha^Q = \text{Res}_{P_\alpha} dQ, \quad \alpha = 1, \dots, N. \quad (3.6)$$

These functions satisfy the relation $\sum R_\alpha^Q = 0$.

The remaining functions are only locally well-defined on $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$. Fix a basis of cycles A_i, B_i in $H_1(\Gamma, \mathbb{Z})$ with canonical intersection form, such that no cycle passes through any of the P_i or the zero-divisors of dE and dQ . Let $\bar{\Gamma}$ denote the Riemann surface cut along these cycles and let $\partial\Gamma$ denote the boundary of $\bar{\Gamma}$. Fix also a choice of paths $\gamma_2, \dots, \gamma_n$ from P_1 to P_2, \dots, P_n not intersecting any of the basis cycles. The fixed branches of E and Q then extend to functions on $\bar{\Gamma} \setminus \cup \gamma_i$.

Let z_α be the unique local coordinate in the jet $[z_\alpha]$ such that for the chosen branch of E we have $E = z_\alpha^{-n_\alpha}$ near P_α . We consider the coefficients of the expansion of $Q dE$ at the marked points, with respect to the chosen local coordinates z_α :

$$Q dE = \left(\sum_{k=0}^{n_\alpha+m_\alpha} T_{\alpha,k} z_\alpha^{-k-1} - n_\alpha R_\alpha^Q z^{-n_\alpha-1} \log z_\alpha + \sum_{j=1}^{\infty} H_{\alpha,j} z_\alpha^{j-1} \right) dz_\alpha. \quad (3.7)$$

We also consider the periods of the differentials dE and dQ along the chosen cycles:

$$\tau_{A_i}^E = \oint_{A_i} dE, \quad \tau_{B_i}^E = \oint_{B_i} dE, \quad \tau_{A_i}^Q = \oint_{A_i} dQ, \quad \tau_{B_i}^Q = \oint_{B_i} dQ, \quad i = 1, \dots, g, \quad (3.8)$$

and the A -periods of the differential $Q dE$:

$$a_i = \oint_{A_i} Q dE, \quad i = 1, \dots, g. \quad (3.9)$$

The sum of the residues of the differential $Q dE$ is equal to its integral over $\partial\Gamma$. That gives the only relation between these functions:

$$2\pi i \sum_{\alpha=1}^N T_{\alpha,0} = \sum_{i=1}^g \left(\tau_{A,i}^Q \tau_{B,i}^E - \tau_{B,i}^Q \tau_{A,i}^E \right), \quad (3.10)$$

so therefore we exclude $T_{1,0}$ from consideration.

As before, these functions are locally well defined on $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$ and have well-defined differentials, and they can be used as holomorphic coordinates on $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$. The corresponding generalization of Theorem 2.1 was also proved in [14]:

Theorem 3.1 *Let $P = (\Gamma, P_\alpha, [z_\alpha], E, Q)$ be a point on $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$ such that the zero-divisors of dE and dQ on Γ do not intersect. Then in a neighborhood of P the collection of $5g + 2N - 2 + \sum(n_\alpha + m_\alpha)$ functions $R_\alpha^Q, T_{\alpha,k}$ (excluding R_1^Q and*

$T_{1,0}$), $\tau_{A_i}^E$, $\tau_{B_i}^E$, $\tau_{A_i}^Q$, $\tau_{B_i}^Q$ and a_i have linearly independent differentials and thus define a local holomorphic coordinate system for $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$.

In the N -point case, the local coordinates z_α and hence the functions $H_{\alpha,k}$ are no longer globally well-defined on $\mathbf{M}_{g,N}(\mathbf{n}, \mathbf{m})$ for $\alpha = 2, \dots, N$. Hence, we cannot talk of their critical points. However, on the smaller subspace $\mathcal{M}_{g,N}^{\mathbf{m}}(\mathbf{n})$ the imaginary parts of these functions are well-defined. Indeed, any two choices of local coordinate z_α in the jet $[z_\alpha]$ are related by Eq. (3.3). On $\mathcal{M}_{g,N}^{\mathbf{m}}(\mathbf{n})$ the coefficients C_j are real, and therefore substituting them into expansion (3.7) does not change the imaginary parts of the functions $H_{\alpha,k}$.

It is now clear how to generalize the results of the previous section to the N -point case. Fix N polynomials $q_\alpha(t)$ of degrees m_α with no constant terms for $\alpha = 2, \dots, N$, and N imaginary numbers R_α^Q adding up to zero. Consider the embedding $\mathcal{M}_{g,N}(\mathbf{n}) \hookrightarrow \mathcal{M}_{g,N}^{\mathbf{m}}(\mathbf{n})$ defined by these data.

We introduce a hierarchy of foliations on $\mathcal{M}_{g,N}(\mathbf{n})$. Let $\Lambda' \subset \mathcal{M}_{g,N}(\mathbf{n})$ be a submanifold along which the functions $T_{\alpha,k}$ for $k = 1, \dots, m_\alpha + n_\alpha$ are constant:

$$\delta T_{\alpha,k} = 0 \quad \forall k > 0 \quad \text{and} \quad \forall \delta \in T(\Lambda'). \quad (3.11)$$

Let $\Lambda \subset \Lambda'$ denote a submanifold along which the functions $T_{\alpha,k}$ for all $k = 0, \dots, m_\alpha + n_\alpha$ are constant:

$$\delta T_{\alpha,k} = 0 \quad \forall k \geq 0 \quad \text{and} \quad \forall \delta \in T(\Lambda). \quad (3.12)$$

Let $\mathcal{L}_E \subset \Lambda$ denote a submanifold along which the periods of dE are constant:

$$\delta \oint_{\gamma} dE = 0 \quad \forall \delta \in T(\mathcal{L}_E) \quad \text{and} \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}), \quad (3.13)$$

let $\mathcal{L}_Q \subset \Lambda$ denote a submanifold along which the periods of dQ are constant:

$$\delta \oint_{\gamma} dQ = 0 \quad \forall \delta \in T(\mathcal{L}_Q) \quad \text{and} \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (3.14)$$

Finally, let $\mathcal{L} = \mathcal{L}_E \cap \mathcal{L}_Q$ denote a submanifold along which the periods of dE and dQ are both constant:

$$\delta \oint_{\gamma} dE = \delta \oint_{\gamma} dQ = 0 \quad \forall \delta \in T(\mathcal{L}) \quad \text{and} \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (3.15)$$

We now define a real-valued function on $\mathcal{M}_{g,N}(\mathbf{n})$ and study its critical points on the leaves of the foliations described above. Let $d_{\alpha,j}$ for $\alpha = 1, \dots, N$, $j = 1, \dots, M_\alpha$ be a finite collection of real constants. As we have noted above, the function

$$\mathcal{H} = \sum_{\alpha,j} d_{\alpha,j} \operatorname{Im} H_{\alpha,j} \quad (3.16)$$

is globally well-defined on $\mathcal{M}_{g,N}(\mathbf{n})$. The critical points of this function on the leaves of the foliations described above are then described by the following theorem, which is the N -point generalization of Theorem 3.1.

Theorem 3.2 *Let $q_\alpha(t)$ be polynomials of degrees m_α , such that $q_\alpha(t)$ does not have a constant term for $\alpha = 2, \dots, N$. Let R_α^Q be imaginary numbers adding up to zero. Let $(\Gamma, P_\alpha, [z_\alpha])$ be a point in $\mathcal{M}_{g,1}(\mathbf{n})$, which we assume to be embedded in $\mathcal{M}_{g,1}^m(n)$ via the data $q_\alpha(z)$ and R_α^Q . Let E and Q be the real-normalized Abelian integrals defined by Eqs. (3.2), (3.4) and (3.5), and let D denote the intersection of the zero-divisors of dE and dQ .*

A. *The point $(\Gamma, P_\alpha, [z_\alpha])$ is a critical point of \mathcal{H} restricted to \mathcal{L} if and only if*

- (i) *There exists a meromorphic function Y on Γ with poles at P_α with principal parts*

$$Y = i \sum_{j=1}^{M_1} d_{1,j} z_1^{-j} + O(z_1) \text{ near } P_1, \quad Y = i \sum_{j=1}^{M_\alpha} d_{\alpha,j} z_\alpha^{-j} + O(1) \text{ near } P_\alpha, \\ \alpha = 2, \dots, N, \quad (3.17)$$

and poles of order μ_s at the common zeros Q_s of the differentials dE and dQ on Γ , where μ_s is the multiplicity of the common zero Q_s , and no other singularities.

B. *The point $(\Gamma, P_\alpha, [z_\alpha])$ is a critical point of \mathcal{H} restricted to \mathcal{L}_E if and only if there exists a function satisfying condition (i) and in addition the following condition:*

- (ii) *the differential $Y dE$ is real-normalized, i.e.*

$$\operatorname{Im} \oint_\gamma Y dE = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (3.18)$$

C. *The point $(\Gamma, P_\alpha, [z_\alpha])$ is a critical point of \mathcal{H} restricted to \mathcal{L}_Q if and only if there exists a function satisfying condition (i) and in addition the following condition:*

- (iii) *the differential $Y dQ$ is real-normalized, i.e.*

$$\operatorname{Im} \oint_\gamma Y dQ = 0 \quad \forall \gamma \in H_1(\Gamma, \mathbb{Z}). \quad (3.19)$$

D. *The point $(\Gamma, P_\alpha, [z_\alpha])$ is a critical point of \mathcal{H} restricted to Λ if and only if there exists a function satisfying conditions (i), (ii) and (iii).*

E. *The point $(\Gamma, P_\alpha, [z_\alpha])$ is a critical point of \mathcal{H} restricted to Λ' if and only if there exists a function satisfying conditions (i), (ii) and (iii), and such that near P_α we have*

$$Y = i \sum_{j=1}^{M_\alpha} d_{\alpha,j} z_\alpha^{-j} + O(z_\alpha) \text{ near } P_\alpha. \quad (3.20)$$

The proof of the theorem is similar to the one-point case. We introduce an extension of the homomorphism (2.21) to the N -point case:

$$f : T_P(\Lambda') \rightarrow \Omega(\Gamma, P_\alpha, E, Q). \quad (3.21)$$

For a tangent vector $\delta \in T_P(\Lambda')$, the differential $f(\delta) = \Omega_\delta = (\delta Q|_E)dE$ has simple poles at P_1, \dots, P_N and jumps along the cuts that are real linear combinations of dE and dQ . The space of such differentials, which we denote by $\Omega(\Gamma, P_\alpha, E, Q)$, has real dimension $6g + 2N - 2$. The image of the tangent subspace $T(\Lambda)$ is the subspace $\Omega(\Gamma, P_1, E, Q)$ of differentials having poles only at the point P_1 .

The corresponding generalization of Proposition 2.2 is

Proposition 3.1 *Let $D = \sum \mu_s Q_s$ denote the intersection of the zero divisors of dE and dQ . Then the image of the tangent space $T_P(\Lambda')$ under the homomorphism f of (3.21) is the subspace $\Omega_D(\Gamma, P_\alpha, E, Q) \subset \Omega(\Gamma, P_\alpha, E, Q)$ of differentials having zeroes at Q_s of orders μ_s . In particular, if the zero divisors of dE and dQ do not intersect, then the map (3.21) is an isomorphism.*

The space $\Omega(\Gamma, P_\alpha, E, Q)$ is spanned over \mathbb{R} by the $6g$ differentials defined by conditions (2.25)–(2.30), and in addition by the following differentials:

- $2N - 2$ single-valued meromorphic differentials $d\Omega_\alpha^R, d\Omega_\alpha^I, \alpha = 2, \dots, N$, which have real periods and have poles at P_α and P_1 with residues ± 1 and $\pm i$, respectively:

$$\text{Res}_{P_\alpha} d\Omega_\alpha^R = 1, \quad \text{Res}_{P_1} d\Omega_\alpha^R = -1, \quad \text{Im} \oint_{\gamma} d\Omega_\alpha^R = 0, \quad \gamma = A_i, B_i, \quad (3.22)$$

$$\text{Res}_{P_\alpha} d\Omega_\alpha^I = i, \quad \text{Res}_{P_1} d\Omega_\alpha^I = -i, \quad \text{Im} \oint_{\gamma} d\Omega_\alpha^I = 0 \quad \gamma = A_i, B_i. \quad (3.23)$$

Proof of Theorem 3.2 Let dY be the unique real-normalized Abelian differential with fixed branch at P_1 such that Y has poles of the form (3.17). Fix a real number $\rho > 0$, and let γ_α denote a path of radius ρ around the point P_α in the chosen local coordinate z_α . Let $\delta \in T(\Lambda')$ be a tangent vector. A direct calculation then shows that

$$\begin{aligned} \delta H &= \text{Re} \left(\sum_{\alpha=2}^N d_{\alpha,0} \delta T_{\alpha,0} \right) - \sum_{\alpha=1}^N \text{Re} \text{Res}_{P_\alpha} Y \delta Q dE \\ &= \text{Re} \left(\sum_{\alpha=1}^N d_{\alpha,0} \delta T_{\alpha,0} \right) - \frac{1}{2\pi} \text{Im} \oint_{\partial\Gamma} Y \Omega_\delta, \end{aligned} \quad (3.24)$$

where $d_{\alpha,0}$ denotes the constant term of Y at P_α . For a tangent vector δ in $T(\Lambda)$ or in any smaller tangent subspace, we have $\delta T_{\alpha,0} = 0$. Therefore, we can use the same reasoning as in Theorem 2.2, which proves parts A through D.

Finally, if P is a critical point of \mathcal{H} restricted Λ' , then $\delta \mathcal{H} = 0$ for all variations of $\delta T_{0,\alpha}$, which can only happen if $d_{0,\alpha} = 0$ for all $\alpha = 2, \dots, N$, which proves part E.

Acknowledgments This work was supported by The Ministry of Education and Science of the Russian Federation (contract 02.740.11.5194).

References

1. Novikov, S.P.: A periodic problem for the Korteweg–de Vries equation. *Funct. Anal. Appl.* **8**(3), 54–66 (1974)
2. Dubrovin, B., Matveev, V., Novikov, S.: Non-linear equations of Korteweg–de Vries type, finite zone linear operators and Abelian varieties. *Uspekhi Mat. Nauk* **31**(1), 55–136 (1976)
3. Krichever, I.M.: The spectral theory of “finite-gap” nonstationary Schrödinger operators, The non-stationary Peierls model. (Russian) *Funct. Anal. Appl.* **20**(3), 42–54, 96 (1986)
4. Boutroux, P.: Recherches sur les transcendentes de M Painlevé et l’etude asymptotique des équations différentielles du second ordre. *Ann Sci Ecol. Norm Supér* **30**, 255–376 (1913), **31**, 99–159 (1914)
5. Krichever, I.M.: On Heisenberg relations for the ordinary linear differential operators. IHES Preprint, Bur-Sur-Yvette (1990)
6. Fucito, F., Gamba, A., Martellini, M., Ragnisco, O.: Non-linear WKB analysis of the string equation. *Int. J. Mod. Phys. B* **6**(11–12), 2123–2147 (1992)
7. Grinevich, P.G., Novikov, S.P.: String equation—2. Physical solution. *St. Petersburg Math. J* **6**, 553 (1995)
8. Kapaev, A.A.: Monodromy approach to the scaling limits in the isomonodromy systems. arXiv.org:nlin/0211022
9. David, F.: Phases of the large-N matrix model and non-perturbative effects in 2D gravity. *Nuclear Phys. B* **348**(4), 507–524 (1991)
10. David, F.: Non-perturbative effects in matrix models and vacua of two dimensional gravity. *Phys. Lett. B* **302**(4), 403–410 (1993)
11. Bertola, M., Mo, M.Y.: Commuting difference operators, spinor bundles and the asymptotics of orthogonal polynomials with respect to varying complex weights (2006). arXiv.org:math-ph/0605043
12. Lee, S.Y.: Teodorescu, R., Wiegmann, P., Shocks and finite-time singularities in Hele–Shaw flow. *Phys D Nonlinear Phenom.* **238**(14), 1113–1128
13. Krichever, I.: The τ -function of the universal Whitham hierarchy, matrix models, and topological field theories. *Commun. Pure Appl. Math.* **47**, 437–475 (1994)
14. Krichever, I., Phong, D.H.: On the integrable geometry of $N = 2$ supersymmetric gauge theories and soliton equations. *J. Differ. Geom.* **45**, 445–485 (1997)
15. Krichever, I., Phong, D.H.: Symplectic forms in the theory of solitons, Surveys in Differential Geometry IV. In: Terng, C.L., Uhlenbeck, K. (eds.) International Press, Boston, pp. 239–313 (1998)
16. Grushevsky, S., Krichever, I.: The universal Whitham hierarchy and the geometry of the moduli space of pointed Riemann surfaces. arXiv:0810.2139
17. Grushevsky S., Krichever, I.: Vanishing of tautological classes on the moduli space of curves (in preparation)
18. Dzhamay, A.: Real-normalized Whitham hierarchies and the WDVV equations. *Int. Math. Res. Notices* (21), 1103–1130 (2000)