

The orbit method, microlocal analysis and applications to L -functions

Paul Nelson

ETH Zurich

(partly joint work with Akshay Venkatesh)

Automorphic Forms and Arithmetic Seminar
Columbia University

Objectives

- ▶ Analyze L -functions on $GL_{n+1} \times GL_n$ and related groups.
- ▶ Determine asymptotics of “special functions” attached to representations of such groups.
- ▶ $n = 1$: Bernstein–Reznikov, Michel–Venkatesh, (...): analysis via explicit formulas, complicated in higher rank.
- ▶ Develop the orbit method in the spirit of microlocal analysis, giving a soft approach that applies in higher rank.
- ▶ Standard problems (moments, subconvexity).

Contents

Special values of L -functions as branching coefficients

The orbit method

Microlocal analysis

Local applications

Global applications

Special values of L -functions as branching coefficients

The orbit method

Microlocal analysis

Local applications

Global applications

Setting

- ▶ Let (G, H) be one of

$$(\mathrm{GL}_{n+1}(\mathbb{R}), \mathrm{GL}_n(\mathbb{R})),$$

$$(\mathrm{U}(p+1, q), \mathrm{U}(p, q)), \quad (\mathrm{SO}(p+1, q), \mathrm{SO}(p, q)).$$

(G, H) is a **strong Gelfand pair**: irreducible representations of G have multiplicity-free restriction to H .

- ▶ Let $\Gamma \leq G$ and $\Gamma_H = \Gamma \cap H \leq H$ be arithmetic lattices.
Example: $(\Gamma, \Gamma_H) = (\mathrm{GL}_{n+1}(\mathbb{Z}), \mathrm{GL}_n(\mathbb{Z}))$.

- ▶ Define

$$[G] := \Gamma \backslash G, \quad [H] := \Gamma_H \backslash H.$$

We assume these quotients are compact.

- ▶ We consider irreducible subrepresentations

$$\pi \subseteq L^2([G]), \quad \sigma \subseteq L^2([H]).$$

Global restriction

Let $v \in \pi \subseteq L^2([G])$ be a smooth vector.

Restrict to $[H]$, spectrally decompose:

$$v|_{[H]} = \sum_{\sigma \subseteq L^2([H])} \underbrace{\sum_{u \in \mathcal{B}(\sigma)} \left(\int_{[H]} v \bar{u} \right) u}_{=: \text{global projection of } v \text{ to } \sigma}$$

Local restriction

Example

$(G, H) = (\mathrm{SO}(3), \mathrm{SO}(2))$, π : irrep of highest weight $T \in \mathbb{Z}_{\geq 0}$.

Weight space decomposition:

$$\pi|_H = \bigoplus_{\ell=-T}^T \sigma_\ell, \quad v = \sum v_\ell, \quad \|v\|^2 = \sum \|v_\ell\|^2.$$

General picture

As abstract unitary representations of H ,

$$\pi|_H = \int_{\sigma \in \mathrm{Irr}(H)} \underbrace{m_\pi(\sigma)}_{\in \{0,1\}} \sigma \, d\sigma, \quad d\sigma = \text{Plancherel}.$$

This means: there are “local” projection maps $\pi \rightarrow \sigma$ such that

$$\|v\|^2 = \int_{\sigma} m_\pi(\sigma) \|\text{local projection of } v \text{ to } \sigma\|^2 \, d\sigma.$$

Branching coefficients

The strong Gelfand property implies

global projection = scalar multiple of local projection.

We may thus define a **branching coefficient** $\mathcal{L}(\pi, \sigma) \in \mathbb{R}_{\geq 0}$ by asking that for all $v \in \pi$,

$$\begin{aligned} \|\text{global projection of } v \text{ to } \sigma\|^2 = \\ \mathcal{L}(\pi, \sigma) \|\text{local projection of } v \text{ to } \sigma\|^2. \end{aligned}$$

It quantifies how automorphic forms in π correlate with those in σ .

Branching coefficients and L -values

Refined Gan–Gross–Prasad conjecture, known in many cases:¹

$$\mathcal{L}(\pi, \sigma) = \text{special value of an } L\text{-function.}$$

Example

In the $\mathrm{GL}_n(\mathbb{Z})$ case,²

$$\mathcal{L}(\pi, \sigma) \approx \frac{1}{4} \left| \frac{L(\pi \times \bar{\sigma}, 1/2)}{L(\mathrm{Ad}, \dots, 1)} \right|^2.$$

¹Hecke, Waldspurger, Ichino–Ikeda, N. Harris, Jacquet–Rallis, W. Zhang, Beuzart-Plessis, Beuzart-Plessis–Liu–Zhang–Zhu, Beuzart-Plessis–Chaudouard–Zydor

²Jacquet–Piatetski-Shapiro–Shalika

Motivation for studying L -values

Problem: estimate special values of L -functions,

- ▶ individually (“subconvexity problem”) or
- ▶ on average over a family (“moment problem”).

Considered fundamental nowadays.

Motivated by questions/applications discovered in the 1980's and 1990's:

- ▶ random matrix heuristics for moments of families
- ▶ existence and distribution of integral solutions of $Q(v) = n$ for a ternary quadratic form Q
- ▶ arithmetic quantum unique ergodicity: $|\varphi_j|^2 d\mu \sim (?)$

Problems

To study $\mathcal{L}(\pi, \sigma)$ using the defining property

$$\begin{aligned} & \|\text{global projection of } v \text{ to } \sigma\|^2 = \\ & \mathcal{L}(\pi, \sigma) \|\text{local projection of } v \text{ to } \sigma\|^2 \end{aligned}$$

we must understand the local and global projections.

Sample local problem

Let $\mathcal{F} \subseteq \text{Irr}(H)$ be a nice family.

Can we construct a nice vector $v \in \pi$ so that the map

$$\text{Irr}(H) \ni \sigma \mapsto \|\text{local projection of } v \text{ to } \sigma\|^2$$

approximates the characteristic function of \mathcal{F} ?

Can we then estimate the matrix coefficients $\langle gv, v \rangle$?

Special values of L -functions as branching coefficients

The orbit method

Microlocal analysis

Local applications

Global applications


G : Lie group, $\mathfrak{g} = \text{Lie}(G)$, $\mathfrak{g}^\wedge = \text{Hom}(\mathfrak{g}, i\mathbb{R})$

Definition

A **coadjoint orbit** is $\mathcal{O} \subseteq \mathfrak{g}^\wedge$ is a G -orbit for the coadjoint action.

Example

$G = \text{SO}(3)$, $\mathfrak{g}, \mathfrak{g}^\wedge \cong \mathbb{R}^3$, $\{\text{coadjoint orbits}\} = \{\text{spheres}\}$.



Theorem (Kirillov)

Lie algebra structure on \mathfrak{g} defines a symplectic structure on \mathcal{O} , hence a symplectic **volume form** ω .

Orbit method: heuristic based on an (approximate) bijection

$$\text{Irr}(G) \approx \{\text{coadjoint orbits}\}$$

$$\pi \longleftrightarrow \mathcal{O}_\pi$$

compatible with natural operations.

\mathcal{O}_π should describe the **character** χ_π : for small $x \in \mathfrak{g}$,

$$\chi_\pi(\exp(x)) = j^{-1/2}(x) \int_{\xi \in \mathcal{O}_\pi} e^{\langle x, \xi \rangle} d\omega(\xi),$$

where $j = \text{Jac}(\exp : \mathfrak{g} \rightarrow G)$, $j(0) = 1$.

Such an identity is called the **Kirillov formula** for π .

Valid for

- ▶ G nilpotent (Kirillov)
- ▶ G compact (Weyl, Kirillov)
- ▶ G reductive, π tempered (Rossmann)

Example

$G = \mathrm{SO}(3)$. Classify by highest weight:

$$\mathrm{Irr}(G) = \{\pi_T : T \in \mathbb{Z}_{\geq 0}\}$$

Kirillov formula for $\pi = \pi_T$ holds with

$$\mathcal{O}_\pi = \text{sphere of radius } T + \frac{1}{2},$$

$$\mathrm{vol}(\mathcal{O}_\pi, d\omega) = 2T + 1 = \dim(\pi).$$

Example

$$G = \mathrm{PGL}_2(\mathbb{R}) \cong \mathrm{SO}(1, 2)$$

$$\mathfrak{g}^\wedge \ni i \begin{pmatrix} x & y+z \\ y-z & -x \end{pmatrix}$$

- ▶ one-sheeted hyperboloid

$$\mathcal{O}^+(r) = \{x^2 + y^2 - z^2 = r^2\}$$


- ▶ two-sheeted hyperboloid

$$\mathcal{O}^-(k) = \{x^2 + y^2 - z^2 = -k^2\}$$

Tempered irreducible representations:

- ▶ principal series $\pi(r, \varepsilon)$
- ▶ discrete series $\pi(k)$ for $k \in \mathbb{Z}_{\geq 1}$

$$\mathcal{O}_{\pi(r, \varepsilon)} = \mathcal{O}^+(r), \quad \mathcal{O}_{\pi(k)} = \mathcal{O}^-(k - 1/2)$$



Special values of L -functions as branching coefficients

The orbit method

Microlocal analysis

Local applications

Global applications

Microlocalization

Heuristic. For each partition of $\mathcal{O}_\pi = \sqcup \mathcal{P}_j$ with $\text{vol}(\mathcal{P}_j) = 1$, there corresponds an orthonormal basis $(v_j)_j$ of π such that for $x \in \mathfrak{g}$,


$$\pi(x)v_j \approx \langle x, \mathcal{P}_j \rangle v_j$$

provided that $\langle x, \mathcal{P}_j \rangle$ is approximately well-defined.

Example

$G = \text{SO}(3)$, π of highest weight T . The decomposition $\pi|_{\text{SO}(2)} = \bigoplus_{\ell=-T}^T \sigma_\ell$ into one-dimensional weight spaces corresponds to the partition of \mathcal{O}_π into horizontal strips

$$\mathcal{P}_\ell = \left\{ \xi \in \mathcal{O}_\pi : (\text{z-coordinate of } \xi) \in \left[\ell - \frac{1}{2}, \ell + \frac{1}{2} \right] \right\}$$



For our purposes, better to take each \mathcal{P} concentrated near some specific $\tau \in \mathcal{O}_\pi$, say $\mathcal{P} = \mathcal{P}_\tau$. We call the corresponding vector v_τ **microlocalized** at τ and \mathcal{P}_τ its **microlocal support**.

Example

$G = \text{SO}(3)$, $\pi = \pi_T$ as above,

$$\mathcal{P}_\tau \approx \mathcal{O}_\pi \cap (\tau + \mathcal{O}(T^{1/2}))$$

for a well-spaced subset $\{\tau\} \subseteq \mathcal{O}_\pi$ of cardinality $\dim(\pi) = 2T + 1$.





Figure: Microlocalized partitions $\mathcal{O}_\pi = \sqcup \mathcal{P}_\tau$ for $T \approx 40,160$

A key feature of microlocalized vectors v_τ is that their matrix coefficients concentrate near the centralizer G_τ :

$$\langle gv_\tau, v_\tau \rangle \approx 0 \text{ unless } g\tau \approx \tau.$$

Informal reason:

- ▶ gv_τ is microlocalized at $g\tau$
- ▶ vectors with disjoint microlocal supports are orthogonal



Microlocal calculus

- ▶ We work with microlocalized vectors implicitly via operators

$$\pi(f) = \int_{g \in G} f(g)\pi(g) dg$$

attached to $f \in C_c^\infty(G)$ supported near the identity.

- ▶ We describe f by pulling it back to the Lie algebra and taking the Fourier transform. We call the result $a \in \mathcal{S}(\mathfrak{g}^\wedge)$, and write

$$\pi(f) =: \text{Op}(a).$$

- ▶ Kirillov formula (ignoring $j^{-1/2}$):

$$\text{trace}(\text{Op}(a)) \approx \int_{\mathcal{O}_\pi} a d\omega.$$

Take a microlocalized basis

$$\pi = \bigoplus \mathbb{C}v_\tau \quad \longleftrightarrow \quad \mathcal{O}_\pi = \sqcup \mathcal{P}_\tau.$$

The orbit method suggests

$$\text{Op}(a)v_\tau \approx a(\mathcal{P}_\tau)v_\tau \approx a(\tau)v_\tau$$

if $a(\mathcal{P}_\tau)$ is approximately well-defined, i.e., a is essentially constant on \mathcal{P}_τ .

Example

Take $G = \text{SO}(3)$, $\pi = \pi_T$ and, as above,

$$\mathcal{P}_\tau \approx \mathcal{O}_\pi \cap (\tau + \mathcal{O}(T^{1/2})).$$

If

$$\partial^\alpha a(\xi) \ll_\alpha T^{(-1/2-\varepsilon)|\alpha|}.$$

then a varies by $\ll T^{-\varepsilon}$ on each \mathcal{P}_τ .

Theorem (NV 2018, N 2020)

Let $\pi = \pi_T$ be a unitary representation of a Lie group G . The assignment $a \mapsto \text{Op}(a)$, restricted to functions $a = a_T$ satisfying estimates as above, enjoys a reasonable microlocal calculus, e.g.:

1. If a is supported on elements of size $\asymp T$ and satisfies

$$\partial^\alpha a(\xi) \ll_\alpha T^{(-1/2-\varepsilon)|\alpha|},$$

then $\text{Op}(a)$ has operator norm $O(1)$.

2. If G is reductive and π is irreducible, then $\text{Op}(a)$ has trace norm $O(T^d)$, where $2d = \dim(\mathcal{O}_\pi) = \dim(G) - \text{rank}(G)$.
3. $\text{Op}(a)\text{Op}(b) = \text{Op}(a \star b)$, where $a \star b \sim ab$.
4. Similar results for polynomials, functions with less regularity transverse to the coadjoint orbits, etc.

Constructing microlocalized vectors using the calculus

Say

$$\pi \hookrightarrow L^2([G]), \quad [G] = \Gamma \backslash G.$$

For $a = a_\tau$ as in the microlocal calculus, consider

$$\sum_{v \in \mathcal{B}(\pi)} \text{Op}(a)v(x) \cdot \overline{v(y)}.$$

Example

If $a|_{\mathcal{O}_\pi} \approx$ characteristic function of \mathcal{P}_τ , then we get

$$\approx v_\tau(x) \overline{v_\tau(y)}.$$

Special values of L -functions as branching coefficients

The orbit method

Microlocal analysis

Local applications

Global applications

Relative characters

For tempered (π, σ) , we consider the hermitian form

$$\mathcal{H}_\sigma : \pi \otimes \bar{\pi} \rightarrow \mathbb{C}$$

$$\mathcal{H}_\sigma(v \otimes v) = \|\text{local projection of } v \text{ to } \sigma\|^2.$$

For a nice function a on \mathfrak{g}^\wedge , we define the **relative character**

$$\mathcal{H}_{\pi, \sigma}(a) := \sum_{v \in \mathcal{B}(\pi)} \mathcal{H}_\sigma(\text{Op}(a)v \otimes v).$$

Example

If $a|_{\mathcal{O}_\pi} \approx$ characteristic function of \mathcal{P}_τ , then

$$\mathcal{H}_{\pi, \sigma}(a) \approx \|\text{local projection of } v_\tau \text{ to } \sigma\|^2.$$

Question

For $a = a_\tau$ as in the microlocal calculus, what is the asymptotic behavior of $\mathcal{H}_{\pi, \sigma}(a)$? Here $(\pi, \sigma) = (\pi_\tau, \sigma_\tau)$.

If $a \rightsquigarrow f \in C_c^\infty(G)$ with $\text{Op}(a) = \pi(f)$, then one can verify that

$$\mathcal{H}_{\pi,\sigma}(a) = \int_{h \in H} \underbrace{\text{trace}(\pi(h)\pi(f))}_{\int_{g \in G} \chi_\pi(hg)f(g) dg} \overline{\chi_\sigma(h)} dh.$$

Heuristic

Pretend \exp were an isomorphism, apply Kirillov formula:

$$\begin{aligned} \mathcal{H}_{\pi,\sigma}(a) &\approx \int_{x \in \mathfrak{g}} \int_{y \in \mathfrak{h}} \chi_\pi(e^{y+x}) a^\vee(x) \overline{\chi_\sigma(e^y)} dx dy \\ &= \int_{\mathcal{O}_\pi \cap \text{preimage}(\mathcal{O}_\sigma)} a? \end{aligned}$$

Stability

We say that (π, σ) is **stable** if

$$\mathcal{O}_{\pi, \sigma} := \mathcal{O}_{\pi} \cap \text{preimage}(\mathcal{O}_{\sigma})$$



is an H -torsor (closed orbit, trivial stabilizer).

Fact

This is a generic condition, depending only upon the infinitesimal characters $(\lambda_{\pi}, \lambda_{\sigma})$, equivalent to

$$\{\text{eigenvalues of } \lambda_{\pi}\} \cap \{\text{eigenvalues of } \lambda_{\sigma}\} = \emptyset.$$

Some $\mathcal{O}_{\pi, \sigma}$'s for $(\text{SO}(3), \text{SO}(2))$ and $(\text{PGL}_2(\mathbb{R}), \text{GL}_1(\mathbb{R}))$:




Relative character asymptotics in the stable case

Theorem (NV 2018)

Let $(\pi, \sigma) = (\pi_T, \sigma_T)$ be tempered and “uniformly stable”:
 $T^{-1}(\lambda_\pi, \lambda_\sigma)$ lies in a fixed compact collection of stable pairs.
Then for $a = a_T$ as in the microlocal calculus,

$$\mathcal{H}_{\pi, \sigma}(a) \sim \int_{\mathcal{O}_{\pi, \sigma}} a,$$

where we integrate by transporting Haar on H to its torsor $\mathcal{O}_{\pi, \sigma}$.



Sketch of proof

Recall that

$$\mathcal{H}_{\pi,\sigma}(a) = \int_{h \in H} \text{trace}(\pi(h) \text{Op}(a)) \overline{\chi_{\sigma}(h)} dh.$$

- ▶ Choose a microlocalized partition $\mathcal{O}_{\pi} = \sqcup \mathcal{P}_{\tau}$.
Key case: a concentrated near some \mathcal{P}_{τ} with $\tau \in \mathcal{O}_{\pi,\sigma}$.
- ▶ $\text{trace}(\pi(h) \text{Op}(a)) \approx \langle h v_{\tau}, v_{\tau} \rangle = \text{negligible unless } h\tau \approx \tau$.
Happens only if h is small, since stability $\implies H_{\tau} = \{1\}$.
- ▶ For small h , the heuristic argument can be pushed through.

Special values of L -functions as branching coefficients

The orbit method

Microlocal analysis

Local applications

Global applications

Overview

1. (N–Venkatesh 2018) asymptotic formulas for first moments

$$\sum_{\sigma \in \mathcal{F}} \mathcal{L}(\pi, \sigma), \quad \mathcal{F} : \text{“long” family}$$

with π fixed. Proof uses Ratner theory.

2. (N 2020) subconvex bounds $\mathcal{L}(\pi, \sigma) \ll C(\pi, \sigma)^{1/4 - \delta_n}$ via asymptotic formulas for (amplified) first moments

$$\sum_{\pi \in \mathcal{F}} \mathcal{L}(\pi, \sigma), \quad \mathcal{F} : \text{“short” family.}$$

Proof uses (amplified) relative trace formula, linear algebra.

“long” $\sim [T, 2T]$, “short” $\sim [T, T + 1]$.

\mathcal{F} must be uniformly stable; equivalently, $C(\pi, \sigma) \asymp T^{2n(n+1)}$.

Role of relative character asymptotics

Both proofs use

- ▶ construction of microlocalized vectors via operator calculus,
- ▶ relative character asymptotics, and
- ▶ the following consequence of the definition of $\mathcal{L}(\pi, \sigma)$:

$$\sum_{v \in \mathcal{B}(\pi)} \sum_{u \in \mathcal{B}(\sigma)} \left(\int_{[H]} \text{Op}(a)v \cdot \bar{u} \right) \left(\int_{[H]} \bar{v} \cdot u \right) = \mathcal{L}(\pi, \sigma) \mathcal{H}_{\pi, \sigma}(a).$$

We construct $a = a_T$ so that

$$\mathcal{H}_{\pi, \sigma}(a) \sim \int_{\mathcal{O}_{\pi, \sigma}} a$$

approximates the characteristic function of the family \mathcal{F} . Then

$$\sum_{\sigma \in \mathcal{F}} \mathcal{L}(\pi, \sigma) \approx \sum_{v \in \mathcal{B}(\pi)} \int_{[H]} \text{Op}(a)v \cdot \bar{v},$$

$$\sum_{\pi \in \mathcal{F}} \mathcal{L}(\pi, \sigma) \approx \sum_{u \in \mathcal{B}(\sigma)} \int_{x, y \in [H]} \overline{u(x)u(y)} \sum_{\gamma \in \Gamma} f(x^{-1}\gamma y) dx dy$$

if $\text{Op}(a) = \pi(f)$.


Estimating averages over σ

$$\sum_{\sigma \in \mathcal{F}} \mathcal{L}(\pi, \sigma) \approx \int_{[H]} [a], \quad [a] := \sum_{v \in \mathcal{B}(\pi)} \text{Op}(a)v \cdot \bar{v}$$

Decompose into localized vectors:

$$[a] \approx \int_{\tau \in \mathcal{O}_\pi} a(\tau) |v_\tau|^2 d\omega(\tau).$$

For **fixed** π , since $\text{supp}(a) \rightarrow \infty$, we can replace \mathcal{O}_π by the nilcone:




Any weak limit of the $|v_\tau|^2$ then attains unipotent invariance, hence (by Ratner) equidistributes. In particular, $[a]$ equidistributes.

Estimating averages over π

$$\sum_{\pi \in \mathcal{F}} \mathcal{L}(\pi, \sigma) \approx \sum_{u \in \mathcal{B}(\sigma)} \int_{x, y \in [H]} \overline{u(x)} u(y) \sum_{\gamma \in \Gamma} f(x^{-1} \gamma y) dx dy$$

$\sum_{\gamma \in \Gamma_H} \rightsquigarrow$ main term $\asymp T^{n(n+1)/2}$; apply amplification method.
Remains to show that $\Gamma - \Gamma_H$ gives a smaller contribution.

We may take a supported near some $\tau \in \mathcal{O}_{\pi, \sigma}$ with $|\tau| \asymp T$:



Key case: u microlocalized at $\tau_H := (\text{restriction of } \tau) \in \mathfrak{h}^\wedge$,
otherwise $\int_{[H]} \text{Op}(a)v \cdot \bar{u} = \text{negligible}$.

- ▶ Microlocalization of u implies approximate equivariance under the centralizer H_{τ_H} of τ_H . Key problem: estimate

$$\max_{y \in H} \int_{x \in H} \int_{z \in H_{\tau_H}} |f(x^{-1}\gamma yz)| dz dx$$

- ▶ f concentrates on G_{τ} , so $f(x^{-1}\gamma yz)$ detects when $x\tau \approx \gamma yz\tau$.
- ▶ Key case: $y = 1$ and γ centralizes τ .
- ▶ Key problem: exhibit some transversality between the subvarieties $H\tau$ and $\gamma H_{\tau_H}\tau$ of \mathcal{O}_{π} .




Figure: The “toy case” $(G, H) = (U(2), U(1))$, in which $H_{\tau_H} = H$.

- ▶ Numerology:

$$\dim(H_{\tau_H}\tau) = n, \quad \dim(H\tau) = n^2, \quad \dim(\mathcal{O}_\pi) = n^2 + n.$$

Maybe $\gamma H_{\tau_H}\tau$ and $H\tau$ literally transverse for generic $\gamma \in \Gamma$?
Need just a bit of transversality, but for every $\gamma \notin \Gamma_H$.

- ▶ Key idea: suffices to establish transversality, but with $H_{\tau_H}\tau$ (n -dimensional) replaced by $Z_{H\tau}$ (1-dimensional).
- ▶ Passing to the Lie algebra, reduce to a linear algebra problem:

Theorem (N 2020)

If $\tau \in \mathfrak{g}^\wedge$ is H -stable, $\gamma \in \mathfrak{g}_\tau - \mathfrak{z}$ and $z \in \mathfrak{z}_H - \{0\}$, then

$$[\gamma, [z, \tau]] \notin [\mathfrak{h}, \tau].$$

Theorem (N 2020)

Let M_n denote the space of $n \times n$ complex matrices. Embed $M_n \hookrightarrow M_{n+1}$ via

$$a \mapsto \begin{pmatrix} a & 0 \\ 0 & 0 \end{pmatrix}.$$

Set $z := \text{diag}(1, \dots, 1, 0) \in M_n$.

Let $\tau \in M_{n+1}$. Write

$$\tau = \begin{pmatrix} \tau_H & b \\ c & d \end{pmatrix}.$$

with $\tau_H \in M_n$. Let $\gamma \in M_{n+1}$. Assume that

- ▶ no eigenvalue of τ is also an eigenvalue of τ_H ,
- ▶ $[\gamma, \tau] = 0$, and
- ▶ $[\gamma, [z, \tau]] = [y, \tau]$ for some $y \in M_n$.

Then γ is a scalar matrix.

Summary

- ▶ We develop the orbit method in analytic form as a microlocal calculus for Lie group representations, sharp up to ε 's.
- ▶ We apply that calculus to determine relative character asymptotics in the stable regime.
- ▶ We deduce moment estimates and subconvex bounds in higher rank via some additional (local and global) arguments.

Theorem (NV 2018)

Assume $[G], [H]$: compact.

Fix π : tempered, generic.

Let $T \rightarrow \infty$. Set

$$\mathcal{F}_T := \left\{ \sigma \subseteq L^2([H]) \left| \begin{array}{l} m_\pi(\sigma) = 1 \\ \lambda_\sigma \in T \cdot \Lambda \end{array} \right. \right\},$$

with Λ : nice fixed compact collection of infinitesimal characters with all eigenvalues nonzero.

Then

$$\sum_{\sigma \in \mathcal{F}_T} \mathcal{L}(\pi, \sigma) \sim \frac{|\mathcal{F}_T|}{\text{vol}(\Gamma \backslash G)}.$$

Remark

Translates to (average L -value) $\sim 2 \prod_p(\dots)$, as predicted via random matrix heuristics.

Theorem (N 2020)

Let $(G, H) = (U_{n+1}, U_n)$. Assume $[H]$: compact.

Let $T \rightarrow \infty$. Let $(\pi, \sigma) = (\pi_T, \sigma_T)$ be Hecke-irreducible and tempered, with $T^{-1}(\lambda_\pi, \lambda_\sigma)$ in a fixed compact stable collection.

Then

$$\mathcal{L}(\pi, \sigma) \ll T^{n(n+1)/2-\delta}$$

for some fixed $\delta = \delta_n > 0$.

Context

- ▶ Stability condition equivalent to “no conductor dropping”:

$$C(\pi, \sigma) \asymp T^{2n(n+1)}.$$

- ▶ Translates to a **subconvex bound** for the corresponding L -function:

$$\mathcal{L}(\pi, \sigma) \ll C(\pi, \sigma)^{1/4-\delta}.$$

In the everywhere-tempered case, $\delta = 1/(16n^5 + O(n^4))$.

Example

Take $(G, H) = (\mathrm{GL}_3(\mathbb{R}), \mathrm{GL}_2(\mathbb{R}))$, coadjoint orbits = nilcones:

$$\mathcal{O}_\pi = \mathcal{N} \subset \mathfrak{g}^\wedge, \quad \mathcal{O}_\sigma = \mathcal{N}_H \subseteq \mathfrak{h}^\wedge.$$

Then

$$\begin{aligned} \mathcal{O}_{\pi,\sigma} &= \mathcal{N} \cap \text{preimage}(\mathcal{N}_H) \\ &= \left\{ \xi = \begin{pmatrix} A & b \\ c & d \end{pmatrix} \in \mathfrak{sl}_3(\mathbb{R}) : \xi, A \text{ are nilpotent} \right\}. \end{aligned}$$

Very far from stable: $\#\{H\text{-orbits on } \mathcal{O}_{\pi,\sigma}\} = \infty!$ Representatives

$$\begin{pmatrix} 0 & 1 & b \\ 0 & 0 & 0 \\ 0 & c & 0 \end{pmatrix}$$

with bc the only invariant. Behavior of $\mathcal{H}_{\pi,\sigma}(a)$ remains unclear.