

## ALGEBRAIC NUMBER THEORY W4043

HOMEWORK, WEEK 10, DUE APRIL 14

The following exercises study the basic properties of the  $p$ -adic norm. Let  $x = \frac{a}{b} \in \mathbb{Q}$ , with  $a, b \in \mathbb{Z}$ . If  $x \neq 0$ , write  $x = p^r \cdot \frac{a'}{b'}$  with  $p$  relatively prime to both  $a'$  and  $b'$  and  $r \in \mathbb{Z}$  (not necessarily positive), and define

$$|x|_p = p^{-r}.$$

Thus  $|p|_p = p^{-1}$ . We also define  $|x|_p = 0$ .

1. Show that  $|\bullet|_p$  has the properties of a metric:
  1. For all  $x \in \mathbb{Q}$ ,  $|x|_p \geq 0$ , with  $|x|_p = 0$  if and only if  $x = 0$ .
  2. For all  $x, y \in \mathbb{Q}$ ,  $|xy|_p = |x|_p|y|_p$ .
  3. For all  $x, y \in \mathbb{Q}$ ,  $|x + y|_p \leq \sup(|x|_p, |y|_p)$ .

Item 3. is the *non-archimedean* (or ultrametric) property; it is stronger than the usual triangle inequality. It allows us to define  $\mathbb{Q}_p$  as the set of equivalence classes of infinite series

$$\sum_{i=0}^{\infty} a_i, a_i \in \mathbb{Q}; \lim_{i \rightarrow \infty} |a_i|_p = 0$$

where  $\sum_{i=0}^{\infty} a_i$  and  $\sum_{i=0}^{\infty} b_i$  are defined to be equivalent if

$$\lim_{N \rightarrow \infty} \left| \sum_{i=0}^N a_i - \sum_{i=0}^N b_i \right|_p = 0.$$

The  $p$ -adic norm extends to  $\mathbb{Q}_p$  by setting

$$\left| \sum_{i=0}^{\infty} a_i \right|_p = \lim_{N \rightarrow \infty} \left| \sum_{i=0}^N a_i \right|_p.$$

2. (a) Show that  $|\sum_{i=0}^{\infty} a_i|_p$  is always either 0 or a power (positive or negative) of  $p$ .

(b) Show that  $|\sum_{i=0}^{\infty} a_i|_p = |\sum_{i=0}^{\infty} b_i|_p$  if  $\sum_{i=0}^{\infty} a_i$  and  $\sum_{i=0}^{\infty} b_i$  are equivalent.

3. (a)  $a \in \mathbb{Q}$ ,  $a \neq 0$ . Show that  $|a|_p = 1$  for all but finitely many prime numbers  $p$ .

(b) (*The product formula*) Let  $a \in \mathbb{Q}$ ,  $a \neq 0$ . Let  $|a|$  be the usual absolute value (equal to  $a$  if  $a > 0$  and to  $-a$  if  $a < 0$ ). Show that

$$|a| \cdot \prod_p |a|_p = 1,$$

where the product is taken over all prime numbers. (By (a), this is actually a finite product.)

4. Define the *adèle group*  $\mathbf{A}$  to be the subgroup of the direct product  $\mathbb{R} \otimes \prod_p \mathbb{Q}_p$ , where the product is taken over all prime numbers, of elements  $(a_{\mathbb{R}}, (a_p))$  such that  $|a_p|_p \leq 1$  for all but finitely many  $p$ . (The number of  $p$  such that  $|a_p|_p > 1$  depends on the element but it must always be finite.

Show that there is an injective homomorphism  $i : \mathbb{Q} \rightarrow \mathbf{A}$ . Show that the set of  $x \in \mathbb{Q}$  such that  $|x|_p \leq 1$  for all  $p$  is equal to  $\mathbb{Z}$ .