

MATH 797W LECTURE NOTES
 Taught by Jenia Tevelev; Notes by Patrick Lei
 University of Massachusetts, Amherst
Spring 2019

Abstract

This will be the first course in algebraic geometry - the study of geometric spaces locally defined by polynomial equations. It is a central subject in mathematics with strong connections to differential geometry, number theory, and representation theory. We will pursue an algebraic approach to the subject, when local data is studied via the commutative algebra of quotients of polynomial rings in several variables. The emphasis will be on basic constructions and examples. Topics will include projective varieties, resolution of singularities, divisors and differential forms. Examples will include algebraic curves of low genus and surfaces in projective 3-space. In addition to theoretical approach, we will also learn how to use computer algebra software, specifically the Macaulay 2 package, to help with basic calculations in commutative algebra and algebraic geometry.

Forms of evaluation: biweekly homeworks (25%), take-home midterms (50%) and computer algebra project (25%).

CONTENTS

1	Organizational/Philosophical	2
2	Affine Plane Curves	3
2.1	Lecture 1 (Jan 22)	3
2.1.1	Enumerative Geometry	3
2.1.2	Basic Notions	4
2.1.3	Abelian Integrals	5
2.2	Lecture 2 (Jan 24)	6
2.2.1	Non-Mathematics	6
2.2.2	Rational Curves Continued	6
2.2.3	Elliptic Curves	7
2.2.4	Non-singularity of plane curves	8
3	Affine Zariski-Closed Sets	8
3.1	Lecture 3 (Jan 29)	8
3.2	Lecture 4 (Jan 31)	10
3.2.1	Zariski Topology on \mathbb{A}^n	10
3.3	Lecture 5 (Feb 5)	11
3.3.1	Macaulay 2 Interlude	12
3.3.2	Morphisms of Affine Closed Sets	12
3.4	Lecture 6 (Feb 7)	13
3.4.1	Morphisms Continued	13
3.5	Lecture 7 (Feb 14 ♡)	15

3.5.1	Rational Maps	15
3.5.2	Dimension	16
4	Projective Space	17
4.1	Lecture 8 (Feb 21, given by Luca Schaffler)	17
4.1.1	Affine Constructions	18
4.1.2	Irreducibility	19
4.1.3	Projective and Quasiprojective Varieties	19
4.2	Lecture 9 (Feb 26)	19
4.3	Lecture 10 (Feb 28)	21
4.3.1	Products	22
4.4	Lecture 11 (Mar 5)	23
4.5	Lecture 12 (Mar 7)	24
5	Local Properties	26
5.1	Lecture 13 (Mar 19)	26
5.1.1	Tangent space	26
5.2	Lecture 14 (Mar 21)	28
5.2.1	Tangent Cone	29
5.3	Lecture 15 (Mar 26)	30
5.4	Lecture 16 (Mar 28)	31
5.4.1	Blowups	32
5.5	Lecture 17 (Apr 02)	32
5.5.1	Local Blowup II	33
5.6	Lecture 18 (Apr 04)	33
5.6.1	Normal Varieties	35
5.7	Lecture 19 (Apr 09)	35
5.7.1	Differential of a regular map	35
5.7.2	Normal Varieties Continued	36
5.8	Lecture 20 (Apr 11)	37
5.8.1	Normalization	38
5.9	Lecture 21 (Apr 16)	39
6	Divisors	39
6.1	Lecture 22 (Apr 18)	41
6.1.1	Divisors of rational maps to \mathbb{P}^n	41
6.2	Lecture 23 (Apr 23)	42
6.3	Lecture 24 (Apr 25)	43
7	Regular Differential Forms	44
7.1	Lecture 25 (Apr 30)	45

1 ORGANIZATIONAL/PHILOSOPHICAL

Warning: All jokes and conversations are reproduced as best as I can remember and my transcription is not necessarily faithful. In addition, footnotes and some definitions are based on my

understanding of algebraic geometry and related topics. Also, I am a terrible person and do not include diagrams. Form your own geometric intuition.

The syllabus is on Moodle. He wanted to project the syllabus on the board, but was unable to get the projector to work. Jenia promised to email everyone the syllabus today.

Jenia considers Shafarevich (hereto referred to as **Shaf**) to be the best way to learn algebraic geometry.¹ It combines the Russian algebraic geometry approach (by Shafarevich, others, and Jenia himself) with the translation by Miles Reid (British, more colloquial approach). He has used the book to teach both undergrads and graduate students. Knowing projective algebraic varieties is very helpful in the future whether you use homological/categorical, complex (holomorphic), or scheme theory methods. The plan is to cover all of Shafarevich in this semester.

There are people who attempt to read Mumford instead (it's fantastic), but the book is much more difficult and he attempts to develop both varieties and schemes at the same time.

Jenia notes that people who do algebraic geometry use computers, and he is going to be teaching us some Macaulay 2 (especially because computer packages can do commutative algebra). Note that this is not a course in computational algebraic geometry (which Jenia, Paul, and Eyal don't know very much about). There is David Cox at Amherst, who is retiring in June, who wrote a book titled *Ideals, Varieties, and Algorithms*. There will be a computer algebra project.

There will be two kinds of homework problems. Some are normal homework assignments while others will be posted on Moodle. Two homeworks will be designated as take-home midterms.

Jenia struggled to turn off the projector but managed to do so.

2 AFFINE PLANE CURVES

2.1 Lecture 1 (Jan 22) Historically, algebraic geometry came from two directions: projective geometry and abelian integrals. These are the two big sources of algebraic geometry, and much early progress was about the two subjects.

2.1.1 Enumerative Geometry

Theorem 1 (Butterfly Theorem). *Consider an ellipse with a chord AC and let B be the midpoint of AC. Draw two chords through B and form a "butterfly." We have two new points P, Q. Then $PB = BQ$.*

Proof. Observe we have three plane curves passing through four points: A conic e given by f and two unions of two lines c_1, c_2 given by f_1, f_2 . We have a linear system of conics passing through four points.

The space of all conics forms an \mathbb{R}^6 . Every point imposes a linear condition on the coefficients, so conics passing through four points, so the space we care about is at least two-dimensional. We will later prove that this is exactly two.

¹He said something about starting with schemes (like Vakil's notes) being a bad idea.

We see that $f_2 = \alpha f + \beta f_1$. Now we restrict our polynomials to the line passing through A, C and assume it is the x -axis and that B is the origin. We see that $f|_{y=0} = (x-c)(x+c)$, $f_1|_{y=0} = x^2$, and $f_2|_{y=0} = (x-p)(x-q)$. Then we must have that $p+q=0$. \square

Jenia learned this theorem in high school.² He told a story about Sheldon Katz and one of his interviews.

Lemma 2. *The dimension of the vector space is exactly 2.*

Proof. Assume $\dim(V) \geq 3$. Let X be another point of the ellipse and consider the subspace $V_0 \subset V$ of conics passing through X . Then we see that $\dim(V_0) \geq 2$. Then we have two linearly independent conics passing C, D passing through five points. This contradicts Bezout's Theorem which states that if C has degree n and D has degree m , then $|C \cap D| \leq nm$ unless the curves have a shared component. \square

Jenia says this is nice argument because it connects to enumerative geometry, the connection between algebraic and geometric objects, and curves on surfaces, which we will talk about later when we talk about divisors.

Another famous example of a similar argument is Pascal's Theorem:

Theorem 3 (Pascal). *Given a conic with an inscribed hexagon, then opposite sides intersect at three collinear points.*

Jenia attempted to draw the diagram on the board, but diagrams are not to scale. The argument is very similar to the butterfly proof and can be found in **Shaf**.

2.1.2 *Basic Notions* We will give some definitions to formalize affine plane curves.

Definition 4 (Affine Plane Curve). Let k be algebraically closed and denote by \mathbb{A}^2 the affine plane. Let $f \in k[x, y]$ be nonconstant of degree $d > 0$. Then the vanishing locus C of f is an affine plane curve of degree d . Examples are lines, conics, cubics, quartics, and quintics (which are only for experts).

Remark 5. Observe that $k[x, y]$ is a UFD (more generally, if R is a UFD, then $R[x]$ is a UFD). This is an application of Gauss's Lemma.

Definition 6 (Irreducible Components). We factor $f = f_1 \cdots f_r$ into irreducibles. Then the vanishing locus c_i of each f_i is an irreducible component of C .

Why do we consider algebraically closed fields? One reason is to have more points. We can see that if our field is algebraically closed, our curve has points.

Lemma 7. *Let $k = \bar{k}$. Then:*

1. C contains infinitely many points.
2. Suppose $f \in k[x, y]$ is irreducible and $g \in k[x, y]$. Then the set $\{f = g = 0\}$ is finite unless f divides g . In particular, an irreducible polynomial is uniquely determined up to a scalar by its curve.

²Unlike Americans, he seems to have gotten an actual Euclidean geometry education.

Proof. We use the fact that k is algebraically closed.

1. Every algebraically closed field is infinite, so we can write $f = \sum g_i(x)y^i$. Then take any x_0 and solve for y . Note each g_i is a polynomial and has finitely many roots, so there are only finitely many x_0 such that $\sum g_i(x_0)y^i$ is a nonzero constant.
2. We use Gauss's Lemma in a different form: Let R be a UFD with field of fractions K . Then let $f \in R[y]$. Then if f is irreducible over R , it must be irreducible over K . We see that $f, g \in k[x, y] = k[x][y] \subset k(x)[y]$. By Gauss's Lemma, we know that f does not divide g in $k(x)[y]$ and that f is irreducible. Note that $k(x)[y]$ is a PID, so f, g are coprime and using Bezout's Lemma, we have $1 = \alpha f + \beta g$. Let p be an LCM of the denominators of the coefficients of α, β . Then we have $p(x) = \alpha_0 f + \beta_0 g$. Suppose $f(x_0, y_0) = g(x_0, y_0) = 0$. Then $p(x_0) = 0$, which has only finitely many solutions for x_0 . By the same argument, there are only finitely many possible values for y_0 .

□

During the proof of the previous lemma, Jenia said, "Every time we run into a problem, we have to use a little bit more commutative algebra. That's how algebraic geometry works." In addition, Jenia's phone rang and made frog noises.

We now have enough vocabulary in plane curves to state Bezout's Theorem, but Jenia now wants to talk about Abelian integrals.

2.1.3 *Abelian Integrals* We want to be able to calculate

$$\int u(x, \sqrt{1-x^2}) dx,$$

where u is a rational function in two complex variables. In Calc 2, we use trig substitutions, but Jenia learned something called Euler's substitutions. We can rewrite the integral over the unit circle, which has a rational parameterization

$$(x, y) = \left(\frac{2t}{1+t^2}, \frac{t^2-1}{t^2+1} \right).$$

We can now express the integral in terms of t and compute the integral using partial fractions.

This method is harder than using trig substitution, but is more general. If your curve is rational, then integrating over it is amenable to this method.

Definition 8 (Rational Curve). An irreducible curve $C = (f = 0)$ is rational if there exist $\varphi(t), \psi(t) \in k(t)$ nonconstant such that $f(\varphi(t), \psi(t)) = 0$.³

Corollary 9. If C is rational, then the integral over the curve $\int u(x, y) dx$, where u is rational, can be computed using partial fractions.

Remark 10. Later in the semester, we will talk about differential forms, which are a way to talk about integration in multiple dimensions.

³We will see that this is equivalent to having function field $k(t)$, or being birational to the line.

2.2 Lecture 2 (Jan 24)

2.2.1 Non-Mathematics Jenia was actually able to make the projector work today. He told us to install Macaulay 2 which you can run with the command M2. Jenia showed us some basic commands in Macaulay 2. Next, he showed us the Moodle and the forum with problems for us to solve.

2.2.2 Rational Curves Continued We continue with our preliminary discussion of plane algebraic curves to discuss some history and motivation. Last time we discussed Abelian integrals focusing on the example of the circle. From the rational parameterization $\left(\frac{2t}{1+t^2}, \frac{t^2-1}{t^2+1}\right)$, we can find all possible Pythagorean triples by choosing rational values for t . We saw last time that we can reduce an integral of the form $\int u(x,y) dx + \int v(x,y) dy$ to an integral of the form $\int \varphi(t) dt$.

We introduce a new notion, the field of rational functions $k(C)$ which is the field of all rational functions on the curve. Observe, however, that rational functions are not everywhere defined. There are finitely many points on the curve where the denominator vanishes. We can say that $u_1 = u_2$ if they agree outside their bad points. Algebraically, we can say that $\frac{p_1}{q_1} \sim \frac{p_2}{q_2} \Leftrightarrow f \mid p_1q_2 - p_2q_1$.

Alternately, we can define the coordinate algebra $k[C] = k[x,y]/(f)$ and then define $k(C)$ to be its field of fractions.⁴

Definition 11 (Regular Function). $u \in k(C)$ is regular at $P \in C$ if there exist $p, q \in k[x,y]$ such that $u = \frac{p}{q}$ such that $q(P) \neq 0$.

Example 12. Let $C = \{x^2 + y^2 = 1\}$ and $u = \frac{1-x}{y}$. Then we can write $u = \frac{y}{1+x}$, so it is regular at the point $(1,0)$.

We can see that $k(C)$ is finitely generated by x, y . We also see that $k(C)$ has transcendence degree⁵ 1 because $f(x,y) = 0$ is an algebraic dependence.

Lemma 13. Every finitely generated field of transcendence degree 1 is isomorphic to $k(C)$ for some irreducible curve C .

Remark 14. Different curves can have the same function field.

Definition 15. We say that curves C, D are birational if $k(C) \simeq k(D)$.⁶

Example 16. Let C be the circle and observe that $k(C) \simeq k(t) \simeq k(\mathbb{A}^1) = k(x)$, so C is birational to \mathbb{A}^1 .

Suppose C is a rational curve $\{f=0\}$. Then there exist $\varphi, \psi \in k(t)$ nonconstant such that $f(\varphi(t), \psi(t)) = 0$.

Lemma 17. If C is a rational curve, then $k(C) \simeq k(t)$.

Proof. Take the obvious map. Then we see that this is defined because if $q(\varphi(t), \psi(t)) = 0$, then $f \mid q$. \square

⁴We will formally define the function field and coordinate for affine varieties later.

⁵This equals its geometric dimension.

⁶This is equivalent to the geometric dimension.

So we see that rational curves have function field a subfield of $k(t)$. However, we invoke a theorem from algebra.

Theorem 18 (Lüroth). *Every subfield of $k(t)$ that is not a subfield of k is isomorphic to $k(t)$.*

Corollary 19. *A curve C is rational if and only if C is birational to a line.*

Now we discuss nonrational curves and try to find a simple example.

2.2.3 Elliptic Curves In 1655, Wallis was interested in computing the arc length of an ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$. Then we see that $y = \frac{b}{a}\sqrt{a^2 - x^2}$ and $y' = \frac{-bx}{a\sqrt{a^2 - x^2}}$. We substitute into the formula for arc length $\int \sqrt{1 + (y')^2} dx$, change variables, and obtain an integral of the form

$$\int \frac{a - ae^2x^2}{\sqrt{(1 - e^2x^2)(1 - x^2)}} dx.$$

Then set $C = \{y^2 = (1 - e^2x^2)(1 - x^2)\}$. We will see that this curve is an elliptic curve.

Let's perform an invertible change of variables by moving 1 to ∞ . Then we set $t = \frac{1}{1-x}$. We now obtain the equation

$$y^2 = \left(1 - e^2 \frac{(t-1)^2}{t^2}\right) \frac{1}{t} \left(1 + \frac{t-1}{t}\right).$$

Now multiply by t^4 and obtain the equation

$$(yt^2)^2 = t^2 - e^2(t-1)^2(2t-1)$$

before setting $s = yt^2$ to obtain an equation of the form $s^2 = f_3(t)$, where $f_3(t)$ is a cubic polynomial. Then, after further simplification,⁷ we obtain an elliptic curve D given by $s^2 = w^3 + aw + b$, which is birational to our original curve C .

Theorem 20. *Elliptic curves are not rational.⁸ Proof of this fact will give us motivation to understand other definitions*

Sketch. First, we projectivize our curves, so we have an elliptic curve $C \hookrightarrow \mathbb{P}^2$ and a line $L \hookrightarrow \mathbb{P}^2$. The next step is to show that the two curves are non-singular. Third, we show that if C, L are birational and non-singular, then they are isomorphic. Then, we see that C has a simple involution φ given by $s \mapsto -s$. Then we see that s has four fixed points. Now we use this involution to create an involution $f \circ \varphi \circ f^{-1}$ with $4 > 2$ fixed points. However, every automorphism⁹ of L is given by the images of three points,¹⁰ so this is impossible. \square

⁷This is only possible if the field is not of characteristic 2,3.

⁸Jenia says Dummit and Foote give an algebraic proof of this fact.

⁹The automorphism group of \mathbb{P}^1 is $\text{PGL}(2, k)$

¹⁰This generalizes to higher dimensional projective spaces.

2.2.4 Non-singularity of plane curves Consider a curve $C = \{f = 0\}$ and let $(a, b) \in C$. Then we have a Taylor expansion $f(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b) + \dots$.

Definition 21 (Non-singularity). A plane curve C is nonsingular at (a, b) if at least one of $f_x(a, b), f_y(a, b) \neq 0$. In this case, $f(x, y) = f(a, b) + f_x(a, b)(x - a) + f_y(a, b)(y - b)$ is called the tangent line.

Definition 22 (Multiplicity). We say that $(a, b) \in C$ has multiplicity m is the smallest positive integer such that an order m partial derivative does not vanish at m .

We see that a point is singular if it has multiplicity at least 2.

Example 23 (Singular Conics). Take $(0, 0) \in C$ and suppose the origin is a singularity. Then $f = \alpha x^2 + \beta y^2 + \gamma xy = (\alpha x + \beta y)(\gamma x + \delta y)$. Thus C must be a union of two lines.

3 AFFINE ZARISKI-CLOSED SETS

3.1 Lecture 3 (Jan 29) We will put plane curves away for a while and then return to them. In this section, we will review several facts from commutative algebra.

Theorem 24 (Noether Normalization Lemma). Let A be a finitely generated k -algebra.¹¹ Then A contains x_1, \dots, x_n algebraically independent over k and such that A is integral over $k[x_1, \dots, x_n]$.

Proof. We will assume that k is infinite. Choose some generators y_1, \dots, y_r . We argue by induction on r . If $r = 0$, then $A = k$, so there is nothing to prove. If y_1, \dots, y_r are algebraically independent, then $A = k[y_1, \dots, y_r]$.

Suppose that $f(y_1, \dots, y_r) = 0$ for some $f \in k[Y_1, \dots, Y_r]$. We can write

$$f = \sum_{i=0}^d h_i(y_1, \dots, y_{r-1})y_r^i.$$

If $h_d = 1$, then y_r is integral over the subring B generated by y_1, \dots, y_{r-1} . By induction, B contains algebraically independent x_1, \dots, x_n such that B is integral over $k[x_1, \dots, x_n]$. Therefore, A must be integral over $k[x_1, \dots, x_n]$ by transitivity of integrality.

Noether's trick is to show that we can always reduce to the simple case by a linear change of variables. Introduce $y'_1 = y_1 - \lambda_1 y_r, \dots, y'_{r-1} = y_{r-1} - \lambda_{r-1} y_r, y'_r = y_r$. We see that y'_1, \dots, y'_r generate A . Then we see that $f(y_1, \dots, y_r) = f(y'_1 + \lambda_1 y'_r, \dots, y'_{r-1} + \lambda_{r-1} y'_r, y'_r) = 0$. Write $f = F + \text{l.o.t.}$ and suppose F is of degree d . Then we see that $F(y'_1 + \lambda_1 y'_r, \dots, y'_{r-1} + \lambda_{r-1} y'_r, y'_r) = F(\lambda_1, \dots, \lambda_{r-1}, 1)y_r'^d$. We simply need $F(\lambda_1, \dots, \lambda_{r-1}, 1) \neq 0$, which is always possible if k is infinite. \square

Recall the fundamental theorem of algebra, which states that \mathbb{C} is algebraically closed. This gives a bijection between \mathbb{C} and maximal ideals $\mathfrak{m} \subset \mathbb{C}[x]$ by $a \leftrightarrow (x - a)$. Indeed, over any field, maximal ideals of $k[x]$ are principal ideals generated by irreducible monic polynomials. If $k = \bar{k}$, then all monic polynomials are linear.

¹¹Note here that k does not have to be closed.

Theorem 25 (Weak Nullstellensatz). Let $k = \bar{k}$. Then there is a bijection between $\mathbb{A}^n = k^n$ and maximal ideals in $k[x_1, \dots, x_n]$ given by $(\mathfrak{a}_1, \dots, \mathfrak{a}_n) \leftrightarrow (x_1 - a_1, \dots, x_n - a_n)$.

Proof. Given a point $(a_1, \dots, a_n) \in \mathbb{A}^n$, consider a homomorphism $k[x_1, \dots, x_n] \rightarrow k$ given by $f \mapsto f(a_1, \dots, a_n)$. Then the kernel of this morphism is a maximal ideal, but it must be equal to (a_1, \dots, a_n)

Now take $\mathfrak{m} \subset k[x_1, \dots, x_n]$ maximal and suppose $k[x_1, \dots, x_n]/\mathfrak{m} \xrightarrow{\varphi} K \supset k$ is a field. If $K = k$, define $a_i = \varphi(x_i)$. Then because φ acts on the variables just like the evaluation morphism, we must have $\mathfrak{m} = \ker \varphi = (x_1 - a_1, \dots, x_n - a_n)$.

We need to show that $K = k$. Note that K must be a finitely generated k -algebra. By Noether, K is integral over a polynomial subring $k[x_1, \dots, x_n]$. Recall the fact that if A is a field and A is integral over B , then B is a field. To show this, we see that $b^{-1} \in A$, so $b^{-1} = (b^{-1})^r + a_1(b^{-1})^{r-1} + \dots + a_r = 0$ where $a_1, \dots, a_r \in B$. Multiplying through by b^{r-1} , we see that $b^{-1} \in B$. Therefore $k[x_1, \dots, x_n]$ is a field and $\mathfrak{n} = 0$ and K is integral over k . Because k is algebraically closed, $K = k$. □

Definition 26. A subset $X \subset \mathbb{A}^n$ is called a closed affine set¹² if $X = \{ \mathbf{a} \in \mathbb{A}^n \mid f_1(\mathbf{a}) = \dots = f_r(\mathbf{a}) = 0 \}$ for some polynomials $f_1, \dots, f_r \in k[x_1, \dots, x_n]$.

Lemma 27. Closed sets have the form $V(I) = \{ (a_1, \dots, a_n) \in \mathbb{A}^n \mid f(a_1, \dots, a_n) = 0 \text{ for all } f \in I \}$ for some ideal $I \subset k[x_1, \dots, x_n]$.

Proof. Take $X = \{ f_1 = \dots = f_r = 0 \}$. Then let $I = (f_1, \dots, f_r) \subset k[x_1, \dots, x_n]$. Then $V(I)$ must equal X . On the other hand, take $V(I)$. Because $k[x_1, \dots, x_n]$ is Noetherian, let f_1, \dots, f_r to be generators of I . Then $V(f_1, \dots, f_r) = V(I) = X$. □

Now let $X \subset \mathbb{A}^n$ be a closed set. Define $I(X) = \{ f \in k[x_1, \dots, x_n] \mid f(\mathbf{a}) = 0 \text{ for all } \mathbf{a} \in X \}$. We now have two maps $\{ \text{Closed Sets} \} \xleftarrow[V(I(X))]{V(I)} \{ \text{Ideals} \}$.

Theorem 28 (Strong Nullstellensatz). $V(I(X))$ and $I(V(I)) = \sqrt{I}$.

Proof. Take $g \in I(V(I))$. Suppose $I = (f_1, \dots, f_r)$. We need to show that if g vanishes at every point (a_1, \dots, a_n) such that $f_i(a_1, \dots, a_n) = 0$, then $g^\ell = h_1 f_1 + \dots + h_r f_r$ for some ℓ . We use a trick of Rabinowitch: Let $B = (f_1, \dots, f_r, 1 - g x_{n+1}) \subset k[x_1, \dots, x_{n+1}]$. We show that $B = k[x_1, \dots, x_{n+1}]$. If not, it is contained in some maximal ideal $B \subset \mathfrak{m} = (x_1 - a_1, \dots, x_{n+1} - a_{n+1})$. However, $f_i(a_1, \dots, a_n) = 0$ and $1 - g(a_1, \dots, a_n) a_{n+1} = 0$. But remember $g \in I(X)$, so we obtain that $1 = 0$.

Thus we can write $1 = \sum h_i f_i + h_{n+1}(1 - g x_{n+1})$. Then we write $x_{n+1} = \frac{1}{g(x_1, \dots, x_n)}$. Then we obtain an expression of the form

$$1 = \sum_{i=1}^r h_i(x_1, \dots, \frac{1}{g(x_1, \dots, x_n)}) f_i.$$

¹²Shafarevich calls these closed sets. Some people call them affine sets and other names. Reid uses algebraic sets.

Then we can clear denominators and obtain that $g \in \sqrt{I}$.

Finally we show that $V(I(X)) = X$. Then there exists an ideal J such that $X = V(J)$. Then $V(I(X)) = V(I(V(J))) = V(\sqrt{J}) = V(J) = X$. \square

Remark 29. $I(X)$ is always radical. In fact $V(\sqrt{I}) = V(I)$.

Corollary 30. *The operations I and V give a bijection between closed sets in \mathbb{A}^n and radical ideals in $k[x_1, \dots, x_n]$.*

3.2 Lecture 4 (Jan 31) Recall that the weak Nullstellensatz gives a bijection between points in \mathbb{A}^n and maximal ideals in $k[x_1, \dots, x_n]$. Also, the strong Nullstellensatz gives a bijection between affine closed sets and radical ideals.

3.2.1 Zariski Topology on \mathbb{A}^n

Proposition 31. *The affine closed sets $V(I)$ form the set of closed sets for the Zariski topology on \mathbb{A}^n .*

Proof. 1. $\mathbb{A}^n = V(0)$ and $\emptyset = V((1))$;

2. $\cap_{i \in I} Y_i = V(\sum I_i)$;

3. $\cup_{i=1}^r Y_i = V(I_1 \cdots I_r) = V(I_1 \cap \cdots \cap I_r)$.

\square

Definition 32 (Irreducible Closed Sets). A closed set is called irreducible if it is not the union of two proper closed subsets.

Example 33. In \mathbb{R}^n with the Euclidean topology only points are irreducible.

Theorem 34. *Under the correspondence between $V(I), I(X)$, we will see that irreducible subsets $Y \subset \mathbb{A}^n$ correspond to prime ideals $\mathfrak{p} \subset k[x_1, \dots, x_n]$, which we denote by $\text{Spec } k[x_1, \dots, x_n]$.*

Proof. Suppose $Y \subset \mathbb{A}^n$ is reducible with $Y = Y_1 \cup Y_2$. Then $I(Y) \subsetneq I(Y_i)$ for $i = 1, 2$ by the Nullstellensatz. Then choose $f_1 \in I(Y_1) \setminus I(Y)$ and $f_2 \in I(Y_2) \setminus I(Y)$. Then $f_1 f_2$ vanishes on Y and therefore $I(Y)$ is not prime.

Suppose $I(Y)$ is not prime. Then there exist $f, g \notin I(Y) = I$ with $fg \in I(Y)$. Then write $Y_1 = V(f) \cap Y$ and $Y_2 = V(g) \cap Y$, so we see that $Y = Y_1 \cup Y_2$. \square

Now we consider this correspondence in the case of the plane.

Irreducible subsets	Prime ideals
Points (a, b)	Maximal ideals $(x - a, y - b)$
Irreducible affine plane curves	(f)
\mathbb{A}^2	0

Figure 1: Irreducible subsets of \mathbb{A}^2 and prime ideals in $k[x, y]$.

Why is there nothing else? Suppose we have a prime ideal $0 \neq \mathfrak{p} \subset k[x, y]$. Then write $\mathfrak{p} = (f_1, \dots, f_r) = f(g_1, \dots, g_r)$ where $f = \gcd(f_1, \dots, f_r)$. Then either $f \in \mathfrak{p}$ or $g_i \in \mathfrak{p}$ for all i . If $f \in \mathfrak{p}$, we have that $\mathfrak{p} = (f)$. Otherwise, $\mathfrak{p} = (g_1, \dots, g_r)$. However, the g_i are coprime so $V(\mathfrak{p})$ is a finite union of points, so it must be a single point. Thus \mathfrak{p} is maximal.

“That’s what most arguments in algebraic geometry look like: some input from algebra and some input from geometry.”

Definition 35 (Irreducible Component). Let $Y \subset \mathbb{A}^n$ be a Zariski closed set. Then an irreducible component of Y is a maximal irreducible subset of Y .

Lemma 36. *There exist only finitely many irreducible components Y_1, \dots, Y_s and $Y = Y_1 \cup \dots \cup Y_s$.*

Proof. We show that Y can be written as a finite union $z_1 \cup \dots \cup z_r$ of irreducible subsets. Given that, we can also assume that $z_i \subsetneq z_j$ for all $i \neq j$. Take some irreducible subset $W \subset Y = Z_1 \cup \dots \cup Z_r$. But in fact $W = \cup(W \cap Z_i)$, so $W \subset Z_i$ for some i . Then in particular, if W is an irreducible component, then $W = Z_i$. Therefore this decomposition is a decomposition into irreducible components.

To prove the claim, if Y is irreducible, then clearly this is true. Then write $Y = Y_1 \cup Y_2$. If Y_1, Y_2 are irreducible, then we are done. Suppose Y_1 is reducible. Then write it as a union and continue to form a binary tree. But this tree must be finite because $k[x_1, \dots, x_n]$ is Noetherian. \square

Remark 37. The algebraic counterpart to that is $I = \mathfrak{p}_1 \cap \dots \cap \mathfrak{p}_s$.

Proof. Take $f \in \mathfrak{p}_1 \cap \dots \cap \mathfrak{p}_s$. Then f vanishes on every y_i , so it vanishes on y . Thus $f \in I$. \square

Theorem 38. *Every radical ideal in $k[x_1, \dots, x_n]$ is the intersection of minimal prime ideals which contain I .¹³*

We now shift to a more general perspective. So far we have considered \mathbb{A}^n and $k[x_1, \dots, x_n]$. Now we will consider $X, k[X]$ where X is an irreducible Zariski-closed subset of \mathbb{A}^n and $k[X] = k[x_1, \dots, x_n]/I(X)$ the coordinate ring of X . We see that X is irreducible if and only if $k[X]$ is an integral domain.

Theorem 39 (Generalized Nullstellensatz). *There is a bijection between points of X and maximal ideals of $k[X]$, between closed subsets of X and radical ideals of $k[X]$, and between irreducible closed subsets of X and prime ideals of $k[X]$.*

Proof. Suppose $Y \subset X$ is a closed subset. Then $I(Y) \supset I(X)$ as radical ideals. Then $I(Y)/I(X) \subset k[X]$ is a radical ideal. \square

3.3 Lecture 5 (Feb 5)

¹³This is true in every Noetherian ring.

3.3.1 Macaulay 2 Interlude Recall that we have algebraic sets $X \hookrightarrow \mathbb{A}^n$ and the coordinate algebra $k[X] = k[x_1, \dots, x_n]/I(X)$. Then for any $Y \subset X$, $I(Y) \subset k[X]$ is a radical ideal and Y irreducible implies $I(Y)$ is prime. Then we can decompose Y as a union of irreducible components and $I(Y)$ as an intersection of minimal primes.

There are algorithms that can do these computations, but they generally use Gröbner bases, which are difficult to compute by hand. Fortunately, we can use computers, such as Macaulay2, to perform these calculations.

```
E = QQ[x] -- polynomial ring *-
ideal(x^2+1) -- ideal of i *-
isPrime oo -- true *-
clearAll
E = QQ[x,I]/ideal(I^2+1) -- i is in the ring *-
A = ideal(x^2+1)
isPrime A -- false *-
L = decompose A -- List of two ideals *-
length L -- 2 *-
L_0 -- x-i *-
L_1 -- x+i *-
clearAll
```

We now consider the Clebsch cubic surface $x^3 + y^3 + z^3 + 1 = (x + y + z + 1)^3$. There is a famous theorem that any smooth cubic surfaces contain exactly 27 lines. We will calculate these 27 lines.

```
R = ZZ/101[x,y,z,x0,y0,z0,a,b,c,t] -- initialize all parameters *-
cubic = x^3+y^3+z^3+1-(x+y+z+1)^3 -- cubic *-
cubt = sub(cubic, {x=>x0+a*t, y=>y0+b*t, z=>z0+c*t}) -- substitute parametric equation of line
--> *-
cubic0 = sub(cubt, t=>0) -- constant term *-
cubic1 = sub(diff(t,cubt), t=>0) -- linear term *-
cubic2 = sub(diff(t,diff(t,cubt)), t=>0) -- quadratic term *-
cubic3 = sub(diff(t,diff(t,diff(t,cubt))), t=>0) -- cubic term *-
Lines = ideal(cubic0,cubic1,cubic2,cubic3) -- set identically zero *-
L=decompose Lines -- find all ideals *-
#L -- should be 25 *-
f = x^3+y^3+z^3-(x+y+z)^3 -- ideal at infinity *-
factor f -- three lines *-
```

We found an interesting phenomenon where the computation took longer and was the computer was unable to find the expected 25 affine lines when we worked over $\mathbb{Z}/103\mathbb{Z}$ because 5 is not a quadratic residue mod 103.¹⁴

3.3.2 Morphisms of Affine Closed Sets Let $f : \mathbb{A}^n \rightarrow \mathbb{A}^m$ be given by $y_i = f_i(x_1, \dots, x_n)$, where $f_i \in k[x_1, \dots, x_n]$ for $i = 1, \dots, m$. Algebraically, this corresponds to a morphism $k[y_1, \dots, y_m] \xrightarrow{\varphi} k[x_1, \dots, x_n]$, which is given by $y_i \mapsto f_i(x_1, \dots, x_n)$.

We know that the kernel of this morphism is a prime ideal because the quotient by it is a subring of $k[x_1, \dots, x_n]$, so is an integral domain. Then $V(\text{Ker } \varphi) = \overline{f(\mathbb{A}^n)}$. To see this, suppose that

¹⁴We went into a digression about quadratic reciprocity.

$(b_1, \dots, b_m) = f(a_1, \dots, a_n)$. Then let $g \in \text{Ker } \varphi$. We see that

$$\begin{aligned} g(b_1, \dots, b_m) &= g(f_1(a_1, \dots, a_n), \dots, f_m(a_1, \dots, a_n)) \\ &= [\varphi(g)](a_1, \dots, a_n) \\ &= 0. \end{aligned}$$

In fact, $\overline{\text{Im } f} = V(\text{Ker } \varphi)$.

Let $X \hookrightarrow \mathbb{A}^n$. Now we define a morphism to be a restriction of a polynomial morphism from $\mathbb{A}^n \rightarrow \mathbb{A}^m$. We pull back $k[y_1, \dots, y_m] \rightarrow k[X]$ in the same way.

Finally, suppose $X \subset \mathbb{A}^n, Y \subset \mathbb{A}^m$. Then let $f : X \rightarrow \mathbb{A}^m$ and suppose $f(X) \subset Y$. Then we have a morphism $X \rightarrow Y$ and a pullback morphism $k[Y] \rightarrow k[X]$.

Theorem 40. *There is a bijection $\text{Hom}(X, Y) \simeq \text{Hom}(k[Y], k[X])$.*¹⁵

3.4 Lecture 6 (Feb 7)

3.4.1 Morphisms Continued Recall the definition of a morphism of affine algebraic sets. Also recall that any morphism $f : X \rightarrow Y$ induces a pullback homomorphism $f^* : k[Y] \rightarrow k[X]$.

Also recall Theorem 40. In particular, $X \simeq Y$ if and only if $k[X] \simeq k[Y]$.

Proof of Theorem 40. Recover f from f^* by $b_i = y_i(b) = y_i(f(a)) = (f^*y_i)(a)$. Then given a morphism $\alpha : k[Y] \rightarrow k[X]$, consider the following diagram.

$$\begin{array}{ccc} k[Y] & \xrightarrow{\alpha} & k[X] \\ \uparrow & & \uparrow \\ k[y_1, \dots, y_m] & & k[x_1, \dots, x_n] \end{array}$$

Then there exist f_i such that the image of f_i under the map $k[x_1, \dots, x_n] \rightarrow k[X]$ is $\alpha(y_i)$. Define $f = (f_1, \dots, f_m)$. Clearly $\alpha = f^*$.

We check that $f(X) \subset Y$. Note $g \in O(Y)$ given by $g(f(a_1, \dots, a_n)) = 0$ if and only if $g(f_1, \dots, f_m) \in I(X)$ if and only if $\alpha(g) \in I(X)$ if and only if g restricted to Y is zero. \square

Recall that $X \hookrightarrow \mathbb{A}^n$ corresponds algebraically to $k[x_1, \dots, x_n] \twoheadrightarrow k[X]$, which is choice of generators of $k[X]$.

Let G be a finite group acting on X . Then G must also act on $k[X]$.

Theorem 41. *If the char does not divide $|G|$, then $k[X]^G$ is finitely generated.*

Choose generators g_1, \dots, g_n of $k[X]^G$ given by $\psi : k[y_1, \dots, y_n] \twoheadrightarrow k[X]^G$. Let $I = \ker \psi$.

Definition 42 (Quotient by a finite group). Define $Y = V(I) \subset \mathbb{A}^n$. Then $k[Y] \simeq k[X]^G$. Then $Y = X/G$. Note that $k[X]^G \hookrightarrow k[X]$, so there is a quotient morphism $\pi : X \rightarrow X/G$.

¹⁵This induces an equivalence of categories between affine algebraic sets and finite k -algebras with no nilpotent elements.

Example 43. Consider the action of $\mathbb{Z}/2\mathbb{Z}$ on \mathbb{A}^2 given by $(x, y) \mapsto (-x, -y)$. This gives an action on $k[x, y]$, so $k[x, y]^{\mathbb{Z}/2\mathbb{Z}} = k[x^2, y^2, xy] = k[u, v, w]/(uv - w^2)$. Then we see $\mathbb{A}^2/(\mathbb{Z}/2\mathbb{Z}) = V(uv - w^2) \hookrightarrow \mathbb{A}^3$ and the quotient morphism is given by $(x, y) \mapsto (x^2, y^2, xy)$.

Now we can translate some geometry into algebra and some algebra into geometry.

Definition 44 (Dominance). A morphism $f : X \rightarrow Y$ is dominant if $f(X)$ is (Zariski-)dense in Y .

Proposition 45. f is dominant if and only if f^* is injective.

Proof. Suppose f is dominant. Let $\varphi \in \ker f^*$. We compute $\varphi(f(a)) = (f^*\varphi)(a) = 0$, so $\varphi|_{f(X)} = 0$. However, $f(X) \subset Y$ is dense, so $\varphi|_Y = 0$.

Now suppose f^* is injective but f is not dominant. Therefore $f(X)$ is not dense in Y , which is possible if and only if there exists $\varphi \in K[Y]$ nonzero such that $\varphi|_{f(X)} = 0$. Therefore $\varphi(f(a)) = 0$ for all $a \in X$ and thus $f^*\varphi = 0$, so f^* is not injective. \square

Definition 46 (Finite Morphism). A dominant morphism $f : X \rightarrow Y$ is called finite if $k[X]$ is integral over $k[Y]$. More generally, a morphism is finite if $k[X]$ is integral over the image of $k[Y]$.

Example 47 (Noether Normalization). An integral extension $k[X] \hookrightarrow k[y_1, \dots, y_m]$ corresponds to a finite morphism $X \rightarrow \mathbb{A}^m$.

Example 48 (Normalization of Cusp). Note the map $\mathbb{A}^1 \rightarrow V(y^2 - x^3) \subset \mathbb{A}^2$ given by $t \mapsto (t^2, t^3)$. Algebraically, $k[x, y]/(y^2 - x^3) \rightarrow k[t]$ given by $x \mapsto t^2, y \mapsto t^3$. Because the morphism is dominant, we have $R \simeq k[t^2, t^3] \hookrightarrow k[t]$ is an integral extension (t is a root of $T^2 - t^2$).

Example 49. Consider $k[X]^G \subset k[X]$. This is always integral, so $X \twoheadrightarrow X/G$ is finite. Consider $\alpha \in k[X]$. It is a root of $\prod_{g \in G} (T - g\alpha)$. Each coefficient is invariant because they are elementary symmetric functions on the $g\alpha$.

Recall the Going-up theorem:

Theorem 50 (Going-up). Let $\mathfrak{p} \subset A$ be a prime ideal and let $A \hookrightarrow B$ be an integral extension. Then there exists a prime ideal $\mathfrak{q} \subset B$ such that $\mathfrak{q} \cap A = \mathfrak{p}$.

Theorem 51. Any dominant finite morphism is surjective and has finite fibers.

Proof. Let α be a finite dominant morphism. Take $y \in Y$, which corresponds to a maximal ideal $\mathfrak{m} \subset k[Y]$. We find a point $x \in X$ (a maximal ideal $\mathfrak{n} \subset k[X]$) such that $\alpha(x) = y$ ($\mathfrak{n} \cap k[Y] = \mathfrak{m}$). By going-up, there exists a prime ideal $\mathfrak{q} \subset k[X]$ such that $\mathfrak{q} \cap k[Y] = \mathfrak{m}$. Take any maximal ideal $\mathfrak{n} \supset \mathfrak{q}$. Then $\mathfrak{n} \cap k[Y] \supset \mathfrak{m}$, so by maximality of \mathfrak{m} , $\mathfrak{n} \cap k[Y] = \mathfrak{m}$.

Now we prove that the fibers are finite. Let $y \in Y$ correspond to a maximal ideal $\mathfrak{m} \subset k[Y]$. Then we show that there are finitely many maximal ideals in $k[X]/(k[x] \cdot \mathfrak{m})$. Then $k[X]$ is integral over $k[Y]$, so $A = k[X]/(k[x] \cdot \mathfrak{m})$ is integral over $k[Y]/\mathfrak{m} = k$. Therefore A is a finitely-generated k -module (a finite-dimensional k -vector space).¹⁶ \square

¹⁶We run into an algebra qual problem (Artinian rings have finitely many maximal ideals).

3.5 Lecture 7 (Feb 14 ♡) We begin with a remark: Let $\alpha : X \rightarrow Y$ be a morphism of affine algebraic sets $X \rightarrow Y$. Then there is a pullback $\alpha^* : k[Y] \rightarrow k[X]$. We note that if $Z \subset Y$ is closed, then $\alpha^{-1}(Z) \subset X$ is also closed. Then if $I(Z) = (f_1, \dots, f_r)$, the ideal of $\alpha^{-1}(Z)$ is given by $(\alpha^* f_1, \dots, \alpha^* f_r)$.

This becomes trickier when α is dominant (which is equivalent to α^* being injective). Then we recall the computation of the fiber over a point to prove that dominant morphisms are surjective with finite fibers.¹⁷

We prove that Artinian rings have finitely many maximal ideals:

Lemma 52. *Artinian rings have finitely many maximal ideals.*

Proof. Consider all finite intersections of maximal ideals. This set contains a minimal element $m_1 \cap \dots \cap m_r$. Therefore $m_1 \cap \dots \cap m_r \subset m$ for all maximal ideals m . Then we see $m = m_i$ for some i . (Otherwise, there exists $x_i \in m_i \setminus m$, and then $x_1 \cdots x_r \in m$, which contradicts maximality.) \square

We recall our discussion of group actions and quotients. Suppose we have a finite group G acting on X . Then we have a finitely generated algebra $k[X]^G \hookrightarrow k[X]$. Then we define X/G to be an affine algebraic set such that $k[X/G] \simeq k[X]^G$. Then there exists a quotient morphism $\pi : X \rightarrow X/G$.

Proposition 53. *The fibers of π are G -orbits.*

Proof. First we show that $y = gx$ implies $\pi(x) = \pi(y)$. Suppose not. Then there exists $f \in k[X/G]$ such that $f(\pi(x)) = 0$ and $f(\pi(y)) = 1$. We have a pullback $\pi^* : k[X/G] \hookrightarrow k[X]$. Then there exists a G -invariant function f such that $f(x) = 0$ and $f(y) = 1$, which is impossible.

Next we show that the fiber is precisely the orbit. Suppose that G -orbits of x, y are disjoint. Then there exists $f \in k[X]$ such that $f|_{G \cdot x} = 0$ and $f|_{G \cdot y} = 1$. Then we define $f^\#(x) = \frac{1}{\#G} \sum_{g \in G} f(g \cdot x)$. This new polynomial is now G -invariant and is always 0 on the orbit of x and 1 on the orbit of y . Then we see that $f^\#(\pi(x)) = (\pi^* f^\#)(x) = f^\#(x) = 0$ and $f^\#(\pi(y)) = (\pi^* f^\#)(y) = f^\#(y) = 1$. Thus $\pi(x) \neq \pi(y)$. \square

3.5.1 Rational Maps Now let X be an affine variety. Then $k[X]$ is an integral domain.

Definition 54 (Function Field). The function field $k(X)$ of an affine variety X is the field of fractions of $k[X]$.

Definition 55 (Regular function). Let $f \in k(X)$. Then f is regular at $x \in X$ if f can be written as a fraction $f = \frac{p}{q}$ where $p, q \in k[X]$ and $q(x) \neq 0$.

Definition 56 (Domain of Definition). The domain of definition of f is the set of all $x \in X$ such that f is regular at x .

Proposition 57. *The domain of definition is open.*

Proof. $D = \bigcup_{f=p/q} (X \setminus V(q))$. \square

¹⁷Jenia discussed this again in class, but I will point you back to Theorem 51.

Proposition 58. $f(x)$ is regular at every $x \in X$ if and only if $f \in k[X]$.

Proof. For all $x \in X$, there exists an expression $f = p_x/q_x$ where $q_x(x) \neq 0$. Then take $I = (q_x)$. The vanishing set $V(I) = \emptyset$, so $\sqrt{I} = (1)$, which implies that $1 \in I$. We can write $1 = a_1 q_{x_1} + a_2 q_{x_2} + \cdots + a_r q_{x_r}$. Then

$$f = f \cdot 1 = \sum_{i=1}^r a_i (q_{x_i} f) = \sum_{i=1}^r a_i p_{x_i} \in k[X].$$

□

Definition 59 (Local ring of a point.). Let $x \in X$. Then the ring \mathcal{O}_x is the set of all functions regular at x . Note that this is the same as $k[X]_{m_x}$. This is a local ring.

Definition 60 (Rational Maps). Let $Y \hookrightarrow \mathbb{A}^n$. Then a rational map is a map $f = (f_1, \dots, f_n)$, where $f_i \in k(X)$ such that $f(x) \in Y$ whenever f_1, \dots, f_n are regular at x . Given this, we have a pullback homomorphism $f^* : k[Y] \rightarrow k(X)$.

Proposition 61. We will see that f^* is injective if and only if $f(X)$ is dense in Y . If so, we induce a field extension $k(Y) \hookrightarrow k(X)$. Additionally, the other way around, given $\alpha : k(Y) \hookrightarrow k(X)$, we can construct a dominant rational map $f : X \dashrightarrow Y$ such that $\alpha = f^*$.

Proof. Take $f = (f_1, \dots, f_n) = (\alpha(y_1), \dots, \alpha(y_n))$. □

Definition 62 (Birational Equivalence). X and Y are birationally equivalent if $k(X) \simeq k(Y)$, or equivalently, there exist dominant rational maps $f, g : X \dashrightarrow Y$ such that $f \circ g, g \circ f$ are the identity wherever they are defined. In particular, X is rational if it is birational to an affine space.

We will skip projective and quasiprojective varieties and return to them next week.

3.5.2 *Dimension* Let $X \subset \mathbb{A}^n$. What is the dimension of X ? Here are some ideas:

- “Maximal number of independent parameters.” To make this rigorous, we define the dimension of X to be the transcendence degree of $k(X)$;
- “Maximum possible dimension of a subspace +1.” Convert this into algebra and we define the dimension of X to be the Krull dimension of $k[X]$.

Theorem 63. If X is an affine variety, then the two definitions of dimension agree.

Proof. By Noether’s Normalization Lemma, $k[X]$ is integral over its polynomial subalgebra $k[y_1, \dots, y_n] = k[\mathbb{A}^n]$. Then $\dim X = \text{trdeg } k(X) = \text{trdeg } k(y_1, \dots, y_n) = n = \dim \mathbb{A}^n$.

First we show that the Krull dimension of $k[y_1, \dots, y_n] = n$, and then we show that Krull dimension is preserved by integral extensions.

To see the second claim, let $p_1 \subsetneq \cdots \subsetneq p_r \subsetneq k[X]$ be a chain of prime ideals. Then $p_1 \cap R \subset \cdots \subset p_r \cap R \subset R$ is a chain of prime ideals. As an exercise, show that $p_i \cap R \neq p_{i+1} \cap R$. Now let $q_1 \subset \cdots \subset q_r \subset R$ be a chain of prime ideals in R . Then we use a stronger going-up theorem to lift

the whole chain to $k[X]$. If the q_i are different, then the lifted ideals are different. Thus the Krull dimensions are equal. \square

4 PROJECTIVE SPACE

4.1 Lecture 8 (Feb 21, given by Luca Schaffler) We will work over an algebraically closed field.

Definition 64 (Projective Space). Let $n \in \mathbb{Z}$ be positive. Then the projective space \mathbb{P}^n is defined as $\mathbb{P}^n = (k^{n+1} \setminus \{0\})/G_n$, which is the set of lines in k^{n+1} through the origin.¹⁸

In \mathbb{P}^n , we denote the equivalence class of (z_0, \dots, z_n) by $(z_0 : \dots : z_n)$ in homogeneous coordinates. Observe that at least one of z_0, \dots, z_n is nonzero.

Definition 65. We say that a polynomial $f \in k[z_0, \dots, z_n]$ vanishes at a point $\xi \in \mathbb{P}^n$ if $f(x_0, \dots, x_n) = 0$ for every choice of homogeneous coordinates $\xi = (x_0 : \dots : x_n)$. In this case, we simply write $f(\xi) = 0$.

Definition 66 (Homogeneous Polynomial). A polynomial $f \in k[z_0, \dots, z_n]$ is homogeneous if every monomial term in f has the same degree.

Example 67. $z_0^3 + z_0z_1^2 + z_1^2z_2$ is homogeneous with degree 3.

Remark 68. Every polynomial $f \in k[z_0, \dots, z_n]$ can be decomposed as the sum of its homogeneous components, i.e. $k[x_1, \dots, x_n]$ is a graded algebra.

Proposition 69. If $f \in k[z_0, \dots, z_n]$ vanishes at $p \in \mathbb{P}^n$, then all of its homogeneous components vanish at p .

Proof. Let $f = \sum_{d \geq 0} f_d$ and $p = (x_0 : \dots : x_n)$. Then for all $\lambda \in k^*$ we have

$$\begin{aligned} f(\lambda x_0, \dots, \lambda x_n) &= \sum_{d \geq 0} f_d(\lambda x_0, \dots, \lambda x_n) \\ &= \sum_{d \geq 0} \lambda^d f_d(x_0, \dots, x_n) = 0. \end{aligned}$$

Because k is infinite, then $f_d(x_0, \dots, x_n) = 0$ for all $d \geq 0$. Therefore all f_d vanish at p . \square

Remark 70. With the same assumptions as the previous proposition, the above proof also implies that $f_0 = 0$.

Definition 71 (Closed Set). Let $X \subset \mathbb{P}^n$ is called closed if X is the vanishing set of some set of polynomials $f_1, \dots, f_r \in k[x_0, \dots, x_n]$.

By the previous proposition, it is not restrictive to assume that f_1, \dots, f_r are homogeneous.

Definition 72 (Homogeneous Ideal). An ideal $I \subset k[z_0, \dots, z_n]$ is called homogeneous if it is generated by homogeneous polynomials. Equivalently, I is closed under taking homogeneous parts.

¹⁸The mean way to define this is $\mathbb{P}^n = \text{Gr}(1, n+1)$

Definition 73 (Vanishing Set of Ideal). Let $U \subset k[z_0, \dots, z_n]$ be an ideal. Define $V(I) = \{p \in \mathbb{P}^n \mid f(p) = 0 \text{ for all } f \in U\}$.

Remark 74. If $I \subset k[z_0, \dots, z_n]$ is a homogeneous ideal, then by the Hilbert Basis Theorem, $I = (f_1, \dots, f_r)$, where each f_i can be assumed to be homogeneous.

Definition 75 (Zariski Topology on \mathbb{P}^n). The closed subsets of \mathbb{P}^n are $V(I)$, where $I \subset k[z_0, \dots, z_n]$ is a homogeneous ideal. This forms a topology. If $X \subset \mathbb{P}^n$ is closed, then the Zariski topology on X is the one induced by \mathbb{P}^n .

4.1.1 *Affine Constructions* We will now consider a useful construction.

Definition 76 (Affine Cone). Let $I \subset k[z_0, \dots, z_n]$ be a homogeneous ideal. We can consider the vanishing set $V(I) \subset \mathbb{P}^n$ or the affine cone $V^a(I) \subset \mathbb{A}^{n+1}$.

Example 77. Consider $V(z_0) \subset \mathbb{P}^n$, which is a point. Then $V^a(z_0)$ is a line.

Remark 78. $V(I) = (V^a(I) \setminus \{0\})/G_m$.

Remark 79. $I(V(I)) = I(V^a(I))$.

Remark 80. In the affine cone construction, then something happens: If $I = k[z_0, \dots, z_n]$, then $V(I) = \emptyset$ and $V^a(I) = \emptyset$. Let $I = (z_0, \dots, z_n)$. Then $V(I) = \emptyset$ but $V^a(I) = \{0\}$. The map $V^a(I) \rightarrow V(I)$ fails to be injective exactly in this case.

Proposition 81. Let $I \subset k[z_0, \dots, z_n]$ be a homogeneous ideal. Then the following are equivalent:

1. $V(I) = \emptyset$;
2. $\sqrt{I} \supset (z_0, \dots, z_n)$;
3. There exists $s \in \mathbb{Z}_{>0}$ such that $I \supset I_s$ and $V(I_s) = \emptyset$ but $V^a(I) = \{0\}$.

Proof. First we prove (1) \Leftrightarrow (2). This is because $V(I) = \emptyset$ if and only if $V^a(I) \subset \{0\}$ if and only if $\sqrt{I} = I(V^a(I)) \supset I(0) = (z_0, \dots, z_n)$. To show that (3) \Rightarrow (2), note that $I_s \subset I$, so $(z_0, \dots, z_n) = \sqrt{I_s} \subset \sqrt{I}$.

To see that (2) \Rightarrow (3), then for all $i = 0, \dots, n$, there exists $p_i \in \mathbb{Z}_{>0}$ such that $z_i^{p_i} \in I$. Let $p = \max\{p_i\}$. Then if $s > (n+1)(p-1)$, $I_s \subset I$. \square

Definition 82. In \mathbb{P}^n consider the open subset $\mathbb{A}_i^n = \{(z_0 : \dots : z_n) \in \mathbb{P}^n \mid z_i \neq 0\}$. Note that there is an identification $\mathbb{A}_i^n \rightarrow \mathbb{A}^n$ given by $(z_0 : \dots : z_n) \rightarrow \left(\frac{z_0}{z_i}, \dots, \frac{z_n}{z_i}\right)$. Moreover, for all $X \subset \mathbb{P}^n$ closed, we can define $X_i := X \cap \mathbb{A}_i^n$.

Explicitly, if $X = V(F_1, \dots, F_r)$, then $X_i = V(f_1^{(i)}, \dots, f_r^{(i)})$, where $f_j^{(i)}(x_0, \dots, x_n) = F_j(x_0, \dots, 1, \dots, x_n)$. this is known as de-homogenization.

Conversely, every closed $Y \subset \mathbb{A}^n$ defines canonically $\bar{Y} \subset \mathbb{P}^n$ where we identify \mathbb{A}^n with \mathbb{A}_0^n . The equations for \bar{Y} are given as follows. If $Y = V^a(f_1, \dots, f_r)$, define $F_j(z_0, \dots, z_n) = z_0^{\deg f_j} f_j\left(\frac{z_1}{z_0}, \dots, \frac{z_n}{z_0}\right)$, which is known as homogenization.

Proposition 83. $\bar{Y} = V(F_1, \dots, F_r)$.

Remark 84. X is not necessarily the closure of its affine patches. For example, consider a line at infinity. It is lost during the procedure.

Remark 85. Let F be a homogeneous polynomial. Then $F = z_1^{\deg F} f^{(i)}$.

4.1.2 Irreducibility

Definition 86. A closed set $X \subset \mathbb{P}^n$ is irreducible if X cannot be written as a union $X = X_1 \cup X_2$, where X_1, X_2 are proper closed subsets.

Remark 87. We have a decomposition into irreducible components like in the affine case. Also, X is irreducible if and only if $I(X)$ is prime.

4.1.3 Projective and Quasiprojective Varieties

Definition 88 (Projective Variety). $X \subset \mathbb{P}^n$ closed is called a projective variety.

Definition 89 (Quasiprojective Variety). A quasiprojective variety is an open subset of a projective variety.

Example 90. Affine and projective varieties are both quasiprojective.

4.2 Lecture 9 (Feb 26) Let $X \subset \mathbb{P}^n$ be a quasiprojective variety. We need to define a regular function on X . We define $k(\mathbb{P}^n) = \left\{ \frac{f(z_0, \dots, z_n)}{g(z_0, \dots, z_n)} \mid f, g \in k[z_0, \dots, z_n]_d \text{ for some } d \right\}$, where z_0, \dots, z_n are homogeneous coordinates on \mathbb{P}^n .

Remark 91. In an affine chart, $\mathbb{A}_0^n \subset \mathbb{P}^n$ given by $z_0 \neq 0$, we can write

$$\frac{f(z_0, \dots, z_n)}{g(z_0, \dots, z_n)} = \frac{z_0^d f(1, z_1/z_0, \dots, z_n/z_0)}{g(1, z_1/z_0, \dots, z_n/z_0)} = \frac{f(1, x_1, \dots, x_n)}{g(1, x_1, \dots, x_n)}.$$

We can reverse this process, so $k(\mathbb{P}^n) = k(\mathbb{A}^n)$.

If $f \in k(\mathbb{P}^n)$, $f = \frac{p}{q}$, then f defines a function in some open neighborhood of $x \in X$.

Definition 92 (Regular at a Point). We call this function f regular at $x \in X$.

Definition 93 (Regular Function). A function $f : X \rightarrow k$ is regular on X if for all $x \in X$, f can be written as $f = \frac{p}{q}$ in some open neighborhood of $x \in X$ with $p, q \in k[z_0, \dots, z_d]$ and $q(x) \neq 0$.

Remark 94. We write $k[X]$ for the set of regular functions on X .

Remark 95. If X is affine, then $k[X] = k[x_1, \dots, x_n]/I(X)$.

Remark 96. If X is projective, then $k[X] = k$.

Remark 97. There are examples of quasi-projective varieties X such that $k[X]$ is not finitely generated.¹⁹

Definition 98. A map $X \rightarrow \mathbb{A}^m$ is regular if $f = (f_1, \dots, f_m)$ and $f_i \in k[X]$.

¹⁹Jenia wrote a paper about this. The first examples were due to Rees and Nagata.

Definition 99. A map $f : X \rightarrow Y \hookrightarrow \mathbb{P}^m$ quasiprojective varieties is called regular if for all $x \in X$ we can choose an affine chart $\mathbb{A}_i^m \subset \mathbb{P}^m$ containing $f(x)$ and an open set $U \ni x$ such that $f(U) \subset \mathbb{A}_i^m$ and the induced map $f|_U$ is regular.

Remark 100. This definition does not depend on the choice of affine chart containing $f(x)$.

Definition 101. Quasi-projective varieties X, Y are isomorphic if there exists regular maps $f : X \rightarrow Y$ and $g : Y \rightarrow X$ such that $f = g^{-1}$.

Lemma 102. Let X be a quasi-projective variety. Then every $x \in X$ has an affine neighborhood.

Proof. Let $X \subset \mathbb{P}^n$ and $x \in \mathbb{A}_i^n$. Then $X \cap \mathbb{A}_i^n = \bar{X} \cap \mathbb{A}_i^n - (\bar{X} \setminus X) \cap \mathbb{A}_i^n$, so $x \in Y - Z \subset \mathbb{A}^n$, where Y, Z are both affine. Choose $0 \neq F \in I(Z) \subset k[Y]$ and set $u = Y - V((F)) =: D(F)$. We will show that $D(F)$ is affine, called a principal open set. Principal open sets form a basis of the Zariski topology.

To prove this, note that $Y = (G_1 = \cdots = G_s = 0) \subset \mathbb{A}^n$. Define $Z := (G_1 = \cdots = G_s = Fy_{n+1} = 0) \subset \mathbb{A}^{n+1}$. Then Z is affine and isomorphic to $D(F)$. Simply take the last coordinate to be $1/F$, which is possible because $F \neq 0$ on Z . \square

Definition 103. A regular map $f : X \rightarrow Y$ of quasi-projective varieties is called finite if all $y \in Y$ have an affine neighborhood such that $f^{-1}(V) = U$ is also affine and the induced map $f|_U$ is a finite map of affine varieties.

Proposition 104. If X, Y are both affine, then this agrees with the previous definition.

Proof. Suppose $f : X \rightarrow Y$ is a map of affine varieties. Let $k[X] = B, k[Y] = A$ and the pullback be $f^* : A \rightarrow B$. For all $y \in Y$, there exists an affine $v \in V \subset Y$ such that $U = f^{-1}(V) \subset X$ is affine and $f|_U$ is finite.

Choose a principal open set $v \in D(F) \subset V \subset Y$ for some $f \in A$. Then $f^{-1}(D(F)) \subset U \subset X$. We see that $f^{-1}(D(F)) = D(f^*F)$, so it is affine.

Now we show that $f|_{D(f^*F)}$ is finite. To prove this claim, note that $k[U]$ is a finite $k[V]$ -module via f^* , so we show that $k[U][\frac{1}{f^*F}]$ is a finite $k[V][\frac{1}{F}]$ -module. Simply take the basis to be the basis of $k[U]$ over $k[V]$.

We need to show that B is a finite A -module via f^* . We know that Y is covered by principal open sets $D(F_\alpha)$ such that $k[D(f^*F)]$ is a finite $k[D(F)]$ -module via f^* . We see that $k[D(F)] = A[\frac{1}{F}]$, so $k[D(f^*F)] = B[\frac{1}{f^*F}]$. Also, Y is covered by $D(F_\alpha)$ if and only if $(F_\alpha) = A$. Thus we can write $1 = \sum h_\alpha F_\alpha$. In particular, we need only finitely many $D(F_\alpha)$.

We know that $B[\frac{1}{f^*F_\alpha}]$ is a finite $A[\frac{1}{F_\alpha}]$ -module. Choose a basis of the form $\omega_{\alpha,i} \in B$. We will show that the $\omega_{\alpha,i}$ form a basis of B over A . Choose $b \in B$. Then for all α we can write $b = \sum \omega_{\alpha,i} \frac{a_{i,\alpha}}{F_\alpha^{n_\alpha}}$. We can still write $1 = \sum H_\alpha F_\alpha^{n_\alpha}$.

We write $b = b \sum H_\alpha F_\alpha^{n_\alpha} = \sum H_\alpha (\sum_i \omega_{\alpha,i} a_{i,\alpha})$. \square

4.3 Lecture 10 (Feb 28) Let $f : X \rightarrow Y$ be a morphism of quasiprojective varieties. Then f is locally given by a polynomial map $f = (f_1, \dots, f_m)$ where $f_i = \frac{p_i(x_0, \dots, x_n)}{q_i(x_0, \dots, x_n)}$ and f_i is of degree 0.

Homogeneizing, we obtain that f is given by $f(x) = [F_0 : \dots : F_m]$, where $F_i(x) \in k[x_0, \dots, x_n]_d$ for some d . Also, for every $x \in X$, there exists a presentation $[f_0 : \dots : f_m]$ such that at least one $F_i(x) \neq 0$.

Proposition 105. *The above two notions are equivalent.*

Proof. Suppose f regular at x and $f(x) \in \mathbb{A}_0^m$. Then $f(x) = [1 : p_1/q_1 : \dots : p_m/q_m]$, so we can clear denominators to get $f = [q_1 \cdots q_m : \dots : \dots]$. \square

Example 106. Consider the projection from $\mathbb{P}^\ell \subset \mathbb{P}^n$. On the affine cones, this is just a projection, so the projection is given by $[x_0 : \dots : x_n] \mapsto [x_{\ell+1} : \dots : x_n]$. In this case the F_i are just coordinates. This gives a regular map $\mathbb{P}^n \setminus \mathbb{P}^\ell \rightarrow \mathbb{P}^{n-\ell-1}$.

Example 107. Consider the d -th Veronese embedding $\mathbb{P}^n \xrightarrow{v_d} \mathbb{P}^N$ given by $x \mapsto [F_i(x)]$ where F_i runs through monomials of degree d and $N + 1 = \binom{n+d}{d}$. This is regular everywhere because one of the powers is nonzero. In fact, v_d is an embedding.

An example of this is the rational normal curve.

Theorem 108. *Let $X \subset \mathbb{P}^n$ be an irreducible projective variety. Then $k[X] = k$.*

Proof. We will deduce this from another theorem, which is given below. Using the theorem, take $f \in k[X]$. Then f is a morphism $X \rightarrow \mathbb{A}^1 \hookrightarrow \mathbb{P}^1$. If $f(X)$ is dense in \mathbb{A}^1 , then it is dense in \mathbb{P}^1 . By the next theorem, $f(X) = \mathbb{P}^1$, which contradicts the fact that $f(X) \subset \mathbb{A}^1$.

If $f(X)$ is not dense in \mathbb{A}^1 , then $f(X) = \{p_1, \dots, p_r\}$. However, X is irreducible, so $f(X) = p$ (consider the fiber above each point). \square

Theorem 109. *Let X be a projective variety and $f : X \rightarrow Y$. Then $f(X) \subset Y$ is closed.*

Proof. To prove this, we prove the main theorem of elimination theory. Now we consider the map $f : X \rightarrow Y$. Then consider the graph $\Gamma_f : X \rightarrow X \times Y$ given by $x \mapsto (x, f(x))$. We show that $\Gamma = \Gamma_f(X)$ is closed for all X, Y . Using this, we reduce to the following:

Theorem 110. *Let $X \subset X \times Y$ be closed. Then $\pi_2(Z) \subset Y$ is closed.*

This motivates the following definition:

Definition 111. A variety is called proper if for every variety Y and closed subvariety $Z \subset X \times Y$, $\pi_2(Z) \subset Y$ is closed.

Next we cover Y by affine open sets Y_i . Then $Z = \bigcup Z \cap (X \times Y_i) = Z_i$ and Z_i is closed in $X \times Y$. Then $\pi_2(Z) = \bigcup \pi_2(Z_i)$. It is enough to show that the projections of the Z_i are closed.

Now we have $X \subset \mathbb{P}^n$ projective and $Y \subset \mathbb{A}^m$ affine. Then Z is defined by equations $g_i(u, y)$ homogeneous in x and arbitrary in y . The second projection of Z is precisely the locus T from the main theorem of elimination theory, which is closed. \square

Theorem 112 (Main Theorem of Elimination Theory). *Let $g_i(u, y)$ be polynomials homogeneous in $u = (u_0, \dots, u_n)$ and arbitrary in $y = (y_1, \dots, y_m)$. Let $T = \{y_0 \in \mathbb{A}^m \mid g_i(u, y_0) = 0 \text{ has a nonzero solution}\}$. Then T is closed.*

Proof. Recall $g_i(u, y_0) = 0$ has a nonzero solution if and only if $(g_1(u, y_0), \dots, g_t(u, y_0)) \not\subseteq I_s$ for all s . Then $T = \bigcap_{s \geq 1} T_s$ where $T_s = \{y_0 \in \mathbb{A}^m \mid (g_1(u, y_0), \dots, g_t(u, y_0)) \not\subseteq I_s\}$. It suffices to check that T_s is closed, or that the complement is open.

Indeed, we see that for all monomials $N_{\alpha \in I_s}$, then $M_\alpha = \sum g_i(u, y_0) F_{i, \alpha}(u)$. Let $N_{i, \beta}$ be all monomials of degree $s - \deg g_i$. Thus $M_\alpha \in I_s$ is in the linear span of $g_i(u, y_0) \cdot N_{i, \beta}$. Therefore, $g_i(u, y_0) \cdot N_{i, \beta}$ span the vector space of all degree s polynomials, which means that at least one of the maximal minors is nonzero. Therefore the complement of T_s is a union of open sets and is thus open. \square

4.3.1 Products We discuss products of projective spaces. To do this, we consider the Segre embedding $\mathbb{P}^n \times \mathbb{P}^m \rightarrow \mathbb{P}^N$, where $N + 1 = (n + 1)(m + 1)$, given by $[x_0 : \dots : x_n], [y_0 : \dots : y_m] \mapsto [x_i y_j]_{ij}$. Note that at least one x_i and at least one y_j are nonzero, so this is well-defined.

Alternately, we present this as $[x], [y] \mapsto [xy^T]$, which gives a linear map $\psi : k^{m+1} \rightarrow k^{n+1}$ with kernel y^\perp and image x . Therefore, the Segre embedding is injective. Matrices of this form are precisely the rank 1 matrices, so the image is defined by equations the 2×2 minors of the matrix. In particular, $\mathbb{P}^n \times \mathbb{P}^m$ is a projective variety. This endows $\mathbb{P}^n \times \mathbb{P}^m$ with the structure of a projective variety.

Remark 113. In the charts $\mathbb{A}_0^n, \mathbb{A}_0^m$, we see $s([1 : x_1 : \dots : x_n], [1 : y_1 : \dots : y_m]) = [1 : x_1 : \dots : x_n : y_1 : \dots : y_m : c_i y_j] \subset \mathbb{A}_0^N$. This image is isomorphic to \mathbb{A}^{n+m} . Therefore this agrees on the charts with a product of affine spaces.

If $X \subset \mathbb{P}^n, Y \subset \mathbb{P}^m$ are projective varieties, then $X \times Y \subset \mathbb{P}^n \times \mathbb{P}^m$ is a projective variety.

To define closed subvarieties in $\mathbb{P}^n \times \mathbb{P}^m$ (or $\mathbb{P}^n \times \mathbb{A}^m$) without referring to the ambient projective space \mathbb{P}^N , we do the following:

Theorem 114. *The closed subvarieties of $\mathbb{P}^n \times \mathbb{P}^m$ are given by multihomogeneous (homogeneous in both variables) polynomials $g_i(u, v) = 0$. For $\mathbb{P}^n \times \mathbb{A}^m$, just remove the assumption that g_i are homogeneous in v .*

Proof. Consider the image under the Segre embedding. Then we see that $Z = (F_\alpha(w_{ij}) = 0)$ is the same as $F_\alpha(x_i, y_j) = 0$ homogeneous in X and Y of the same degree. If the g_i are not of the same degree in u, v ($s > t$), we can multiply by v_j^{s-t} to obtain equivalent equations. \square

4.4 Lecture 11 (Mar 5) We continue our discussion of dimension (from Valentine's Day). Let $X \subset \mathbb{P}^n$ be an irreducible quasi-projective variety. Then we define the local ring $\mathcal{O}_X = \{f \in k(\mathbb{P}^n) \mid f \text{ is regular at } x \in X\}$. Then we can define $k(X) = \mathcal{O}_X/\mathfrak{m}_X$ where \mathfrak{m}_X is the maximal ideal of functions that vanish on X .

Then if $U \subset X$ is open, $k(U) = k(X)$. Thus we can assume X is affine. Then we see that $X \subset \mathbb{A}^n \subset \mathbb{P}^n$ is the vanishing $X = V(\mathfrak{p})$ and $\mathcal{O}_X = k[x_1, \dots, x_n]_{\mathfrak{p}}$. Then $\mathfrak{m}_X = \mathfrak{p}R_{\mathfrak{p}}$, and this agrees with the old definition by a "qualifying exam problem."

Then, if we define $\dim X := \text{tr.deg} k(X)$, it is easy to see that $\dim X = \dim U$ for an open subset $U \subset X$.

Theorem 115. *If X is affine and irreducible, then $\dim X = \text{Kr.dim} k[X]$.*

Proof. We use Noether normalization. Then there exists $k[x_1, \dots, x_n] \subset k[X]$, which is an integral extension. First it is easy to see that $\dim X = \dim \mathbb{A}^n = n$ because the field extension is algebraic.

To see that the Krull dimensions are equal, any chain $\mathfrak{q}_0 \subset \mathfrak{q}_1 \subsetneq \dots \subsetneq \mathfrak{q}_r \subsetneq k[x_1, \dots, x_n]$ gives a chain $\mathfrak{p}_0 \subsetneq \dots \subsetneq \mathfrak{p}_r \subsetneq k[X]$ by going-up. However, all \mathfrak{p}_i are distinct, so their intersections with $k[x_1, \dots, x_n]$ are distinct (quotient by \mathfrak{p}_i , then take the field of fractions of the smaller ring).

Now we show that the Krull dimension of $k[x_1, \dots, x_n]$ is n . Clearly it is at least n because we have the following chain of prime ideals: $(0) \subset (x_1) \subset \dots \subset (x_1, \dots, x_n)$. Take a chain $\mathfrak{p}_0 \subsetneq \dots \subsetneq \mathfrak{p}_s$. Then $V(\mathfrak{p}_s) \subsetneq \dots \subsetneq V(\mathfrak{p}_0) \subsetneq \mathbb{A}^n$. We prove a lemma which implies what we want. \square

Lemma 116. *Let $X \subset Y$ be irreducible affine varieties. Then $\dim X \leq \dim Y$. If $X \neq Y$, then the inequality is strict.*

Proof. Let $X \subsetneq Y \subsetneq \mathbb{A}^N$ and let $n = \dim Y$. Then given $t_1|_Y, \dots, t_n|_Y$, any $n+1$ are algebraically independent. However, $t_1|_X, \dots, t_n|_X$ generate $k[X]$ and therefore $k(X)$, so $\dim X \leq n$.

Suppose $\dim X = \dim Y = n$. Then some n coordinates $t_1|_X, \dots, t_n|_X$ are algebraically independent in $k(X)$. Thus $t_1|_Y, \dots, t_n|_Y$ are also algebraically independent. Choose $0 \neq u \in k[Y]$ such that $u|_X = 0$. Then there is a relation $a_0(t_1, \dots, t_n)u^m + \dots + a_m(t_1, \dots, t_n)$ which vanishes along Y . We can also assume that $a_m \neq 0$ along Y . However, it becomes 0 when restricted to X because $u|_X = 0$. \square

We use the Krull Principal Ideal Theorem:

Theorem 117 (Krull Principal Ideal). *If X is affine, irreducible, and $f \in k[X]$, then all irreducible components of $V(f) \subset X$ have codimension 1.*

Corollary 118. *In \mathbb{A}^n , irreducible hypersurfaces correspond exactly to irreducible polynomials $f \in k[x_1, \dots, x_n]$.*

Proof. Let $f \in k[x_1, \dots, x_n]$ be irreducible. Then because $k[x_1, \dots, x_n]$ is a UFD, (f) is prime, so $V(f)$ is irreducible and has the correct dimension by PIT.

In the other direction, let $X \subset \mathbb{A}^n$ be an irreducible hypersurface. Then choose $f \in k[x_1, \dots, x_n]$ such that $f|_X = 0$. Factor $f = f_1 \dots f_r$ into irreducibles. Then $X \subset V(f_i)$ because $V(f_i)$ are irreducible hypersurfaces. Because they have the same dimension, they are equal. \square

Corollary 119. Let $X \subset \mathbb{P}^n$ be projective and $F \in k[x_0, \dots, x_n]$ be homogeneous of positive degree d . Then $X \subset V(F)$ is non-empty and all irreducible components have codimension 1.

Proof. Work in the charts. Then use the PIT on the irreducible affine components. To show that the intersections in the affine charts are nonempty, pass to the affine cone. By the PIT, the intersection has codimension 1 and is non-empty because it contains the origin. Then the same is true for X . \square

Remark 120. This implies that if $X, Y \subset \mathbb{P}^2$ are projective curves, $X \cap Y$ are nonempty. This implies that every irreducible curve of degree at least 3 has inflection points.

Theorem 121. Let $f : X \rightarrow Y$ be a regular surjective map of irreducible varieties. Suppose $\dim X = n, \dim Y = m$. Then:

1. $\dim F \geq n - m$ for every component F of every fiber $f^{-1}(y)$.
2. There exists $U \subset Y$ open, nonempty such that $\dim F = n - m$ for all $y \in U$.
3. Sets $Y_k = \{y \in Y \mid \dim f^{-1}(y) \geq k\}$ are closed.

Proof. To prove the first part, take $y \in Y$. Then we can take. We can assume X, Y are affine by taking affine charts. Choose $f_1 \neq 0$ such that $f_1(y) = 0$. Then $\dim(f_1 = 0) = m - 1$ by the PIT. Choose $f_2|_D \neq 0$ ($f_2(y) = 0$) for all irreducible components D of $V(f_1)$. Then $\dim(f_1 = f_2 = 0) = m - 2$. Then $y \in (f_1 = \dots = f_m = 0)$, which is a finite set of points. Now we pass from Y to $D(F) \subset Y$ where $F(\tilde{y}) = 0$ for all $y \in (f_1 = \dots = f_m = 0) \setminus \{y\}$. With this new $Y, y = (f_1 = \dots = f_m = 0)$ where $m = \dim Y$, we see that $f^{-1}(y) = V(f^*(f_1), \dots, f^*(f_m))$. By the PIT, the dimension is at least $n - m$. \square

4.5 Lecture 12 (Mar 7) Last time we discussed Krull's Principal Ideal Theorem and began the proof of the theorem on dimension of fibers.

Corollary 122. Let X be irreducible of dimension n and $f_1, \dots, f_r \in k[X]$. Then every irreducible component of $V(f_1, \dots, f_r) \subset X$ has dimension at least $n - r$.

Sketch. We induct on r . If $r = 1$, we use the P.I.T. The rest is left as an exercise. \square

Conclusion of proof of Theorem 121. To prove the second part, we do something similar to the below theorem. We may assume X, Y are affine and $f : X \rightarrow Y$ dominant. Then we can decompose $f^* : k[Y] \hookrightarrow k[Y][z_1, \dots, z_r] \hookrightarrow k[X]$. This corresponds to a map $Y \xleftarrow{\pi_1} Y \times \mathbb{A}^r \xleftarrow{g} X$. Then $f^{-1}(y) = g^{-1}(Y \times \mathbb{A}^r)$.

We will show that there exists $D(F) \subset Y$ such that for all $y \in D(F)$, $p_1|_{f^{-1}(y)}, \dots, p_\ell|_{f^{-1}(y)}$ are algebraic over $z_1|_{f^{-1}(y)}, \dots, z_r|_{f^{-1}(y)}$, so every component of $f^{-1}(y)$ has dimension at most r . By part 1, the dimension is at least r , so it must equal r .

To prove the claim, write the equations of algebraic dependence $F_i(p_i, z_1, \dots, z_r, q_1, \dots, q_s) = 0$ with p_i appearing in the polynomial. Then restrict to a specific y and show that not all coefficients become zero (which is because there is an open set where the product of all coefficients does not vanish).

Now we prove the third part, we induct on the dimension of Y . If Y is a point, there is nothing to prove. Also, by part (1), $Y_{n-m} = Y$ is closed. By part (2), there exists $U \subset Y$ open where $\dim f^{-1}(y) = n - m$ for all $y \in U$. Then Y_j for $j > n - m$ is contained in $f^{-1}(Y \setminus U) = Y'$. Now consider f restricted to $f^{-1}(Y')$. Because $\dim Y' < \dim Y$, Y_j is closed. \square

Theorem 123. *Let $f : X \rightarrow Y$ be a regular map of quasi-projective varieties. Then $f(X)$ contains an open subset $U \subset \overline{f(X)}$.*

Proof. We may assume that f is dominant and that Y is irreducible by considering the irreducible components of Y . Considering the irreducible components of X , we may assume that X is irreducible. Passing to an affine chart $U \subset Y$ and its preimage, we may assume that Y is affine and irreducible. Take any affine chart in the preimage of U and we may assume that X is affine.

Thus we have $X \xrightarrow{X} Y$ and $f^* : k[Y] \hookrightarrow k[X]$ a map of domains. Then we can decompose $f^* : k[Y] \hookrightarrow k[Y][z_1, \dots, z_r] \hookrightarrow k[X]$. This corresponds to a map $Y \xrightarrow{\pi_1} Y \times \mathbb{A}^r \xrightarrow{g} X$. If α is integral, then g (and thus f) is surjective.

We use a trick that there exists a principal affine $D(F) \subset Y \times \mathbb{A}^r$ such that $g|_{g^{-1}(D(F))} \rightarrow D(F)$ is finite. Let $p_1, \dots, p_\ell \in k[X]$ be generators of the algebra. Then they are algebraic over $k[Y][z_1, \dots, z_r]$ so we can write $a_0 p_i^{m_i} + a_1 p_i^{m_i-1} + \dots + a_{m_i} = 0$, where $a_i \in k[Y][z_1, \dots, z_r]$. If we invert $F = a_0^1 a_0^2 \dots a_0^\ell$, then p_i are all integral over $k[Y][z_1, \dots, z_r][1/F] = k[D(F)]$.

Then the image of g contains $D(F)$ because finite maps are surjective. Then $D(F) \hookrightarrow Y \times \mathbb{A}^r \rightarrow Y$. We find $U \subset Y$ such that $U \subset \pi_1(D(F))$. Note that $F = \sum b_{i_1 \dots i_r} z_1^{i_1} \dots z_r^{i_r}$. Define $U = D(b_{i_1 \dots i_r})$ for some coefficient. If $y \in U$, then $F(y, z_1, \dots, z_r) \neq 0$ for some choice of y, z_1, \dots, z_r . Thus $(y, z_1, \dots, z_r) \in D(F)$. \square

Theorem 124. *Let $f : X \rightarrow Y$ be a regular map between projective varieties such that $f(X) = Y$ and suppose Y is irreducible. Suppose $f^{-1}(y)$ is irreducible of the same dimension ($\dim X - \dim Y$). Then X is irreducible.*

Proof. Decompose $X = \cup X_i$. Then $f(X_1) = \dots = f(X_s) = Y$ and $f(X_i) \subsetneq Y$ for all $i > s$. Define $f_i = f|_{X_i}$. Then there exists $U_i \subset Y$ open such that $f^{-1}(y)$ has smallest dimension n_i . Set $U = \cap_{i=1}^s U_i \setminus \cup_{j>s} f(X_j)$. Fix $y_0 \in U$. Then $f^{-1}(y_0) \subset \cup f_i^{-1}(y_0)$. WLOG we see that $f^{-1}(y_0) = f_1^{-1}(y_0)$, which implies $n = n_1$. Then for $y \in Y$, we see that $f^{-1}(y) \subset f_1^{-1}(y)$. Because Y is irreducible, $X = X_1$. \square

Jenia apologizes that this is too abstract and wants to calculate some actual examples.

Theorem 125. *Every cubic surface contains a line.*

Proof. We postulate the existence of the Grassmanian. One property of $\text{Gr}(k, n)$ is that the incidence variety $\mathcal{L} = \{p \in \mathbb{P}^{n-1}, L \in \text{Gr}(k, n) \mid p \in L\} \subset \mathbb{P}^{n-1} \times \text{Gr}(k, n)$ is closed. It is easy to see that \mathcal{L} is irreducible because $p_2 : \mathcal{L} \rightarrow \text{Gr}(k, n)$ and all fibers are \mathbb{P}^{k-1} .

Define $\text{Cub} = \mathbb{P}^{19} = \mathbb{P}(\text{Sym}^3(k^4))$ and $W = \{p \in \mathbb{P}^3, S \in \text{Cub} \mid p \in S\}$. Clearly W is a projective variety. Define $Z = \{L \in \text{Gr}(2, 4), S \in \text{Cub} \mid L \subset S\} \subset \text{Gr}(2, 4) \times \text{Cub}$. This is also a closed set

(construct a map such that Z is the locus of points with 1-dimensional fibers). Therefore Z is projective.

Now consider $\pi_2(Z) \subset \text{Cub}$. Because the Grassmanian is projective, it is proper. Then $\pi_2(Z)$ is closed. This is the set of cubic surfaces containing at least one line. If the image is not all cubic surfaces, we may use the theorem on the dimension of the fibers. We see that each cubic contains a positive-dimensional set of lines. But the Clebsch cubic surface contains only 27 lines, which is a contradiction. \square

5 LOCAL PROPERTIES

5.1 Lecture 13 (Mar 19) We begin our discussion of local properties. Let $p \in X$ be a point on a quasi-projective variety. We can assume X is affine and $k[X] = A$. Then $\mathfrak{m}_p \subset A$ is a maximal ideal. The local ring is $\mathcal{O}_p = A_{\mathfrak{m}_p}$, which is independent of the affine chart. If X is irreducible, then $\mathcal{O}_p \subset k(X)$. If X is not irreducible, then A has zero-divisors and localization is harder to define.

5.1.1 Tangent space Let $X \subset \mathbb{A}^n$. Then $X = V(I)$, $I = (F_1, \dots, F_s)$, $p = (x_1^0, \dots, x_n^0)$.

Definition 126 (Tangent Line). We say that a line L passing through p is tangent to X at p if the multiplicity of $L \cap X$ is at least 2.

Definition 127 (Tangent space, ambient version). The tangent space $T_p X$ is the locus of all lines tangent to X at p .

Definition 128 (Multiplicity). Let L be defined parametrically. Then $I|_L \subset k[t]$ is a principal ideal (f) . Then $f = t^m u$, where $u(0) \neq 0$. The multiplicity of intersection is m .

Example 129. Let $X = (y = x^2)$ the parabola. Then if $L = \begin{cases} x = x_0 + at \\ y = y_0 + bt \end{cases}$ we see that $f(t) = (b - 2ax_0)t + O(t^2)$. Thus the line is tangent if $b = 2ax_0$, or $b/a = 2x_0$.

Example 130. Let $X = (y^2 = x^3)$ and $p = x_0$. Then $f(t) = (bt)^2 - (at)^3 = t^2(b^2 - a^3t)$, so every line is a tangent line through the origin, and the tangent space is \mathbb{A}^2 .

Again let $I = (F_1, \dots, F_s)$ and $p = (x_1^0, \dots, x_n^0)$. Then

$$F_\ell(x) = F_\ell(p) + \sum \frac{\partial F_\ell}{\partial x_i}(p)(x_i - x_i^0) + \text{h.o.t.},$$

where $\sum \frac{\partial F_\ell}{\partial x_i}(p)(x_i - x_i^0)$ is the linearization of $d_p F_\ell$. Then $F_\ell|_L = \frac{\partial F_\ell}{\partial x_i}(p)a_i t + \text{h.o.t.}$ Thus L is a tangent line if and only if

$$\sum_{i=1}^n \frac{\partial F_\ell}{\partial x_i}(p)a_i = 0$$

for all $\ell = 1, \dots, s$. Thus the defining equations for $T_p X$ are

$$\sum \frac{\partial F_\ell}{\partial x_i}(p)(x_i - x_i^0), \ell = 1, \dots, s.$$

Note that $T_p X$ is an affine subspace of \mathbb{A}^n . One can also view the linear subspace parallel to $T_p X$ as the tangent space.

Theorem 131. *There is a canonical isomorphism $T_p^* \simeq \mathfrak{m}_p/\mathfrak{m}_p^2$, where either we take $\mathfrak{m}_p \subset A = k[X]$ or $\mathfrak{m}_p \subset \mathcal{O}_p$. This is known as the Zariski cotangent space.*

Proof. Let $g \in \mathfrak{m}_p$. Then $g = G + I$ for some $G \in k[x_1, \dots, x_n]$. Then $d_p G = \sum \frac{\partial G}{\partial x_i} (x_i - x_i^0)$, but for all $F \in I$, $d_p F|_{T_p X} = 0$. Therefore we can define $d_p g = d_p G|_{T_p X}$. This gives a map $d_p : \mathfrak{m}_p \rightarrow T_p^*$. Then d_p is clearly k -linear and surjective (linear function is its own differential). Then $\mathfrak{m}_p^2 \subset \text{Ker} d_p$ because \mathfrak{m}_p^2 is generated by $(x_i - x_i^0)(x_j - x_j^0)$, which vanish under the differential by Leibniz.

Thus d_p induces a surjective map $\mathfrak{m}_p/\mathfrak{m}_p^2 \rightarrow T_p^*$. We show that the kernel of d_p is \mathfrak{m}_p^2 . WLOG let $p = (0, \dots, 0)$. Suppose $d_p g = 0$. Then $g = G|_X$, $G \in k[x_1, \dots, x_n]$ and $d_p G|_{T_p} = 0$. Then $T_p = \{d_p F_\ell = 0 \mid \ell = 1, \dots, s\}$, so $d_p G = \lambda_1 d_p F_1 + \dots + \lambda_s d_p F_s$. Set $G_1 = G - \sum_{i=1}^s \lambda_i F_i$. Then $g_1|_X = g$ and $d_p G_1 = 0$, so $G_1 \in (x_1, \dots, x_n)^2$. Therefore $G_1|_X \in \mathfrak{m}_p^2$. \square

Corollary 132. T_p and therefore its dimension is a local invariant of $p \in X$.

Remark 133. Suppose $X \subset \mathbb{P}^n$ is projective. Then $X = (F_1 = \dots = F_s = 0)$. Choose $p \in X$, and suppose WLOG that $p \in \mathbb{A}_0^n$, so $p = (1 : x_1^0 : \dots : x_n^0)$. Then $T_p X \subset \mathbb{A}_0^n$ is given by $\sum_{i=1}^n \frac{\partial F_\ell}{\partial x_i}(p)(x_i - x_i^0) = 0$ where $\ell = 1, \dots, s$.

We can take $\overline{T_p X} \subset \mathbb{P}^n$ the projective tangent space. Recall Euler's formula $\sum_{i=0}^n \frac{\partial F_\ell}{\partial x_i}(p)x_i^0 = (\text{deg} F_\ell)F_\ell(p) = 0$. Therefore the projective tangent space is given by

$$\sum_{i=1}^n \frac{\partial F_\ell}{\partial x_i}(p)x_i = 0, \ell = 1, \dots, s.$$

Theorem 134. *Suppose X is irreducible. Then $X^{\text{sm}} = \{p \in X \mid \dim T_p X = \dim X\}$ is a non-empty open subset of "smooth" or "non-singular" points. The complement is known as the singular locus X^{sing} , and for all $p \in X^{\text{sing}}$, $\dim T_p X > \dim X$.*

Proof. We assume $X \subset \mathbb{A}^N$ is affine. Then define $TX = \{(a, p) \in \mathbb{A}^N \times X \mid a \in T_p X\}$. This is called the tangent bundle (or fiber space). Its equations are F_1, \dots, F_s the equations of X and the equations of tangency $\sum \frac{\partial F_\ell}{\partial x_i}(x_1, \dots, x_n)(y_i - x_i) = 0$. Thus TX is algebraic.

Now the second projection $\pi_2 : TX \rightarrow X$ has fiber the tangent space of X . By the theorem on the dimension of fibers, $X^{\text{sm}} = \{p \in X \mid \dim T_p X = s\}$, where s is the minimum possible, is an open subset. It remains to be seen that $s = n$.

Note every irreducible algebraic variety is birational to a hypersurface. Also, we show that X is birational to Y , if and only if there exist $U \subset X, V \subset Y$ open, nonempty such that $U \simeq V$.²⁰

Given both claims, we can assume $X \subset \mathbb{A}^{n+1}$ is a hypersurface. Then $X = (F = 0)$ for some irreducible $F \in k[x_0, \dots, x_{n+1}]$. Then $T_p X$ is given by $\sum_{i=0}^n \frac{\partial F}{\partial x_i}(p)(x_i - x_i^0) = 0$. Then the dimension of the tangent space is n unless $\frac{\partial F}{\partial x_i}(p) = 0$ for all i . This implies that $\frac{\partial F}{\partial x_i} \in (F)$

²⁰The second claim is on the midterm.

and thus the partial derivatives are zero. In characteristic 0, this implies $F = 0$, which is a contradiction, and in positive characteristic, F is a p th power, so it is not irreducible. Thus we have a contradiction. \square

Lemma 135. *Every irreducible variety is birational to a hypersurface.*

Proof. We work in characteristic zero. Choose a transcendence basis $k(x_1, \dots, x_n) \subset k(X)$ a finite extension. This extension is separable, so we use the primitive element theorem. Then $k(x_1, \dots, x_n, y)$, where y has minimal polynomial $F \in k[x_1, \dots, x_n, y]$. Thus $k(X) \simeq k(H)$, where $H = V(F)$. \square

If X is reducible, then $p \in X$ is nonsingular if $\dim T_p X = \max(\dim Y)$ where $Y \ni p$ is an irreducible component of X . In fact, there is a theorem that if p lies on several irreducible components, then p is singular.

5.2 Lecture 14 (Mar 21) Last time we considered local properties at a point. Today we will prove more theorems. Let p be a non-singular point of X .

Theorem 136. $\mathcal{O}_p \hookrightarrow k[[u_1, \dots, u_n]]$.

Corollary 137. \mathcal{O}_p is an integral domain. Thus X has only one irreducible component passing through p .

Remark 138. This is an analogue of the result from complex analysis that holomorphic functions are analytic and the map $\mathcal{O}_U \rightarrow \mathbb{C}[[z]]$ is injective.

Definition 139. $u_1, \dots, u_n \in \mathfrak{m}_p$ are called local parameters if $\{u_i\}$ is a basis of $\mathfrak{m}_p/\mathfrak{m}_p^2$. Equivalently, $d_p u_1, \dots, d_p u_n$ form a basis of T_p^\vee .

Fix $u_1, \dots, u_p \in \mathfrak{m}_p$ a system of local parameters.

Lemma 140. $\mathfrak{m}_p = (u_1, \dots, u_n)$.

Proof. Let $U = (u_1, \dots, u_n) \subset \mathfrak{m}_p$. Then $\mathfrak{m}_p(\mathfrak{m}_p/I) = \mathfrak{m}_p/I$, so by Nakayama's lemma, $\mathfrak{m}_p/I = 0$. \square

Definition 141. A formal power series $\phi = \sum_{k \geq 0} \phi_k(u_1, \dots, u_n)$ is called a Taylor series of $f \in \mathcal{O}_p$ if $(f - \sum_{k=0}^n \phi_k) \in \mathfrak{m}_p^{n+1}$.

Lemma 142. *Every $f \in \mathcal{O}_p$ has a Taylor series.*

Proof. Note $f = f(p) + g_1$, where g_1 vanishes at p , so $g_1 \in \mathfrak{m}_p$. Then we can write $g_1 = \sum_{i=1}^n a_i u_i$ where $a_i \in \mathcal{O}_p$. Thus we can write $a_i = a_i(p) + \bar{a}_i$, where $\bar{a}_i \in \mathfrak{m}_p$. Then we can write $g_1 = \sum_{i=1}^n a_i(p)u_i + \sum \bar{a}_i u_i$. Set the second sum to be $g_2 \in \mathfrak{m}_p^2$.

Now suppose $g_k \in \mathfrak{m}_p^k$. Thus $g_k = \sum_i h_1^i \cdots h_k^i$. Noting that $h_j^i = \sum a_{j,k}^i u_k$. Decompose this as above and then the claim follows by induction. \square

Lemma 143. *A Taylor series is unique. Thus we have a morphism $\mathcal{O}_p \rightarrow k[[u_1, \dots, u_n]]$.*

Lemma 144. *This morphism is injective.*

Proof. Take $f \in \mathcal{O}_p$ and suppose f has Taylor series 0. This happens if and only if $f \in \mathfrak{m}_p^n$ for every n . We know that by Krull's intersection theorem, if (R, \mathfrak{m}) is a Noetherian local ring and $I \subset R$ is some ideal, then $\bigcap_{n \geq 1} I^n = 0$. Therefore $f = 0$. \square

Theorem 145 (Krull's Intersection Theorem). *If (R, \mathfrak{m}) is a Noetherian local ring and $I \subset R$ is some ideal, then $\bigcap_{n \geq 1} I^n = 0$.*

Proof of Krull's intersection Theorem. Consider the Rees ring $\tilde{R} = R \oplus I \oplus I^2 \oplus \dots$ with graded multiplication. Then \tilde{R} is Noetherian. Consider $N = \bigcap_{n \geq 1} I^n$ and $\tilde{N} = N \oplus N \oplus N \oplus \dots \subset \tilde{R}$, which is an ideal. Thus \tilde{N} has a finite generating set. Thus all generators live in the first k components for some k . Then any $n \in N_{k+1}$ can be written $n = \sum \tilde{r}_i n_i$ where $\tilde{r}_i \in I \oplus I^2 \oplus \dots$. Then $N = I \cdot N$. I is contained in a maximal ideal, so $N = \mathfrak{m}N$, and therefore $N = 0$ by Nakayama. \square

Remark 146. This leads to the Artin-Rees lemma.

Proof of Lemma 143. Suppose $f \in \mathcal{O}_p$ has 2 different Taylor series. Equivalently, suppose 0 has a nontrivial Taylor series. Then $\phi_0 \in \mathfrak{m}_p$, so $\phi_0 = 0$. Then $\phi_1 = \sum a_i u_i \in \mathfrak{m}_p^2$. However, u_1, \dots, u_n are linearly independent mod \mathfrak{m}_p^2 by definition of local parameters. Thus $a_1 = \dots = a_n = 0$. Thus $\phi_i = 0$.

Now $\phi_2 = \sum a_{ij} u_i u_j \in \mathfrak{m}_p^3$. We show that $\phi_k(u_1, \dots, u_n) \in \mathfrak{m}_p^{k+1}$ if and only if ϕ_k is a trivial polynomial. If $n = 1$, then $\mathfrak{m}_p = (u)$ and $\phi_k(u) = \alpha u^k \in (u)^{j+1}$. If $\alpha \neq 0$, then $(u)^k = (u)^{k+1}$, which implies $(u) = 0$ by Nakayama.

From the proof of Noether's normalization, WLOG $\phi_k = u_1^k + \text{other monomials}$. We restrict to a hypersurface $Y = V(u_n) \subset X$. Then all irreducible components of Y have codimension 1 by the PIT. We claim that Y is non-singular at p and $u_1|_Y, \dots, u_{n-1}|_Y$ are local parameters of Y at p .

Given the claim, $\phi_k|_Y = \phi(u_1|_Y, \dots, u_{n-1}|_Y, 0)$ is a nontrivial polynomial. It contains $(u_1|_Y)^k$. On the other hand, $\phi_k \in \mathfrak{m}_p^{k+1}$, so $\phi_k|_Y \in (\mathfrak{m}_{p,Y})^{k+1}$. By induction, this is a contradiction. \square

Theorem 147. *Let $u_1, \dots, u_n \in \mathfrak{m}_p \subset \mathcal{O}_p$ be local parameters. Shrink X so that $u_1, \dots, u_n \in k[X]$. Then $X_i = V(u_i)$ is nonsingular at p and $\{u_j|_{X_i}\}_{j \neq i}$ are local parameters of X_i . Also, $\bigcap_{i=1}^n T_{p,X_i} = 0$.*

Proof. Note $u_i \in I(x_i) \subset k[X]$. Then $T_p X_i = \{df_p = 0 \text{ for all } f_p \in I(x_i)\} \subset L_i = \{d_p u_i = 0\} \subset T_p X$. By the PIT, $\dim_p X_i = n - 1$. However, $\dim T_p X_i \leq n - 1$. Thus x_i is smooth and $T_p X_i = L_i$. Also, $\{d_p u_j|_{L_i}\}_{j \neq i}$ is a basis of L_i^\vee . Therefore $\{u_j|_{X_i}\}_{j \neq i}$ are local parameters on X_i . \square

This completes the proof of Lemma 143.

5.2.1 Tangent Cone Consider the cusp $X = \{y^2 = x^3\}$, $T_0 X = \mathbb{A}^2$. Rescale $x \rightarrow tx, y \rightarrow ty$, so the equation becomes $t^2 y^2 = t^3 x^3$, so $y^2 = tx^3$. Note that $X \simeq X_t$ for $t \neq 0$. Then we set $X_0 = \{y^2 = 0\}$.

Let $0 \in X \subset \mathbb{A}^n$. Then $X = V(I)$. For all $f \in I$ we can write $f = \text{in}(f) + \text{h.o.t.}$. Then define $I_0 = \text{in}(I) = (\text{in}(f) \mid f \in I)$. For example, $\text{in}(y^2 - x^3) = (y^2)$.

Definition 148. The Tangent cone is defined to be $V(I_0)$, at least if I_0 is a radical ideal.

Theorem 149. *The tangent cone has the same dimension as the original variety. There is a family of varieties $X_t \simeq X$ such that “ $\lim_{t \rightarrow 0} X_t$ ” = X_0 .*

More precisely, there exists an affine variety $\mathcal{X} \subset \mathbb{A}^n \times \mathbb{A}_t^1$ such that $\pi_2^{-1}(t) \simeq X$ for $t \neq 0$, $\pi_2(1) = X$, and $\pi_2^{-1}(0) = X_0$.

5.3 Lecture 15 (Mar 26) We can consider lines (or curves on a variety X as maps $\mathbb{P}^1 \dashrightarrow X$). Also, we can consider maps $f : X \dashrightarrow \mathbb{P}^1$. Then $Y = f^{-1}(a) \subset X$ is a hypersurface.

Definition 150 (Locally Principal). A hypersurface $Y \subset X$ is called locally principal if it is locally given by 1 equation.

Definition 151 (Local Equations). $f_1, \dots, f_r \in \mathcal{O}_p$ are called local equations of a subvariety $Y \subset X$ if there exists an affine neighborhood $U \subset X$ of p such that $f_1, \dots, f_r \in k[U]$ and $I(Y) = (f_1, \dots, f_r)$.

Lemma 152. f_1, \dots, f_r are local equations of Y at p if and only if $\mathcal{L}_{Y,p} \subset \mathcal{O}_p$ is generated by f_1, \dots, f_r . Here $\mathcal{L}_{Y,p} = \{f \in \mathcal{O}_p \mid f|_{Y \cap U} = 0 \text{ for some neighborhood } p \in U\}$. Equivalently, if X is already affine, $I(Y) \subset k(X)$ and $\mathcal{L}_{Y,p} = I(Y)_{\mathfrak{m}_p}$.

Proof. If $f_1, \dots, f_r \in \mathcal{O}_p$ are local equations, then after shrinking X to U , we have $I(Y) = (f_1, \dots, f_r) \subset k[X]$. Then $\mathcal{L}_{Y,p} = (f_1, \dots, f_r)$ because localization preserves generators.

Now suppose that $\mathcal{L}_{Y,p} = (f_1, \dots, f_r) \subset \mathcal{O}_p$. By shrinking, we can assume that $f_1, \dots, f_r \in k[X]$ and that X is affine. We know that $I(Y) = (g_1, \dots, g_s) \subset K[X]$. We can write $g_i = \sum_{j=1}^r h_{ij} f_j$ in \mathcal{O}_p . We can shrink X : Choose a principal open set $U = D(w) \subset X$ such that $h_{ij} \in k[U]$. We claim that in U , $I(Y \cap U) = (g_1, \dots, g_s) \subset k[U] = k[X][1/w]$ is also generated by f_1, \dots, f_r . We know $(g_1, \dots, g_s) \subset (f_1, \dots, f_r)$. Then $f_i|_Y = 0$, so $(f_1, \dots, f_r) \subset (g_1, \dots, g_s)$. \square

Theorem 153 (Criterion for Smoothness). *Let X be a variety of dimension n . Then suppose $p \in Y \subset X$ and Y is locally principal at p , as in $\mathcal{L}_{Y,p} = (g) \subset \mathcal{O}_p$. If Y is nonsingular at p , then so is X .*

Proof. Shrink X to be affine. Then $I(Y) = (g) \subset k[X]$. We see that $I(X) = (F_1, \dots, F_s)$ and $\overline{I(Y)} = (F_1, \dots, F_s, G)$. Recall that $T_p X = \{d_p F_1, \dots, d_p F_s = 0\}$ with dimension at least n . Then $T_p Y = \{d_p F_1 = \dots = d_p F_s = d_p G = 0\}$. Then we see that $T_p Y$ has dimension $n - 1$, so $T_p X$ must have dimension n , so X is nonsingular at p . \square

Remark 154. It is not enough to assume that $Y = V(g)$. For example, take $X = \{z^2 = xy\}$ and $Y = V(x)$. Y is nonsingular at the origin but X is singular. In fact, this implies that $Y \subset X$ is not locally principal, so $I(Y) = (x, z)$ needs at least 2 generators.

Theorem 155. *Suppose X is nonsingular at p and suppose $p \in Y \subset X$ is an irreducible hypersurface. Then Y is locally principal at p .²¹*

Theorem 156 (Equivalent Theorem). *Let $p \in X$ be a nonsingular point. Then \mathcal{O}_p is a UFD.*

²¹This does not imply that $I(Y) \subset K[X]$ is principal. For an example, take $p \in E = \{y^2 = x^3 + ax + b\}$ where the discriminant is nonzero. Then $\mathfrak{m}_p \subset \mathcal{O}_p = (t)$ is principal, but $I(p) \subset k[E]$ is not principal. If $I(p) = (t) \subset k[E]$, then we have a map $f : E \rightarrow \mathbb{A}^1$ where $f^{-1}(0) = p$. This induces a map $f : \overline{E} \rightarrow \mathbb{P}^1$ such that $f(\infty) = \infty$. This implies f is birational, which is a contradiction.

Proof of Equivalence. Suppose $p \in Y \subset X$ where X is affine. Then consider $f \in \mathcal{L}_{Y,p}$. This is a UFD, so we can factor $f = f_1 \cdots f_r$. We can assume that $f \in k[X]$ and $f|_Y = 0$. If Y is irreducible, then f_1 vanishes on Y (up to renumbering). We will show that $\mathcal{L}_{Y,p} = (f_1 = g)$.

We know that $Y \subset V(g)$. We will show that Y is the only irreducible component of $V(g)$ passing through p . Suppose Y' is another component passing through p . Then there exists $h|_Y = h'|_{Y'} = 0$, and $hh'|_{V(g)} = 0$. Thus $g|(hh')^k$ for some k . Thus g divides h or h' in a UFD, which leads to a contradiction. This implies that $Y = V(g)$ is irreducible. Take $s \in I(Y)$. Then $g|s^k$ for some k . Because g is prime, it divides s . Thus $s \in (g)$. In fact, $I(Y) = (g)$.

In the other direction, note that \mathcal{O}_p is Noetherian, which implies existence of prime factorization. We prove Euclid's Lemma. Let g be irreducible. Let $Y = V(g)$. Then Y is irreducible at p . Take $Y_1 \subset Y$ an irreducible component. By Krull, Y_1 has codimension 1. Thus Y_1 is locally principal, so $Y_1 = V(g_1)$ after shrinking. We also know $I(Y_1) = (g_1)$. Then $g|_{Y_1} = 0$, so $g \in (g_1)$. Thus $g = g_1 h_1 \in \mathcal{O}_p$, so h_1 is a unit. Then $(g) = (g_1)$ and thus $Y = Y_1$. Thus Y is irreducible and g is prime. \square

Proof of Theorem 156. The idea is to use the injection $\mathcal{O}_p \subset k[[T_1, \dots, T_n]]$ where $p \in X$ nonsingular and $n = \dim X$. We check that Euclid's Lemma holds. We know that this holds in $k[[T_1, \dots, T_n]]$. Then if $a, c \in \mathcal{O}_p$ with $a|c$ in $\widehat{\mathcal{O}}_p$. We also check that if a, b are coprime in \mathcal{O}_p then they are coprime in $\widehat{\mathcal{O}}_p$. We show that if $I \subset \mathcal{O}_p$, then $(I\widehat{\mathcal{O}}_p) \cap \mathcal{O}_p = I$.

To prove the claim, let $I \subset \mathcal{O}_p$ where $I = (f_1, \dots, f_s)$ and $x \in (I\widehat{\mathcal{O}}_p) \cap \mathcal{O}_p$. Then $x = \sum f_i \alpha_i$, $\alpha_i \in \widehat{\mathcal{O}}_p$. Then for all $n \geq 1$ we see that $\alpha_i = a_i^{(n)} + \xi_i^{(n)}$ where $a_i \in \mathcal{O}_p$ and $\xi_i \in \widehat{m}_p^n$. Thus $x = \sum a_i^{(n)} f_i + \sum \xi_i^{(n)} f_i$. We know $\xi = x - a \in \widehat{m}_p^n \cap \mathcal{O}_p$, so $\xi \in m_p^n$. Therefore $x \in I + m_p^n$ for all n , and thus $x \in I$ by Krull's Intersection Theorem.

Given the claim, we see that $c \in (a)\widehat{\mathcal{O}}_p$. However, $c \in \mathcal{O}_p$, so $c \in (a)$. Now we show that gcd is preserved by the embedding. Suppose $a = \gamma\alpha, b = \gamma\beta$ where $(\alpha, \beta) \in \widehat{\mathcal{O}}_p$. Choose n such that $\alpha, \beta \notin \widehat{m}_p^n$. Then $\alpha = x_n + u_n, \beta = y_n + v_n$ where $x_n, y_n \in \mathcal{O}_p$ and $u_n, v_n \in \widehat{m}_p^n$. Because $a\beta - b\alpha = 0$, we see that $ay_n - bx_n = a(\beta - v_n) - b(\alpha - u_n) = -av_n + bu_n \in (a, b)\widehat{m}_p^n$. By the claim, $ay_n - bx_n \in (a, b)m_p^n$, so $ay_n - bx_n = a_n + bs_n$ where $t_n, s_n \in m_p^n$. By algebraic manipulation, $\alpha(y_n - t_n) = \beta(x_n + s_n)$. Because α, β are coprime, then $\alpha | x_n + s_n$, so $x_n + s_n = \alpha\lambda$. Thus $\lambda = 1 + h.o.t.$ is a unit. Therefore $x_n + s_n | \alpha$, so $x_n + s_n | a$ in \mathcal{O}_p . Then $a = (x_n + s_n)h$ and therefore $h(y_n - t_n) = b$, so $b = h(y_n - t_n)$. Therefore h is a unit. In the power series ring, $\alpha | a$ and $a | \alpha$, so γ is a unit. \square

5.4 Lecture 16 (Mar 28) We prove that if $p \in X$ is smooth, then \mathcal{O}_p is a UFD. We will finish the proof in last time's section.


Corollary 157. *Let $f : X \dashrightarrow Y$ is a rational map with X nonsingular, Y projective. Then the indeterminacy locus has codimension at least 2. In particular, if X is nonsingular, f is regular.*

Proof. We work near $p \in X$ and assume $Y = \mathbb{P}^n$. Then $f = (f_0 : \dots : f_n)$ where $f_i \in k(X)$. We can assume that $f_i \in \mathcal{O}_p$ with no common factor. We show that $V(f_0, \dots, f_n) \subset X$ has codimension 2 near p .

Suppose $Y \subset V(f_0, \dots, f_n)$ and Y has codimension 1. Then $\mathcal{L}_{Y,p} = (h)$ because \mathcal{O}_p is a UFD. Then $f_i|_Y = g_i h$. This contradicts the fact that the f_i have no common factor. \square

5.4.1 Blowups

Example 158. Consider $\xi = [1 : 0 : \dots : 0] \in \mathbb{P}^n$ and take the map $\pi : \mathbb{P}^n \dashrightarrow \mathbb{P}^{n-1}$ given by $[x_0 : \dots : x_n] \mapsto [x_1 : \dots : x_n]$. We know π is regular at $\mathbb{P}^n \setminus \xi$. We want to resolve π :



Consider the map $\mathbb{P}^n \times \mathbb{P}^{n-1} \supset \text{Bl}_\xi \mathbb{P}^n = \overline{\Gamma_\pi}$, where $\Gamma_\pi = \{(x, y) \mid y = \pi(x)\}$. By construction, $\sigma = \pi_1$ and the right arrow is π_2 . We also know that $\text{Bl}_\xi \mathbb{P}^n \setminus \sigma^{-1}(\xi) \simeq \mathbb{P}^n \setminus \xi$. Also, the blowup is irreducible.

Fix $[y_1 : \dots : y_n] \in \mathbb{P}^{n-1}$. We know $\pi^{-1}(y) \supset \{[1 : t_{y_1} : \dots : t_{y_n}]\}$. Then the closure contains $\xi, [y_1 : \dots : y_n]$. Therefore $\sigma^{-1}(\xi) = \mathbb{P}^{n-1}$ and is called the exceptional divisor of the blowup.

Now we find equations for the blowup. We know that $\mathbb{P}^n \times \mathbb{P}^{n-1}$ is given by $x_i y_j = x_j y_i$. Then if $x_1 = \dots = x_n = 0$ (ξ), we get \mathbb{P}^{n-1} . If $x_1 \neq 0$, then $[x_1 : \dots : x_n] \sim [y_1 : \dots : y_n]$ and we can use equations for Γ_π .

5.5 Lecture 17 (Apr 02) We finish the discussion of blowups from last time. Let $f : X \rightarrow Y$ be a birational regular surjective morphism. Then f restricts to an isomorphism $U \simeq V$ of open subsets. Using the theorem on dimension of the fibers, there exists an open $W \subset Y$ such that f has finite fibers over points in W . Away from W , every component of every fiber is positive dimensional.

Definition 159 (Exceptional Locus). $\text{Exc}(f) = f^{-1}(Y \setminus W)$ is the exceptional locus of f .

Theorem 160. Suppose $f : X \rightarrow Y$ is a surjective birational regular map. Suppose $f(x) = y$ and assume that Y is smooth at y (more generally, Y is locally factorial at y , meaning that \mathcal{O}_y is a UFD). Assume that $g = f^{-1}$ is not defined at y . Then there exists a subvariety $Z \subset X$ of codimension 1 passing through x such that $Z \subset \text{Exc}(f)$.

Corollary 161. If $f : X \rightarrow Y$ is birational, surjective, regular, and $\text{codim}(\text{Exc}(f)) \geq 2$, then Y is singular and in fact not even locally factorial.

Proof. Pass to an affine neighborhood of x at X , and assume that $X \subset \mathbb{A}^n$ with $g = f^{-1}$. Then denote $g_i = g^*(f_i) \in k(Y)$. Then g is not defined at $y \in Y$, so one of the g_i , say g_1 is not regular at $y \in Y$. Thus $g_1 \notin \mathcal{O}_y$.

If \mathcal{O}_y is a UFD we can write $g_1 = u/v$ where u, v are coprime and $v(y) = 0$. We know that t_1 is a regular function on X , where $t_1 = f^*(g^*(t_1)) = f^*(g_1) = \frac{f^*(u)}{f^*(v)}$.

Define $Z = V(f^*v) \subset X$, which is a hypersurface. We know $f^*(u)|_Z = (t_1 f^*v)|_Z = 0$. Thus $f(Z) \subset V(u) \cap V(v)$. We know $V(u, v) \subset Y$ has codimension 2 at y because the local ring at y is a UFD. \square

We observe that the blowup is a local construction. Take $\mathbb{A}^n_{x_1, \dots, x_n}$ and let $\xi = 0$. Then $\text{Bl}_\xi = \bar{\Gamma}_\pi \subset \mathbb{A}^n \times \mathbb{P}^{n-1} \subset \text{Bl}_\xi \mathbb{P}^n$. To find equations, we dehomogenize to find $x_i y_j = x_j$ if we take $y_i = 1$. Also, other equations are $x_j y_k = x_k y_j$. Also $\text{Bl}_\xi \mathbb{A}^n = \bigcup_{i=1}^n U_i$ where $U_i = \{y_i \neq 0\}$. Thus $U_i \simeq \mathbb{A}^n_{y_1, \dots, x_i, \dots, y_n}$ and there is a map $U_i \rightarrow \mathbb{A}^n$ given by $(y_1, \dots, x_i, \dots, y_n) \mapsto (x_i y_1, \dots, x_i, x_i y_n)$.

Corollary 162. *The blowup of \mathbb{A}^n is smooth, covered by \mathbb{A}^n 's.*

Definition 163 (Proper Transform). Take $X \subset \mathbb{A}^n$. Then $\sigma^{-1}(X) = E \cup \overline{\sigma^{-1}(X \setminus \xi)}$. the components are $E \simeq \mathbb{P}^n$ and \tilde{X} , which is called a proper transform of X .

Example 164. Consider $X = (x_2 = tx_1) \subset \mathbb{A}^2$. Then consider $U_1 \simeq \mathbb{A}^2_{x_1, y_2}$ and $U_2 = \mathbb{A}^2_{y_1, y_2} /$ Them if $\alpha : U_1 \rightarrow \mathbb{A}^2$ is the covering map, $\alpha^{-1}(x) = x_1 y_2 - tx_1 = 0$, so either $x_1 = 0$ or $y_2 = t$. On U_2 , we see $x_1 \mapsto x_2 y_1, y_2 \mapsto y_2$, so $\beta^{-1}(x) = x_2 - tx_2 y_1 = 0$, so either $x_2 = 0$ or $1 - ty_1 = 0$.

Example 165. Consider $X = (x_2^2 - x_1^3 = 0)$ in U_1 . Then $\alpha^{-1}(X) = (x_1 y_2)^2 - x_1^3 = 0$, so either $x_1 = 0$ (E) or $x_1 = y_2^2$ (\tilde{X}).

5.5.1 *Local Blowup II* Let $\xi \in X$ be a smooth point and the dimension of X be n . We find the blowup of X at ξ ,

Definition 166. Choose local parameters $u_1, \dots, u_n \in \mathfrak{m}_\xi$. Near ξ , a rational map $X \dashrightarrow \mathbb{P}^n$ given by $x \mapsto [u_1(x) : \dots : u_n(x)]$ is not regular only at ξ . Then $\text{Bl}_\xi X = \bar{\Gamma}_\pi \subset X \times \mathbb{P}^1$ which is irreducible, smooth and the map $\sigma : \text{Bl}_\xi(X) \rightarrow X$ is an isomorphism away from ξ with equations as before.

Theorem 167. *Let $\zeta \in X \subset \mathbb{A}^n$ with X smooth. Then $\tilde{X} \subset \text{Bl}_\xi \mathbb{A}^n$ is isomorphic to $\text{Bl}_\xi X$.*

Theorem 168. *$\text{Bl}_\xi X$ is independent of the choice of local parameters. For two choices of local parameters, there exists a unique isomorphism ψ that commutes with the σ .*

5.6 **Lecture 18 (Apr 04)** Recall that local parameters are essentially a substitute for homolorphic local coordinates in complex geometry.

Lemma 169 (Weak Inverse Function Theorem). *Let $p \in Y \subset X$ be a smooth point of both X and Y . Then there exist local parameters u_1, \dots, u_n such that $u_1|_Y, \dots, u_m|_Y$ are local parameters on Y and $I(Y) = (u_{m+1}, \dots, u_n)$ locally in some affine neighborhood of p . Equivalently, a smooth subvariety of a smooth variety is a local complete intersection.*

Proof. Observe the following short exact sequences:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
0 & \longrightarrow & I_{Y,X}/(\mathfrak{m}_{p,X}^2 \cap I_{Y,X}) & \longrightarrow & T_{p,X}^* & \longrightarrow & T_{p,Y}^* \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
0 & \longrightarrow & I_{Y,X} & \longrightarrow & \mathfrak{m}_{p,X} & \longrightarrow & \mathfrak{m}_{p,Y} \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
0 & \longrightarrow & \mathfrak{m}_{p,X}^2 \cap I_{Y,X} & \longrightarrow & \mathfrak{m}_{p,X}^2 & \longrightarrow & \mathfrak{m}_{p,Y}^2 \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
& & 0 & & 0 & & 0
\end{array}$$

Choose a basis u_1, \dots, u_m of $T_{p,Y}^*$ and lift to X . Then choose u_{m+1}, \dots, u_n which generate the quotient. This gives local parameters u_1, \dots, u_n in $\mathfrak{m}_{p,X}$ such that $u_1|_Y, \dots, u_m|_Y$ are local parameters on Y . Define $Y' = V(u_{m+1}, \dots, u_n)$. By previous results, Y' is smooth at p and has dimension m . Then $Y \subset Y'$ is smooth with the same dimension, so they must be equal in some neighborhood of p . \square

Remark 170. Y is a local complete intersection at p if $\mathcal{L}_Y \subset \mathcal{O}_p$ is generated by $s = \text{codim}_Y X$ equations.

Now let $p \in X$ be smooth and u_1, \dots, u_n be local parameters. Then $\bar{X} = \text{Bl}_p X \subset X \times \mathbb{P}^{n-1}$. We check that this is smooth. We know that $\text{Bl}_p X \setminus E \simeq X \setminus \{p\}$. Therefore we check for a point $y^0 \in E$. We show that $\mathfrak{m}_{y^0} \bar{X}$ is generated by n elements. Take the chart U_i of \bar{X} given by $U_i \subset X \times \mathbb{A}_{y_2, \dots, y_n}^{n-1}$. Then the equations are $u_j = u_1 y_j$. Then

$$\mathfrak{m}_{y^0} U_i = \mathfrak{m}_p X|_{U_i} + \mathfrak{m}_{(y_2^0, \dots, y_n^0)} \mathbb{A}^{n-1}|_{U_i} = (u_1, \dots, u_n, y_2 - y_2^0, \dots, y_n - y_n^0)|_{U_i} = (u_1, y_2 - y_2^0, \dots, y_n - y_n^0).$$

Thus \bar{X} is smooth at Y^0 .

Theorem 171. *Let $0 \in X \subset \mathbb{A}^n$ be a smooth point. The proper transform of X in $\text{Bl}_0 \mathbb{A}^n$ is isomorphic to $\text{Bl}_0 X$.*

Proof. Choose local parameters u_1, \dots, u_n such that $I(X) = u_{m+1}, \dots, u_n$ locally at 0 . Then $u_1|_X, \dots, u_m|_X$ are local parameters at $0 \in X$. We know the blowup is independent of the choice of local parameters.

Therefore $\text{Bl}_0 \mathbb{A}^n \subset \mathbb{A}^n \times \mathbb{P}^{n-1}$ with equations $u_i y_j = u_j y_i$ and $\text{Bl}_0 X \subset X \times \mathbb{P}^{m-1}$ is given by $u_i y_j = u_j y_i$ (with appropriate indices). To get equations of $\text{Bl}_0 X$ from equations of $\text{Bl}_0 \mathbb{A}^n$, take equations $u_{m+1} = \dots = u_n = y_{m+1} = \dots = y_n = 0$ to cut out X and \mathbb{A}^n .

To see that this is the proper transform \tilde{X} , note that $u_{m+1} = \dots = u_n$ along \tilde{X} because they cut $X \times \mathbb{P}^{n-1}$ from $\mathbb{A}^n \times \mathbb{P}^{n-1}$. Then choose one of the u_i , say $u_1 \neq 0$ at some part of $X - 0$. Then $u_1 y_j = u_j y_1$ for all j in $\text{Bl}_0 \mathbb{A}^n$. Since $u_j = 0$ for $j = m+1, \dots, n$, $y_j = 0$. \square

5.6.1 *Normal Varieties* Let $p \in X$ be an irreducible variety. Then X is normal at p if $\mathcal{O}_p \subset k(X)$ is integrally closed. Then X is normal if it is normal at every point.

Lemma 172. *If $p \in X$ is a smooth point, or factorial at p , then X is normal at p .*

Proof. \mathcal{O}_p is a UFD and any UFD is integrally closed. To see this suppose $f \in k(X)$, $f = ab$ where $a, b \in \mathcal{O}_p$ and a, b are coprime, is integral over \mathcal{O}_p . Then $f^s + a_1 f^{s-1} + \dots + a_s = 0$. Clearing denominators, we see that $a^s + a_1 a^{s-1} b + \dots + a_s b^s = 0$. Then $b \mid a^s$, so b is a unit. Thus $f \in \mathcal{O}_p$. \square

Lemma 173. *Let X be affine. Then X is normal if and only if $k[X] \subset k(X)$ is integrally closed.*

Proof. Let $R \subset f.f.R$ be integrally closed. Then we know that $R_p \subset f.f.R$ is integrally closed for every prime $p \subset R$.

In the other direction, note that $R_m \subset K$ is integrally closed. Then note that $R = \bigcap R_m$ where the intersection is over all maximal ideals m . Then if $x \in K$ is integral over R , it must be integral over R_m for all m , so it must be in R_m for all m . Thus $x \in R$.

To prove that $R = \bigcap_m R_m$, define $I = \{b \in R \mid xb \in R\}$. If $I = R$, then $1 \in I$ and thus $x \in R$. Otherwise, $I \subset m$ for some maximal ideal m . This is impossible because we can write $x = \frac{a}{b}$ where $b \notin m$ because $x \in R_m$. \square

Remark 174. Every irreducible surface in A^3 with isolated singularities is normal.

Theorem 175 (Serre’s Criterion). *Let R be a Noetherian domain and K the fraction field of R . Then $R \subset K$ is integrally closed if and only if*

$$R = \bigcap_{\substack{p \subset R \\ \text{minimal prime}}} R_p$$

and for all minimal primes $p \subset R$, R_p is a DVR.

Now let $R = k[X]$ for some affine X . Then we find when $p \subset R$ is a minimal prime (of height 1) if $\dim R/p = \dim R - 1$, or $Y = V(p) \subset X$ is a hypersurface.

Note that $R_p = \mathcal{O}_Y = \{f \in k(X) \mid f \text{ is regular at some point of } Y\}$. Then note that $\bigcap R_p = \{f \in k(X) \mid f \text{ is regular at some point of every hypersurface}\}$. Therefore $R = \bigcap R_p$ if and only if every rational function regular at some point of every hypersurface is regular everywhere.

Theorem 176 (Hartog’s Extension Principle). *Let $Z \subset X$ of codimension at least 2. Then if $f \in k(x)$ is regular on $X \setminus Z$, f is regular at some point of every hypersurface, so f is regular.*

5.7 Lecture 19 (Apr 09)

5.7.1 *Differential of a regular map* Let $\varphi : X \rightarrow Y$ be regular. Then we want to define $d_x \varphi : T_{X,x} \rightarrow T_{Y,y}$. We may assume that X, Y are affine, so choose charts. Then recall that $T_{X,x} = (d_x f_1 = \dots = d_x f_r = 0)$ where $I(X) = (f_1, \dots, f_r)$, so we define $d_x \varphi = (d_x \varphi_1, \dots, d_x \varphi_m)$.

Proposition 177. $d_x \varphi(T_{X,x}) \subset T_{Y,y}$.

Proof. $(d_y g_i)(d_x \varphi) = d_x(f_i \circ \varphi) = d_x 0 = 0$. \square

More intrinsically, the dual map $T_{Y,y}^\vee \rightarrow T_{X,x}^\vee$ is induced by the pullback $\varphi^* : \mathcal{O}_{Y,y} \rightarrow \mathcal{O}_{X,x}$.

Example 178. Consider $X = (xy = t) \subset \mathbb{A}^3$ and $\varphi : X \rightarrow \mathbb{A}^1$ given by $(x, y, t) \mapsto t$. Note that $T_p X = \{y_0 dx + x_0 dy - dt = 0\} \subset \mathbb{A}^3$. Therefore X is smooth and $T_p X$ has coordinates dx, dy (basis of $T_p X^\vee$) and $d\varphi = dt = y_0 dx + x_0 dy$, which is surjective unless $x_0 = y_0 = 0$.

To see the geometry of this map, we see that the fiber above every nonzero point is smooth (a torus), but the fiber at 0 is the union of two coordinate axes.

Lemma 179. Let $\varphi : X \rightarrow Y$ be a regular map of smooth varieties with $\varphi(x) = y$. Suppose that $d_x \varphi : T_{X,x} \rightarrow T_{Y,y}$ is surjective. Then $\varphi^{-1}(y)$ is nonsingular at x .

Proof. Consider the differential $d_x \varphi : T_{X,x} \rightarrow T_{Y,y}$. We show that $T_{\varphi^{-1}(y),x} \subset \text{Ker } d_x \varphi$. Given the claim, we see that $\dim T_{\varphi^{-1}(y),x} \leq \dim \text{Ker } d_x \varphi = \dim T_{X,x} - \dim T_{Y,y} = \dim X - \dim Y$. On the other hand, $\dim T_{\varphi^{-1}(y),x} \geq \dim_x \varphi^{-1}(y) \geq \dim X - \dim Y$. Thus all inequalities are equalities and $\varphi^{-1}(y)$ is smooth at x .

To prove the claim, consider the sequence of maps $T_{\varphi^{-1}(y),x} \xrightarrow{d_x t} T_{X,x} \xrightarrow{d_x \varphi} T_{Y,y}$. Then $T_{\varphi^{-1}(y),x} \rightarrow T_{Y,y}$ is a differential of the map $\varphi^{-1}(y) \rightarrow Y$. This is constant, so the differential is zero. \square

5.7.2 Normal Varieties Continued

Theorem 180. A nonsingular variety is normal.

Proof of this theorem comes from Lemma 173.

Definition 181 (Discrete Valuation). A discrete valuation on a field k is a function $v : k^* \rightarrow \mathbb{Z}$ such that $v(fg) = v(f) + v(g)$ and $v(x + y) \geq \min\{v(x), v(y)\}$.

Define $R = \{f \in K \mid v(f) \geq 0\}$ and $\mathfrak{m} = \{f \in R \mid v(f) > 0\}$.

Lemma 182. R is a local ring with maximal ideal \mathfrak{m} (called a DVR) with field of fractions k .

Proof. Clearly by definition of discrete valuation, R is a ring and \mathfrak{m} is an ideal. It is easy to see that everything with valuation 0 is a unit in R , so \mathfrak{m} is a maximal ideal. Then we see that for all $x \in K$, at least one of $v(x), v(x^{-1})$ is nonnegative, so at least one of x, x^{-1} is in R . \square

Lemma 183. Every DVR is a PID and in fact, every ideal has form (t^n) for some fixed $t \in \mathfrak{m} \setminus \mathfrak{m}^2$.

Proof. Choose t such that $v(t) = 1$. Then clearly $t \in \mathfrak{m}$. Take $I \subset R$ an ideal. Then let n be the minimum valuation of any element of I , so $v(g/t^n) \geq 0$ and thus $g \in (t^n)$. Then choose a minimizing f , and we see that f/t^n is a unit, so $t^n \in I$. \square

Theorem 184. Let R be a ring. Then the following are equivalent:

1. R is a DVR.
2. R is a Noetherian local domain which is integrally closed and has Krull dimension 1.

Corollary 185. *Let R be a Noetherian integrally closed domain and let $\mathfrak{p} \subset R$ is a height 1 prime ideal. Then $R_{\mathfrak{p}}$ is a Noetherian, integrally closed, local domain of Krull dimension 1.*

Proof of Theorem 184. First let R be a DVR. Then R is a PID, so it is Noetherian and integrally closed. Also R is local with prime ideals (0) and \mathfrak{m} , so it has Krull dimension 1.

In the other direction, let $\mathfrak{m} \subset R$ be the maximal ideal. Take $x \in R$ nonzero. Then there exists a unique $n \geq 0$ such that $x \in \mathfrak{m}^n \setminus \mathfrak{m}^{n+1}$ by Krull intersection theorem. We need to show that \mathfrak{m} is principal. Given the claim, take $x \in R \setminus 0$. Then $x \in (\mathfrak{m}^n) \setminus (\mathfrak{m}^{n+1})$ for a unique n . Thus $x = ut^n$ for some unit u , and thus n defines a valuation $k^* \rightarrow \mathbb{Z}$.

To prove the claim, $\mathfrak{m} \neq \mathfrak{m}^2$ by Krull or by Nakayama. Take $t \in \mathfrak{m} \setminus \mathfrak{m}^2$. Since the only prime ideals are $0, \mathfrak{m}$, and every radical ideal is an intersection of primes, $\mathfrak{m} = \sqrt{(t)}$. Thus $\mathfrak{m}^n \subset (t)$ for some $n \geq 1$. Choose n to be the smallest such n , so choose $b \in \mathfrak{m}^{n-1} \setminus (t)$. Take $x = \frac{t}{b} \in K$. Then $b \in (t)$, so $x^{-1} \notin R$. Because R is integrally closed, x^{-1} is not integral over K . This implies that $x^{-1}\mathfrak{m} \not\subset \mathfrak{m}$ (otherwise, choose $\mathfrak{m} = (e_1, \dots, e_s)$ and write $x^{-1}e_i = \sum a_{ij}e_j$. Then $\sum (x^{-1}\delta_{ij} - a_{ij})e_j = 0$, so $\det|x^{-1}\delta_{ij} - a_{ij}| = 0$. Thus x^{-1} is integral).

However, $x^{-1}\mathfrak{m} \subset R$ because if $z \in \mathfrak{m}$ then $bz \in \mathfrak{m}^n \subset (t)$, so we can write $bz = rt$, so $x^{-1}z = r \in R$. We see that $x^{-1}\mathfrak{m} \subset R$, so it is an ideal R but not contained in a maximal ideal. Therefore $x^{-1}\mathfrak{m} = R$, so $\mathfrak{m} = (x)$. \square

Let X be a normal variety. Then we have local rings $R_{\mathfrak{p}} \subset k(X)$ and a DVR for every irreducible subvariety $Y \subset X$ of codimension 1. Thus we have a collection of discrete valuations $v_Y : k(X)^* \rightarrow \mathbb{Z}$. There are called orders of zeroes or poles of $f \in k(X)$ along Y .

5.8 Lecture 20 (Apr 11) Recall that if X is a normal variety and $Y \subset X$ is an irreducible subvariety, then $\mathcal{O}_{Y,X}$ is a DVR. Last time we defined the order of zeroes or poles as the valuation.

Corollary 186. *Let X be normal and $Y \subset X$ be irreducible. Then there exists an affine chart $U \subset X$ with $U \cap Y \neq \emptyset$ such that $I(Y) \subset k[U]$ is principal.*

Proof. Assume that X is affine. Then $\mathfrak{p} = I(Y) = (u_1, \dots, u_m) \subset k[X] = R$. Then we note that $\mathfrak{p}_{\mathfrak{p}} \subset R_{\mathfrak{p}}$ is the maximal ideal in the DVR and it is principal. Then we can write $t = \frac{a}{b}$ where $b \notin \mathfrak{p}$. We can write $u_i = tv_i$, where $v_i \in R_{\mathfrak{p}}$, where $v_i = \frac{a_i}{b_i}$ where $b_i \notin \mathfrak{p}$. Now we know $t \in k[U]$ and $u_i = tv_i$, where $v_i \in k[U]$. Then $I(Y \cap U) = (t) \subset k[U]$. \square

Corollary 187. *The singular locus of a normal variety has codimension at least 2.*

Proof. Suppose there exists an irreducible hypersurface Y in the singular locus. As a variety, Y has an open smooth locus. Thus we can shrink X to an affine chart $U \subset X$ such that $U \cap Y$ is nonempty and smooth. By the previous corollary, we can also assume that X is affine. Then $I(Y) = (t)$. This implies X is smooth along Y by Theorem 153. This gives a contradiction. \square

Corollary 188. *For curves, smooth is equivalent to normal.*

5.8.1 *Normalization* Trying to resolve singularities is not very well resolved, and has led to several Fields Medals. On the other hand, normalization is relatively easy to understand and always available.

Example 189. Consider the curve $X = \{y^2 = x^3\}$. This is singular, so not normal. We see that $t \in k(X)$ is a root of a monic equation $T^2 - x$, but $t \notin k[X]$. We know that the rational parameterization from \mathbb{A}^1 is birational and finite, and the source is normal.

Definition 190. A morphism $\nu : X^\vee \rightarrow X$ is called a normalization if it is birational and finite and X^\vee is normal.

Theorem 191. Every irreducible variety has a normalization, which is unique up to isomorphism.

Construction for affine X. Let $k[X] \subset k(X)$, so define R to be the integral closure of $k[X]$ in $k(X)$. We show that R is a finitely generated $k[X]$ -module. Given the claim, R is also a finitely-generated k -algebra, so $R = k[Y]$ for some variety Y , which must be birational to X because $k[X] \subset k[Y] \subset k(X)$. The extension $k[Y]/k[X]$ is integral, so $Y \rightarrow X$ is finite. Also, we know R is integrally closed, so Y is normal.

To prove the claim, use Noether’s normalization lemma. We have a diagram

$$\begin{array}{ccccc}
 k[X] & \hookrightarrow & R & \hookrightarrow & k(X) \\
 \uparrow & & & & \uparrow \\
 k[T_1, \dots, T_n] & \hookrightarrow & & \hookrightarrow & k(T_1, \dots, T_n)
 \end{array}$$

We know that A is a UFD, so it is integrally closed. Thus A^n is normal, so R is integral over A . This implies that R is the integral closure of A in L . It remains to prove the following lemma. \square

Lemma 192. Let A be an integrally closed Noetherian domain in its field of fractions K . Then if L/K is a finite separable extension and B is the integral closure of A in L , then B is a finitely generated A -module.

Proof. Consider the map $L \rightarrow K$ given by the trace of the map given by multiplication by x . Because L/K is separable, then $\text{Tr}(xy)$ is a nondegenerate quadratic form. It suffices to find a basis v_1, \dots, v_n of L over K such that $B \subset Av_1 + \dots + Av_n$ (this is because A is Noetherian).

To find the basis, first choose some basis u_1, \dots, u_n . After rescaling by elements of A , we can assume that $u_1, \dots, u_n \in B$. Then the u_i are algebraic over K , so it is a root of some polynomial $a_0u_i^{d_i} + \dots + a_{d_i} = 0$. Then we see that $a_0u_i \in B$. Finally, let v_1, \dots, v_n be a dual basis via the trace form. Take $x \in B$ and write it as $x_1v_1 + \dots + x_nv_n$. First note that $\text{Tr}(xu_j) = x_j$. On the other hand, $x \in B$, $u_i \in B$ implies that $xu_i \in B$, so $\text{Tr}(xu_i) \in A$. Therefore $x_i \in A$ for all i . \square

Theorem 193 (Universal Property of Normalization). 1. Suppose $g : Y \rightarrow X$ is a finite birational regular map. Then there exists a unique morphism $f : X^\vee \rightarrow Y$ such that $\nu = g \circ f$.

2. If Y is a normal variety with a dominant regular map $g : Y \rightarrow X$, then there exists a unique morphism $f : Y \rightarrow X^\vee$ such that $g = \nu \circ f$.

Proof. Assume X is affine. In 1, we can also assume Y is affine. Then note that $k(Y) = k(X)$, so we must conclude that $k[Y] \subset k[X^v]$, and this inclusion is unique.

In 2, we can also assume Y is affine. Then we again know that $k[X] \subset k[Y]$ and $k[X] \subset k[X^v]$ and that $k(X) \subset k(Y)$. We know $k[Y]$ is integrally closed, so $k[Y] \supset k[X^v]$ and this containment is unique. \square

5.9 Lecture 21 (Apr 16) Homework is now due next Tuesday as opposed to this Thursday. There will be one more homework and then the second take-home midterm.

6 DIVISORS

Let X be an irreducible variety.

Definition 194 (Prime Divisor). A prime divisor $D \subset X$ is an irreducible subvariety of codimension 1 (hypersurface).

Definition 195 (Divisor Group). The divisor group $\text{Div } X$ is a free abelian group generated by symbols $[D]$ for each prime divisor $D \subset X$.

Definition 196. A divisor $D = \sum n_i H_i$ is effective, or $D \geq 0$, if $n_i \geq 0$ for all i .

Suppose X is normal (or at least non-singular in codimension 1) and let $H \subset X$ a prime divisor. Then $\mathcal{O}_{H,X} \subset k(X)$ is a DVR with valuation $v_H : k(X)^* \rightarrow \mathbb{Z}$. If $v_H(f) > 0$ then we say that f has a zero of order $v_H(f)$ along H . If $v_H(f) < 0$, then f has a pole. To every $f \in k(X)^*$ we associate a divisor $(f) = \sum_{H \subset X} v_H(f)[H]$ called the divisor of zeros and poles.

Lemma 197. $v_H(f) = 0$ for all but finitely many prime divisors.

Proof. We can shrink X as much as we want because if $U \subset X$ is open, then $X \setminus U$ contains only finitely many hypersurfaces. We can also assume that X is affine and that $f \in k[X]$. We can also remove $V(f)$ and assume that f is invertible. Then it is easy to see that $(f) = 0$. Indeed, f is invertible on every hypersurface, so its valuation must be 0. \square

Remark 198. Divisors of the form (f) are called principal divisors. Principal divisors form a subgroup in $\text{Div } X$ because the valuation is like a logarithm (takes products to sums).

Example 199. If X is normal and projective and $f \in k(X)$, $(f) \geq 0$ if and only if f is constant.

Definition 200. The divisor class group $\text{Cl } X := \text{Div } X / \{(f)\}$ is the quotient of the divisor group by the subgroup of principal divisors.

Example 201. Take $X = \mathbb{A}^n$. Then a prime divisor $H \subset X$ is an irreducible hypersurface given by an irreducible polynomial $F \in k[x_1, \dots, x_n]$. We claim that $\text{div}(F) = H$. In particular, all divisors are principal, so $\text{Cl}(\mathbb{A}^n) = 0$. To see the claim, we compute $v_H(F) = 1$ and then for any other hypersurface, $v_{H'}(F) = 0$.

Remark 202. Suppose X is normal, $f \in k(X)$ and suppose $(f) \geq 0$. Then f is regular outside of that divisor and at general points of irreducible components of (f) , so f is not regular only in codimension 2. Therefore f is regular everywhere.

Proposition 203 (Homework). *Let X be an affine variety. Then the class group of X is trivial if and only if $k[X]$ is a UFD.*

Example 204. Let $X = \mathbb{P}^n$. Then $H \subset X$ irreducible hypersurface is given by an irreducible homogeneous polynomial $F \in k[T_0, \dots, T_n]$ of degree k . However, $F \notin k(\mathbb{P}^n)$. However, we know $I(H \cap \mathbb{A}_i^n) = \left(\frac{F}{T_i^k} \right)$. Then we know $v_{\tilde{H}}(f)$ is 1 when $\tilde{H} = H$, 0 when $\tilde{H} \neq H$ and $\tilde{H} \cap \mathbb{A}_i^n \neq \emptyset$, and unknown when \tilde{H} is the hyperplane at infinity.

To compute this, choose a different affine chart which gives $-k$. Now take any $f \in k(\mathbb{P}^n)$. We know $f = \frac{\prod F_i^{n_i}}{\prod G_j^{m_j}}$, so we can write $(f) = \sum n_i [F_i = 0] - \sum m_j [G_j = 0]$.

Corollary 205. $D \in \text{Div } \mathbb{P}^n$ is principal if and only if $\deg(D) = 0$, where $\text{def}(D) = \sum n_i \deg(F_i)$, where $F_i \in k[T_0, \dots, T_n]$ is a homogeneous equation of H_i . Therefore the class group of \mathbb{P}^n is \mathbb{Z} .

Definition 206. Divisors D, D' on X are called linearly equivalent ($D \sim D'$) if $D - D'$ is a principal divisor.

Remark 207. Every divisor on \mathbb{P}^n is linearly equivalent to a unique multiple of a hyperplane.

The notion of a divisor we just introduced is called a Weil divisor.

Definition 208 (Cartier Divisor). A Cartier divisor on an irreducible variety X is the following data:

1. A cover $X = \cup U_i$ by open sets;
2. Rational functions $f_i \in k(X)^*$ "defining a divisor on U_i " satisfying a compatibility condition: f_i/f_j is invertible on $U_i \cap U_j$.

Two such data (U_i, f_i) and (V_j, g_j) are considered equivalent if f_i/g_j is an invertible regular function on $U_i \cap V_j$.

Now suppose X is normal, or at least nonsingular in codimension 1. Then to each Cartier divisor (U_i, f_i) we associate a Weil divisor F such that $D \cap U_i = (f_i) \cap U_i$. Concretely, $D = \sum n_j H_j$, where $n_j = v_{H_j}(f_i)$ as long as $H_j \cap U_i \neq \emptyset$.

Example 209. Take a prime divisor $H \subset \mathbb{P}^n$ and take the standard cover $\mathbb{P}^n = \cup U_i$. We see that $I(H \cap U_i) = (f_i)$ where $f_i \in k[x_0, \dots, x_n]$. Thus H is in fact given by a Cartier divisor (U_i, f_i) .

Why are the f_i compatible? If H is given by homogeneous F of degree k , then $f_i = F/T_i^k$. Then $f_i/f_j = (T_j/T_i)^k$, which is regular and invertible on $U_i \cap U_j$.

We can turn CDiv , the group of Cartier divisors into a group $(U_i, f_i) \cdots (V_j, g_j) = (U_i \cap V_j, f_i g_j)$. By the above construction, we have a homomorphism $\text{CDiv} \rightarrow \text{Div}$. If $f \in k(X)^*$, there is a Cartier divisor (X, f) , so principal divisors lie in the Cartier divisors. If X is normal, this is an inclusion.

Definition 210 (Picard Group). The Picard group $\text{Pic } X := \text{CDiv} / \{(f)\}$ is the quotient of the group of Cartier divisors by principal divisors.

Note that the Picard group is contained in the class group if X is normal. We want to determine when every Weil divisor is Cartier. If X is nonsingular (more generally, $\mathcal{O}_{x,X}$ is a UFD) for all $x \in X$, choose $H \subset X$ a prime Weil divisor. We need to construct an open cover: first take

$U_0 = X \setminus H, f_0 = 1$. For all $x \in X$, there exists some affine neighborhood $U \supset X$, which depends on $x \in X$, such that H is principal in this neighborhood. This gives an infinite open cover, but we can choose a finite subcover. Thus H is a Cartier divisor.

Corollary 211. *If X is locally factorial, every divisor is Cartier.*

6.1 Lecture 22 (Apr 18) We will continue our discussion of divisors. Here we will assume that X is nonsingular and study the behavior of divisors under regular maps.

Let $D = (U_\alpha, f_\alpha)$ where $Y = \cup U_\alpha, f_\alpha \in k(Y)^*$. Then we define $\varphi^*D = (\varphi^{-1}U_\alpha, \varphi^*f_\alpha)$. We need to see that the pullback of each f_α is well defined and nonzero. This happens with $\overline{\varphi(X)}$ is not contained in the divisor of zeroes and poles of f_α , so when $\overline{\varphi(X)} \not\subseteq D$. Fortunately, we can pass to a linearly equivalent divisor.

Lemma 212. *Let X be nonsingular. Then for any divisor $D \subset X$ there exists $D' \sim D$ such that $x_1, \dots, x_m \notin D'$.*

Proof. We argue by induction on m . We can assume that $x_1, \dots, x_{m-1} \notin D, x_m \in D$. We can also assume that D is prime and assume that X is affine. Then D has a local equation at x_m : $(\pi) = \mathcal{L}_D \subset \mathcal{O}_{x_m}$, where $\pi = p/q$ for some $p, q \in k[X]$. Near x_m , we see that $D = (p)$. Then $D' = D - (p) \sim D$ does not contain x_m , but it may contain x_1, \dots, x_{m-1} .

Choose $g_i \in k[X]$ such that $g_i(x_i) \neq 0$. Then for all $i = 1, \dots, m-1, g_i|_{D \cup \{x_1, \dots, x_m\}} = 0$. Then choose constants $\alpha_1, \dots, \alpha_{m-1}$ such that for $f = p + \sum \alpha_i g_i^2, f(x_i) \neq 0$. We claim that $D' = D - (f)$ does not contain x_1, \dots, x_m . We know that $f(x_i) \neq 0$, so $x_1, \dots, x_{m-1} \notin (f)$. Now we need to know that f is a local equation of D . Then because $f = \sum \alpha_i g_i^2$, we know that $p|g_i$, so $f = p(1 + \sum \alpha_i p w_i^2)$, so it is also a local equation of D . □

This defines a pullback on the level of Picard groups.

6.1.1 Divisors of rational maps to \mathbb{P}^n Suppose $\varphi : X \dashrightarrow \mathbb{P}^n$ is a rational map defined by $(\varphi_0 : \dots : \varphi_n)$. Suppose know $(\varphi) = D_i - D$ where D_0, \dots, D_n, D are effective. Choose D to be the smallest possible.

We want to know where φ is regular. Choose $x \in X$. Then $\varphi_i = p_i/q$, where p_i is a local equation of D_i and q is a local equation of D . Thus $\varphi = (p_0 : \dots : p_n)$. If $p_i(x) \neq 0$ for some x , φ is regular at p . The converse is also true because \mathcal{O}_x is a UFD.

Theorem 213. *The indeterminacy locus of φ is $D_0 \cap \dots \cap D_n$, where $D_i = (p_i = 0)$.*

Example 214. Consider the rational normal curve $\mathbb{P}^1 \rightarrow \mathbb{P}^n$. Then $\varphi_i = x^i$ and $(x^i) = i[0] - i[\infty]$. In another chart, $x = \frac{1}{y}$. Then $D = n[\infty]$, so $D_i = i[0] + (n-i)[\infty]$ and then $D_0 \cap \dots \cap D_n = \emptyset$, so the map is indeed regular everywhere.

Suppose X is a nonsingular variety and D is a divisor. Then set $\mathcal{L}(D) = \{f \in k(X) \mid (f) + D \geq 0\}$ the complete linear system.

Theorem 215 (Serre). *If X is projective, $\mathcal{L}(D)$ has dimension $\ell(D) < \infty$.*

We can define $\varphi_D : X \dashrightarrow \mathbb{P}^{\ell(D)-1}$. First choose a basis $\varphi_1, \dots, \varphi_{\ell(D)}$ and then set $\varphi_D = [\varphi_1 : \dots : \varphi_{\ell(D)}]$.

Lemma 216. *If $D \sim D'$ then $\mathcal{L}(D) \simeq \mathcal{L}(D')$ and $\varphi_D = \varphi_{D'}$, so we have distinguished maps to projective spaces parameterized by the Picard group.*

Proof. If $D \sim D'$ then $D = D' + (f)$. Therefore there are isomorphisms between the linear systems given by multiplying and dividing by f , so they are isomorphic. Then if $\varphi_1, \dots, \varphi_{\ell(D)}$ is a basis of $\mathcal{L}(D)$, $f\varphi_1, \dots, f\varphi_{\ell(D)}$ is a basis of $\mathcal{L}(D')$. Then the two maps are the same. \square

Remark 217. If X is projective then if $(f) + D = (g) + D$, we must have $(f) = (g)$, so $(f/g) = 0$ and thus f/g is a constant.

If X is projective, then $\mathbb{P}^{\text{ell}(D)-1} = \mathbb{P}(\mathcal{L}(D)) = |D|$, which is defined to be the set of effective divisors linearly equivalent to D .

Example 218. If $X = \mathbb{P}^r$, then $\text{Pic } X = \text{Cl } X = \mathbb{Z} = \mathbb{Z}[H]$, where H is a hyperplane. Then $|dH|$ is the set of hypersurfaces of degree d , which is $\mathbb{P}^{\binom{r+d}{d}-1}$. We see that $\mathcal{L}(dH) = \mathcal{L}(dH_0) = \{f \in k(\mathbb{P}^r) \mid (f) + dH_0 \geq 0\}$. Note that $\mathbb{P}^r \setminus H_0 = \mathbb{A}_{x_1, \dots, x_r}^r$. Then we have that all poles of f are at infinity. Thus $f \in k[x_1, \dots, x_r]$, so $v_{H_0}(f) = -\deg(f)$. After homogenizing, this becomes the set of degree d hypersurfaces, so the map given by the divisor is the Veronese embedding.

6.2 Lecture 23 (Apr 23) We continue with our discussion of linear systems of divisors. We will always assume that X is projective and nonsingular. Suppose $\varphi : X \dashrightarrow \mathbb{P}^r$ is a rational map.

Remark 219. Given $\varphi : X \rightarrow \mathbb{P}^r$, it can be degenerate, say $\varphi(X) \subset \mathbb{P}^{r-1} \subset \mathbb{P}^r$. In this case, $\varphi_0, \dots, \varphi_r$ are linearly dependent. This does not happen for φ_D , so we can assume that $\varphi : X \dashrightarrow \mathbb{P}^r$ is not degenerate.

For a basis f_0, \dots, f_r of $\mathcal{L}(D)$,

1. We can assume that D is effective because $\varphi_D = \varphi_{D'}$ for all $D \sim D'$.
2. If D is effective, $(f_i) = D_i - D$. However, it can happen that the D_i have a common prime divisor.

Definition 220 (Fixed Part). A fixed part of a linear system $\mathcal{L}(D)$ is the largest effective divisor F such that $D_i - F \geq 0$. We can write $D_i = F + M_i$, where M_i is the moving part of the linear system.

We can write $\text{abs } D = \{D' \sim D \mid D' \geq 0\} = \mathbb{P}(\mathcal{L}(D))$. Another way to phrase this is to introduce the base locus.

Definition 221 (Base Locus). The base locus BL_D of a divisor D is defined by $\text{BL}_D = \bigcap_{D' \in |D|} D' \supset F$.

We know that $\mathcal{L}(D) \simeq \mathcal{L}(D - F)$ and that $\varphi_D = \varphi_{D-F}$. Thus we can always reduce to linear systems without fixed part.

Definition 222 (Base-point-free). A divisor D is base-point-free if its base locus is empty. Note this implies φ_D is regular.

Definition 223 (Very Ample). D is very ample if it is base-point-free and φ_D is an embedding.

Definition 224 (Ample). D is called ample if kD is very ample for some $k > 0$.

When we start with $\varphi : X \dashrightarrow \mathbb{P}^r$, $\varphi_0, \dots, \varphi_r \in \mathcal{L}(D)$. Now we consider incomplete linear systems: a subspace $V \subset \mathcal{L}(D) \subset k(X)$. Choose a basis f_0, \dots, f_s of V . Then $\varphi_V = [f_0 : \dots : f_s] : X \dashrightarrow \mathbb{P}^s$. We can complete f_0, \dots, f_s to a basis f_0, \dots, f_r of $\mathcal{L}(D)$ and recover φ_V as the composition of φ_D and the projection.

We can talk about the base locus, fixed part of V , define global generation, and very ampleness of V .

Example 225. Consider \mathbb{P}^2 with coordinates x, y, z and let $H = (Z = 0)$. Then $\mathcal{L}(2H) = \{f \in k(\mathbb{P}^2) \mid (f) + 2H \geq 0\} = \{f \in k[x, y] \mid \deg f \leq 2\} = k[x, y, z]_2$. Thus $|2H|$ is the set of conics in \mathbb{P}^2 and $\varphi_{2H} = [z^2 : xz : yz : x^2 : xy : y^2]$, the Veronese embedding. Thus $2H$ is very ample.

Now we consider an incomplete linear system. Choose points $p = [1 : 0 : 0]$, $q = [0 : 1 : 0]$. Consider V the set of conics passing through a, b . Then $\mathcal{L}(2H) = k^6$ and $\dim V \geq 4$. In fact, V is spanned by $1, x, y, xy$. Then $\varphi_V : \mathbb{P}^2 \dashrightarrow \mathbb{P}^3$ is given by $[x : y : 1] \mapsto [1 : x : y : xy]$. In particular, φ_V is birational. The equation of the image is actually $AD - BC = 0$, which makes φ_V a birational map $\mathbb{P}^2 \dashrightarrow Q$ to a quadric surface in \mathbb{P}^3 . We see that the base locus is p, q .

We consider the image of H . For any point $[x : y : 0]$, the image is $[0 : 0 : 0 : xy] = [0 : 0 : 0 : 1]$. Thus H collapses to the point $[0 : 0 : 0 : 1]$. In the hyperplane at infinity $Q \cap (A = 0)$ is given by $BC = 0$, which is a union of two lines. We have two points where φ_V is not regular, so we resolve by blowing up at p, q . We blowup at p and consider two charts of the local blowup.

In the chart $x = 1$, we see $y = u, z = uv$, so $v = z/y$. Thus the chart $u = 0$ is exceptional. In this chart, the map is $[z^2 : z : yz : y] = [u^2v^2 : uv : u^2v : u] = [uv^2 : v : uv : 1]$. We see this is regular. Note v is a coordinate along \mathbb{P}^1 , so restricting to the exceptional divisor E , we get the map $[0 : v : 0 : 1]$, which maps E onto a line $A = C = 0$. The exceptional divisor over q will map to another line $A = B = 0$.

Thus φ_V is given by blowing up p, q and then contracting the proper transform of H .

Example 226. We will do similar calculations without being able to do it too explicitly. Choose 6 points p_1, \dots, p_6 on \mathbb{P}^2 that are distinct, in general linear position, and not coconic. Consider the linear system $\mathcal{L}(3H)$ and φ_{3H} . Note $|3H|$ is the space of cubics in \mathbb{P}^2 , so consider V to be the set of cubics through p_1, \dots, p_6 . We check that the dimension of V is 4. If this is not the case, then every cubic that passes through p_2, \dots, p_6 also passes through p_1 . This is false because we can take a conic through p_1, \dots, p_6 and add a line not through p_1 . Thus we get a rational map $\varphi_V : \mathbb{P}^2 \dashrightarrow \mathbb{P}^3$.

First we find the base locus, which is $\{p_1, \dots, p_6\}$. For any other point q , we can take the conic through five of the points and a line containing p_6 but not q . If q is on the conic, just change the five points and then note that q cannot be on the new conic because both conics are smooth and they already have four intersections points.

6.3 Lecture 24 (Apr 25) Last time we were building up to a theorem of Clebsch:

Theorem 227 (Clebsch). *For a smooth projective surface X , the following are equivalent:*

1. $X \simeq \text{Bl}_6\mathbb{P}^2$ where the six points are in general position (no three are collinear and the six points are not coconic).
2. X is a smooth cubic surface in \mathbb{P}^3 .
3. X is a del Pezzo surface of degree 3 (Fano variety of dimension 2, which means the anticanonical divisor is ample).

We continue Example 226 from last time.

Example (Continuation of Example 226). We show that for all $p, q \in \mathbb{P}^2 \setminus \{p_1, \dots, p_6\}$, $\varphi_V(p) \neq \varphi_V(q)$. To see this, take the conic through 1, 2, 3, 4, 5 and the line through 6, p and a similar combinatorial argument from the base locus calculation tells us that at least one choice works.

Next we show that the induced rational map $\text{Bl}_6\mathbb{P}^2 \rightarrow \mathbb{P}^3$ is regular. It suffices to show that it is regular at every point of $E \simeq \mathbb{P}^1$ over p_1 . To do this, we find two cubics $D_1, D_2 \in V$ transversal at p_1 . To do this, we take the conic C through p_2, \dots, p_6 and then two distinct lines L_1, L_2 through p_1 . Then $D_1 = C \cup L_1, D_2 = C \cup L_2$. To see that this is enough, choose a basis f_1, \dots, f_4 of V such that $D_1 = (f_1 = 0)$ and $D_2 = (f_2 = 0)$. Assume $p_1 = [0 : 0 : 1]$. We know φ_V is given by $[f_1 : f_2 : f_3 : f_4] = [f_1 : z^3 : \dots : f_4/z^3]$. We can dehomogenize in the standard chart and write $\varphi_V = [y + \alpha : x + \beta : g_3 : g_4]$, where $\alpha, \beta \in \mathfrak{m}_{(0,0)}^2$ and $g_3(0, 0) = g_4(0, 0) = 0$.

Next we blow up and consider the chart $x = u, y = uv$. Then $\varphi_V = [uv + u^2g_1 : u + u^2g_2 : ug_3 : ug_4] = [v + ug_1 : 1 + ug_2 : g_3 : g_4]$. The equation of E_1 is $u = 0$, so we substitute $u = 0$ and in the second component we have 1, so φ_V is regular at every point of E_1 .

Remark 228. We consider π^*V where $\pi : \text{Bl}_6\mathbb{P}^2 \rightarrow \mathbb{P}^2$. Then for all $D \in V$, we see $\pi^*D = E_1 + \dots + E_6 + D'$, so $\pi^*V = (E_1 + \dots + E_6) + |D'|$. Thus $D' \sim \pi^*(3H) - E_1 - \dots - E_6$. The D' are the preimages of hyperplanes in \mathbb{P}^3 . Because $\varphi(E_i)$ are not points (it intersects a general hyperplane at one point), it must be a curve. In fact it must be a line because a general hyperplane will give a cubic in \mathbb{P}^2 smooth at p_i , so its proper transform intersects E_i in one point.

Next we will show that $\varphi(\text{Bl}_6\mathbb{P}^2)$ is smooth. Indeed, if not, T_x has dimension 3 for some $x \in X$. (To be continued next lecture).

7 REGULAR DIFFERENTIAL FORMS

Now we will attempt to perform calculus on algebraic varieties. Let X be a variety and $f \in k[X]$. Then $d_x f$ is a linear function on T_x .

Definition 229. $\Phi[X] = \{ \text{functions that assign every } x \in X \text{ an element of } T_x^\vee \}$. Note that $\Phi[X]$ is a $k[X]$ -module. For $f \in k[X], \alpha \in \Phi[X], (f\alpha)(x) = f(x)\alpha(x)$.

Definition 230. The regular differential forms $\Omega[X] \subset \Phi[X]$ are $\alpha \in \Phi[X]$ such that for every $x \in X$, there exists a neighborhood $U \ni x$ such that $\alpha|_U = \sum_i f_i dg_i$. We have a map $d : k[X] \rightarrow \Omega[X]$ and the Leibniz rule $d(fg) = f(dg) + g(df)$.

Lemma 231. *If X is affine, then $\Omega[X]$ is generated by differentials.*

Proof. We use another algebraic partition of unity. For all $x \in X$, we can write $\alpha = \sum f_{i,x} dg_{i,x}$ is some neighborhood of x . Clearing denominators, $p_x \alpha = \sum_i r_{i,x} dg_i, x$ where $p_x, r_{i,x}, g_{i,x} \in k[X]$

and $p_x(x) \neq 0$. Then the ideal $(p_x) = k[X]$ by the Nullstellensatz, so we can choose finitely many of them and write $\sum p_x q_x = 1$. Then multiply by q_x and add, and we get $\alpha = \sum_{i,x} r_{i,x} q_x dg_{i,x}$. \square

Corollary 232. *Let X be affine and g_1, \dots, g_s generate $k[X]$. Then dg_1, \dots, dg_s generate $\Omega[X]$ as a $k[X]$ -module (use the Leibniz rule to prove this).*

In the simplest case, $\Omega[\mathbb{A}^n]$ is a free $k[x_1, \dots, x_n]$ -module with generators dx_1, \dots, dx_n (if $\sum f_i dx_i = 0$ then $\exists x \in \mathbb{A}^n$ where one $f_i \neq 0$, and then dx_1, \dots, dx_n are linearly dependent in $T_x \mathbb{A}^3$, which is a contradiction).

Theorem 233. *Suppose $x \in X$ is a nonsingular point with local parameters u_1, \dots, u_n . Then there exists an affine neighborhood U of x such that $\Omega[U]$ is a free $k[U]$ -module generated by du_1, \dots, du_n .*

Proof. Without loss of generality, $x \in \mathbb{A}^N$. Then $I(X) = (F_1, \dots, F_m)$, so $k[X]$ is generated by $f_i = T_i|_X$, so $\Omega[X]$ is generated by dt_1, \dots, dt_N . We know $F_i|_X = 0$, so $\sum_j \frac{\partial F_i}{\partial T_j} dt_j = 0$ for all i . The Jacobian matrix has rank $N - n$ at all nonsingular $x \in X$, so it has a nondegenerate minor with the last variables.

Thus t_1, \dots, t_n are local parameters of x . From We get that $dt_j = \sum_{i=1}^n f_j^i dt_i$ for all $j > n$ using Cramer's rule, where $f_j^i \in \mathcal{O}_x$. Now we shrink X to an affine neighborhood U such that all $f_j^i \in k[U]$. After this shrinking, $\Omega[U]$ is generated by dt_1, \dots, dt_n and t_1, \dots, t_n are local parameters for all $y \in U$. \square

7.1 Lecture 25 (Apr 30) There will be extra office hours tomorrow from 2 to 4.

We will continue our study of multivariable calculus. Last time we discussed regular differential forms.

Example 234. We calculate $\Omega^1[\mathbb{P}^1]$. We know $\Omega^1[\mathbb{A}^1] = k[x]dx$. Take another chart with coordinate $y = 1/x$, we can write $\omega = f(x)dx = f\left(\frac{1}{y}\right) d\frac{1}{y} = \frac{g(y)}{y^n} \left(-\frac{dy}{y^2}\right) = \frac{g(y)dy}{y^{n+2}}$ where $g(y) \neq 0$. This is not a polynomial multiple of dy , so $\Omega^1[\mathbb{P}^1] = 0$.

Now we talk about differential r -forms. We introduce $\Phi^r[X] = \{x \in X \mapsto \omega(x) \in \wedge^r T_x^*\}$ and $\Omega^r[X]$ as the subspace that can be written locally as $\omega = \sum f_{i_1, \dots, i_r} dg_{i_1} \wedge \dots \wedge dg_{i_r}$.

Theorem 235. *Let X be nonsingular at x and $U \ni x$ an affine neighborhood with a system of local parameters u_1, \dots, u_n . Then $\Omega^r[U]$ is a free $k[U]$ -module generated by $\binom{n}{r}$ forms of the form $du_{i_1} \wedge \dots \wedge du_{i_r}$.*

Proof of this is the same as for Ω^1 . Now we discuss the case of $\Omega^n[X]$ the space of canonical differentials. Here we have $\Omega^n[U] = k[U]du_1 \wedge \dots \wedge du_n$. If u'_1, \dots, u'_n is another system of local parameters, then $du_1 \wedge \dots \wedge du_n = J(u_1, \dots, u_n, u'_1, \dots, u'_n) du'_1 \wedge \dots \wedge du'_n$ where J is an invertible function.

We discuss rational canonical differentials $\Omega^n(X)$ where every pair is a form (U, ω) where $U \subset X$ is open and $\omega \in \Omega^n[U]$. They are subject to equivalence by equality on intersections. We can assume from now on that X is nonsingular and $\omega \in \Omega^n(X)$ is regular on some affine open with local parameters u_1, \dots, u_n , so $\omega = f(x)du_1 \wedge \dots \wedge du_n$ where $f \in k(X)$ is regular on U .

Lemma 236. $\Omega^n(X)$ is a one-dimensional vector space over $k(X)$ generated by du_1, \dots, du_n for all choices of local parameters somewhere.

Definition 237. Choose $\omega \in \Omega^n(X)^*$. We will define (ω) as a Cartier divisor. Cover $X = \cup U_\alpha$ such that for all alpha there exist $u_1^\alpha, \dots, u_n^\alpha$ are local parameters on U_α . Then $\omega|_{U_\alpha} = f^\alpha du_1^\alpha \wedge \dots \wedge du_n^\alpha$ for $f^\alpha \in k(X)$. This gives a Cartier divisor (U_α, f_α) .

To see this is Cartier, on the overlaps $U_\alpha \cap U_\beta$ we see that $\omega|_{U_\alpha \cap U_\beta} = f^\alpha du_1^\alpha \wedge \dots \wedge du_n^\alpha = f^{\alpha'} du_1^{\alpha'} \wedge \dots \wedge du_n^{\alpha'} = f^\alpha \mathcal{J} du_1^{\alpha'} \wedge \dots \wedge du_n^{\alpha'}$, so $f^\alpha = \mathcal{J} f^{\alpha'}$, where the Jacobian is invertible.

Note that if $\omega' = f\omega$ then $(\omega') = (f) + (\omega)$, so they are equivalent. Thus the class of K_X in the Picard group only depends on X .

Example 238. We calculate the canonical divisor of \mathbb{P}^n . Let $\mathbb{A}^n = \{z_0 \neq 0\}$ and choose $\omega = dx_1 \wedge \dots \wedge dx_n$. This has no zeroes or poles on \mathbb{A}^1 and let $H = \mathbb{P}^n \setminus \mathbb{A}^n$ be the plane at infinity. We see that $[1 : x_1 : \dots : x_n] = [1/x_1 : 1 : x_2/x_1 : \dots : x_n/x_1]$, so writing $y_1 = 1/x_1, x_2 = y_2, \dots, x_n = y_n/y_1$, we have

$$\begin{aligned} \omega &= d(1/y_1) \wedge d(y_2/y_1) \wedge \dots \wedge d(y_n/y_1) \\ &= -\frac{dy_1}{y_1^2} \wedge \frac{(dy_2)y_1 - y_2 dy_1}{y_1^2} \wedge \dots \wedge \frac{y_1 dy_n - y_n dy_1}{y_1^2} \\ &= \pm \frac{dy_1}{y_1^2} \wedge \frac{dy_2}{y_1} \wedge \dots \wedge \frac{dy_n}{y_1} \\ &= \pm \frac{1}{y_1^{n+1}} dy_1 \wedge \dots \wedge dy_n \end{aligned}$$

Thus $K_{\mathbb{P}^n} = -(n+1)H$.

Definition 239. X is Fano is $-K_X$ is ample.

Corollary 240. \mathbb{P}^n is Fano. $\varphi_{-K_X} = \varphi_{(n+1)H}$ is the $(n+1)$ -th Veronese embedding of \mathbb{P}^n .

For example, $-K_{\mathbb{P}^2} = 3H$. and $|-K_{\mathbb{P}^2}|$ is the set of cubic curves in \mathbb{P}^2 . We now determine the relationship between K_S and K_X if S is a blowup of X at a point p . Choose local parameters u, v at p and $\omega = du \wedge dv$ on X . Then (ω) is disjoint from p , and we know $k(S) = k(X)$, so $\omega \in \Omega^n(S)$. We compute its divisor on S .

Working in a chart of the blowup with local parameters x, y where $u = x, v = xy$, we have $\omega = du \wedge dv = dx \wedge (xdy + ydx) = xdx \wedge dy$. Near E , (ω) is precisely E . This proves the following:

Theorem 241. Let $S \xrightarrow{\pi} X$ be the blowdown. Then $K_S = \pi^* K_X + E$, where E is the exceptional divisor.

Corollary 242. Let S be a blowup of \mathbb{P}^2 at 6 points. Then $K_S = \pi^* K_{\mathbb{P}^2} + E_1 + \dots + E_6 = -3H + E_1 + \dots + E_6$ and thus $-K_S = 3H - E_1 - \dots - E_6$.

Last time, we constructed the map $\varphi : S \rightarrow \mathbb{P}^3$. We have the following:

Theorem 243. φ is an isomorphism onto its image, which is a smooth cubic surface.

Proof. We already know that φ is birational and bijective away from E_1, \dots, E_6 . Also, $\varphi|_{E+i}$ is a line in \mathbb{P}^3 . Therefore φ has finite fibers. We claim that φ is finite. Instead of a proof of the claim, we state the Stein Factorization Theorem (after the end of this proof).

Given Stein, φ is finite and birational, so we need to check that Y is normal. Note X is smooth, so it is normal. We use Serre's criterion. If X is a hypersurface in \mathbb{P}^n then X is Cohen-Macaulay and is thus S_1 by Hartog's principle. Now we need to show that $X = \varphi(S) \subset \mathbb{P}^3$ has isolated singularities. If not, then X is singular along $C \subset X$, so every plane section $X \cap H$ is singular at $C \cap H$. Thus $X \cap H$ is a singular cubic curve, which means it is rational. However, on S , $X \cap H$ is a member of $|3H| - p_1 - \dots - p_6$, which for some H is a smooth cubic curve, which is not rational. \square

Corollary 244. $-K_S$ is very ample, so S is a Fano surface, or a del Pezzo surface.

Theorem 245 (Stein Factorization Theorem). *Let $f : X \rightarrow Y$ be a proper regular map. Then there exists a commutative diagram*

$$\begin{array}{ccc} X & \xrightarrow{\quad} & Y \\ & \searrow & \nearrow \\ & Z & \end{array}$$

where $X \rightarrow Z$ has connected fibers and $Z \rightarrow Y$ is finite. In particular, if f is proper and has finite fibers, then f is finite.